

Abbreviations Table and References

In this guide

[In this guide](#)

1. [Fifth draft statement on the safety of Titanium Dioxide \(E171\) as a Food Additive- Introduction](#)
2. [Fifth draft statement on the safety of Titanium Dioxide \(E171\) as a Food Additive- Executive Summary](#)
3. [Fifth draft statement on the safety of Titanium Dioxide \(E171\) as a Food Additive- Exposure Assessment](#)
4. [Fifth draft statement on the safety of Titanium Dioxide \(E171\) as a Food Additive- Methodology of the COT review](#)
5. [Fifth draft statement on the safety of Titanium Dioxide \(E171\) as a Food Additive- Physicochemical Characterisation of nano grade TiO₂](#)
6. [Fifth draft statement on the safety of Titanium Dioxide \(E171\) as a Food Additive- Studies used to review the toxicokinetics and absorption of the E171 form of TiO₂](#)
7. [Fifth draft statement on the safety of Titanium Dioxide \(E171\) as a Food Additive- EFSA review and conclusions on ADME of TiO₂](#)
8. [Fifth draft statement on the safety of Titanium Dioxide \(E171\) as a Food Additive- Summary of the EOGRT study \(LPT, 2020\)](#)
9. [Fifth draft statement on the safety of Titanium Dioxide \(E171\) as a Food Additive- Results](#)
10. [Fifth draft statement on the safety of Titanium Dioxide \(E171\) as a Food Additive- Studies using the E171 form of TiO₂ \(in mice\)](#)
11. [Fifth draft statement on the safety of Titanium Dioxide \(E171\) as a Food Additive- COM review and conclusions](#)
12. [Fifth draft statement on the safety of Titanium Dioxide \(E171\) as a Food Additive- Reproductive and developmental studies using the nanoparticle form of TiO₂](#)
13. [Fifth draft statement on the safety of Titanium Dioxide \(E171\) as a Food Additive- Neurotoxicity](#)
14. [Fifth draft statement on the safety of Titanium Dioxide \(E171\) as a Food Additive- Exposure Estimation](#)

15. [Fifth draft statement on the safety of Titanium Dioxide \(E171\) as a Food Additive- Abbreviations Table and References](#)
16. [Fifth draft statement on the safety of Titanium Dioxide \(E171\) as a Food Additive- Annex B](#)
17. [Fifth draft statement on the safety of Titanium Dioxide \(E171\) as a Food Additive- Annex C](#)
18. [Fifth draft statement on the safety of Titanium Dioxide \(E171\) as a Food Additive- Annex D](#)
19. [Fifth draft statement on the safety of Titanium Dioxide \(E171\) as a Food Additive- Annex E](#)

This is a draft statement for discussion. This does not represent the views of the Committee and should not be cited.

Abbreviations Table

ABC	Aberrant Crypts
ACF	Aberrant Crypt Foci
COM	Committee on Mutagenicity
COT	Committee on Toxicity
E171	Titanium Dioxide (food additive)
EFSA	European Food Safety Authority
EOGRT	Extended One-Generation Reproductive Toxicity (study)
HBGV	Health-based Guidance Value
OECD	Organisation for Economic Co-Operation and Development (Guideline 443)

POD	Point of Delivery
TiO ₂	Titanium Dioxide
TiO ₂ -P25	Anatase:Rutile : 85:15, Diameter: 10-40nm
BET	Brunauer-Emmett-Teller

References

Akagi JI, Mizuta Y, Akane H, Toyoda T, Ogawa K. (2023) Oral toxicological study of titanium dioxide nanoparticles with a crystallite diameter of 6 nm in rats. Part Fibre Toxicol. 20;20(1):23.

Ammendolia MG, Iosi F, Maranghi F, Tassinari R, Cubadda F, Aureli F, Raggi A, Superti F, Mantovani A and De Berardis B, (2017) Short-term oral exposure to low doses of nano-sized TiO₂ and potential modulatory effects on intestinal cells. Food and Chemical Toxicology. 102 pp 63-75.

ANSES (Agence nationale de sécurité sanitaire de l'alimentation, de l'environnement et du travail) (2017) Expert Committee (commissioned by ANSES.). www.nanoreg.eu

Bachler, G., von Goetz, N., & Hungerbuhler, K. (2015). Using physiologically based pharmacokinetic (PBPK) modeling for dietary risk assessment of titanium dioxide (TiO₂) nanoparticles. Nanotoxicology, 9(3) pp 373-380.

Balcaen, L., Bolea-Fernandez, E., Resano, M. and Vanhaecke, F. (2014) Accurate determination of ultra-trace levels of Ti in blood serum using ICP-MS/MS. Analytica chimica acta, 809, pp.1-8.

Bates, B.; Collins, D.; Jones, K.; Page, P.; Roberts, C.; Steer, T.; Swan, G. (2020) National Diet and Nutrition Survey Results from years 9, 10 and 11 (combined) of the Rolling Programme (2016/2017 to 2018/2019) Available at: National Diet and Nutrition Survey (publishing.service.gov.uk).

Bates, B.; Cox, L.; Nicholson, S.; Page, P.; Prentice, A.; Steer, T.; Swan, G. (2016) National Diet and Nutrition Survey Results from Years 5 and 6 (combined) of the Rolling Programme (2012/2013 – 2013/2014) Available at: Main heading (publishing.service.gov.uk).

Bates, B.; Lennox, A.; Prentice, A.; Bates, C.; Page, P.; Nicholson, S.; Swan, G. (2014) National Diet and Nutrition Survey Results from Years 1, 2, 3 and 4 (combined) of the Rolling Programme (2008/2009 – 2011/2012) Available at: Main heading (publishing.service.gov.uk).

Bettini et al., 2017. Food-grade TiO₂ impairs intestinal and systemic immune homeostasis, initiates preneoplastic lesions and promotes aberrant crypt development in the rat colon. *Scientific Reports*, 7, 40373.

Bischoff et. al., 2020. Possible Adverse Effects of Food Additive E171 (Titanium Dioxide) Related to Particle Specific Human Toxicity, Including the Immune System. *International Journal of Molecular Science*, 22(1):207.

Blevins et al., 2019. Evaluation of immunologic and intestinal effects in rats administered an E171-containing diet, a food grade titanium dioxide (TiO₂). *Food and Chemical Toxicology: An International Journal Published for the British Industrial Biological Research Association*, 133.

Canli EG, Gumus C, Canli M and Ila HB (2020): The effects of titanium nanoparticles on enzymatic and non-enzymatic biomarkers in female Wistar rats, *Drug and Chemical Toxicology*, 1–9.

Centres for Disease Control and Prevention (2011) Occupational Exposure to Titanium dioxide: [2011-160.pdf](#)

Chen, Z., Zheng, P., Han, S., Zhang, J., Li, Z., Zhou, S. and Jia, G. (2020) Tissue-specific oxidative stress and element distribution after oral exposure to titanium dioxide nanoparticles in rats. *Nanoscale*, 12(38), pp.20033-20046.

Cho, W.S., Kang, B.C., Lee, J.K., Jeong, J., Che, J.H. and Seok, S.H. (2013) Comparative absorption, distribution, and excretion of titanium dioxide and zinc oxide nanoparticles after repeated oral administration. *Particle and fibre toxicology*, 10(1), pp.1-9.

Cheng L and Lai MD (2003). Aberrant crypt foci as microscopic precursors of colorectal cancer. *World J Gastroenterol*. Dec, 9(12).

Clapper, M.L., Chang, W.C.L. and Cooper, H.S., 2020. Dysplastic aberrant crypt foci: biomarkers of early colorectal neoplasia and response to preventive intervention. *Cancer prevention research*, 13(3), pp.229-240.

Colucci, P., Turgeon, J. and Ducharme, M.P. (2011) How critical is the duration of the sampling scheme for the determination of half-life, characterization of exposure and assessment of bioequivalence? *Journal of Pharmacy & Pharmaceutical Sciences*. 14(2) pp.217-226.

Comera C, Cartier C, Gaultier E, Catrice O, Panouille Q, El Hamdi S, Tirez K, Nelissen I, Theodorou V and Houdeau E, (2020). Jejunal villus absorption and paracellular tight junction permeability are major routes for early intestinal uptake of food-grade TiO₂ particles: an in vivo and ex vivo study in mice. *Part Fibre Toxicol*, 17, 26.

Commission Regulation (EU) 2022/63 of 14 January 2022 amending Annexes II and III to Regulation (EC) No 1333/2008 of the European Parliament and of the Council as regards the food additive titanium dioxide (E171). [Publications Office](#).

COT Statement on Fluorine in the 1997 Diet Study: [fluoride.PDF](#)

COT Interim Position Paper, 2022 2021: [TiO₂ COT Interim position paper \(food.gov.uk\)](#)

Demir E, Akça H, Turna F, Aksakal S, Burgucu D, Kaya B, Tokgün O, Vales G, Creus A, Marcos R. (2015). Genotoxic and cell-transforming effects of titanium dioxide nanoparticles. *Environ Res*. 136:300-8.

Department of Health (2013). Diet and Nutrition Survey of Infants and Young Children (DNSIYC), 2011. Available at: [Diet and nutrition survey of infants and young children, 2011 - GOV.UK \(www.gov.uk\)](#)

Di Bucchianico, S., Cappellini, F., Le Bihanic, F., Zhang, Y., Dreij, K. and Karlsson, H.L. (2017) Genotoxicity of TiO₂ nanoparticles assessed by mini-gel comet assay and micronucleus scoring with flow cytometry. *Mutagenesis*, 32(1) pp.127-137.

Disdier C, Devoy J, Cosnefroy A, Chalansonnet M, Herlin-Boime N, Brun E, Lund A and Mabondzo A, (2015). Tissue biodistribution of intravenously administered titanium dioxide nanoparticles revealed blood-brain barrier clearance and brain inflammation in rat. *Particle and Fibre Toxicology*, 12, 24.

Donner EM, Myhre A, Brown SC, Boatman R, Warheit DB. (2016) In vivo micronucleus studies with 6 titanium dioxide materials (3 pigment-grade & 3

nanoscale) in orally-exposed rats. Regul Toxicol Pharmacol. 74:64-74.

Dudefoi, W., Moniz, K., Allen-Vercoe, E., Ropers, M.H. and Walker, V.K., (2017b) Impact of food grade and nano-TiO₂ particles on a human intestinal community. Food and Chemical Toxicology, 106 pp.242-249.

European Commission, Directorate-General for Health and Food Safety, Opinion on titanium dioxide (nano form) coated with cetyl phosphate, manganese dioxide or triethoxycaprylylsilane as UV-filter in dermally applied cosmetic, (2017) Publications Office. [Opinion on titanium dioxide \(nano form\) coated with cetyl phosphate, manganese dioxide or triethoxycaprylylsilane as UV-filter in dermally applied cosmetic - Publications Office of the EU \(europa.eu\)](#)

ECHA (2021) Guide on the classification and labelling of titanium dioxide. [ECHA Guide on the classification and labelling of titanium dioxide](#)

EFSA ANS Panel (EFSA Panel on Food Additives and Nutrients Sources added to Food), 2016. Re-evaluation of titanium dioxide (E171) as a food additive. EFSA Journal 2016;14(9):4545, 83 pp.

EFSA ANS Panel (EFSA Panel on Food Additives and Nutrients Sources added to Food), 2017. Approach followed for the refined exposure assessment as part of the safety assessment of food additives under re-evaluation. EFSA Journal 2017;15(10):5042, 9 pp.

EFSA FAF Panel (EFSA Panel on Food Additive and Flavourings), 2019. Scientific opinion on the proposed amendment of the EU specification for titanium dioxide (E171) with respect to the inclusion of additional parameters related to its particle size distribution. EFSA Journal 2019;17(7):5760, 23 pp.

EFSA FAF Panel (EFSA Panel on Food Additives and Flavourings) (2021) Scientific Opinion on the safety assessment of titanium dioxide (E171) as a food additive. EFSA Journal 2021;19(5):6585, 130 pp.

Farrell TP and Magnuson B. (2017). Absorption, Distribution and Excretion of Four Forms of Titanium Dioxide Pigment in the Rat. J Food Sci. 82(8):1985-1993.

Titanium Dioxide as a Food Additive Food Standards Australia New Zealand Report (2022) [FSANZ TiO₂ Assessment report.pdf](#).

Geiss, O., Ponti, J., Senaldi, C., Bianchi, I., Mehn, D., Barrero, J., Gilliland, D., Matissek, R. and Anklam, E. (2020) Characterisation of food grade titania with respect to nanoparticle content in pristine additives and in their related food

Heringa MB, Peters RJB, Bleys R, van der Lee MK, Tromp PC, van Kesteren PCE, van Eijkeren JCH, Undas AK, Oomen AG and Bouwmeester H, (2018). Detection of titanium particles in human liver and spleen and possible health implications. *Particle and Fibre Toxicology*, 15, 15.

Huang C, Sun M, Yang Y, Wang F, Ma X, Li J, Wang Y, Ding Q, Ying H, Song H, Wu Y, Jiang Y, Jia X, Ba Q, Wang H. (2017). Titanium dioxide nanoparticles prime a specific activation state of macrophages. *Nanotoxicology*. 11(6):737-750.

Hummel, T.Z., Kindermann, A., Stokkers, P.C., Benninga, M.A. and ten Kate, F.J., (2014) Exogenous pigment in Peyer patches of children suspected of having IBD. *Journal of pediatric gastroenterology and nutrition*, 58(4) pp.477- 480.

IARC (International Agency for Research on Cancer)/WHO, 2010. IARC Monographs on the Evaluation of Carcinogenic Risks to Humans. Volume 93. Carbon Black, Titanium Dioxide, and Talc. Available online: [IARC Publications Website - Carbon Black, Titanium Dioxide, and Talc](#)

Janer, G., Del Molino, E.M., Fernández-Rosas, E., Fernández, A. and Vázquez-Campos, S. (2014) Cell uptake and oral absorption of titanium dioxide nanoparticles. *Toxicology Letters*, 228(2) pp.103-110.

JECFA (1969) Joint FAO/WHO Expert Committee on Food Additives. Toxicological evaluation of some food colours, emulsifiers, stabilizers, anti-caking agents and certain other substances. FAO Nutrition Meetings Report Series No. 46A WHO/FOOD ADD/70 pp 36. [FSANZ TiO2 Assessment report.pdf](#)

FAO/WHO (Joint FAO/WHO Expert Committee on Food Additives). 2023. Risk assessment of titanium dioxide. Available at: Joint [FAO/WHO Expert Committee on Food Additives risk assessment of titanium dioxide risk released – background information](#).

Jensen DM, Løhr M, Sheykhzade M, Lykkesfeldt J, Wils RS, Loft S, Møller P. (2019). Telomere length and genotoxicity in the lung of rats following intragastric exposure to food-grade titanium dioxide and vegetable carbon particles. *Mutagenesis*. 29;34(2):203-214.

Jones, K., Morton, J., Smith, I., Jurkschat, K., Harding, A.H. and Evans, G., (2015) Human in vivo and in vitro studies on gastrointestinal absorption of titanium dioxide nanoparticles. *Toxicology letters*, 233(2) pp.95-101.

Kirkland D, Aardema M, Henderson L, Müller L. (2005) Evaluation of the ability of a battery of three in vitro genotoxicity tests to discriminate rodent carcinogens and non-carcinogens I. Sensitivity, specificity and relative predictivity. Mutation Research/Genetic Toxicology and Environmental Mutagenesis. 584(1-2) pp 1-256.

Kirkland D, Pfuhler S, Tweats D, Aardema M, Corvi R, Darroudi F, Elhajouji A, Glatt H, Hastwell P, Hayashi M, Kasper P, Kirchner S, Lynch A, Marzin D, Maurici D, Meunier JR, Müller L, Nohynek G, Parry J, Parry E, Thybaud V, Tice R, van Benthem J, Vanparys P, White P. (2007) How to reduce false positive results when undertaking in vitro genotoxicity testing and thus avoid unnecessary follow-up animal tests: Report of an ECVAM Workshop. Mutation Research/Genetic Toxicology and Environmental Mutagenesis. 628(1) pp 31-55.

Karimipour et al. 2018. Oral administration of titanium dioxide nanoparticle through ovarian tissue alterations impairs mice embryonic development. International Journal of Reproductive Biomedicine (Yazd, Iran), 16, 397–404.

Khorsandi et al., 2016. Morphometric and stereological assessment of the effects of titanium dioxide nanoparticles on the mouse testicular tissue. Bratislavske lekarske listy, 117, 659–664.

Khorsandi et al., 2017. Beneficial effects of quercetin on titanium dioxide nanoparticles induced spermatogenesis defects in mice. Environmental Science and Pollution Research International, 24, 5595–5606.

Koivisto AJ, Trabucco S, Ravegnani F et al (2022). Nanosized titanium dioxide particle emission potential from a commercial indoor air purifier photocatalytic surface: A case study [version 1; peer review: 4 approved]. Open Res Europe 2022, 2:84 [Nanosized titanium dioxide particle emission potential from a commercial indoor air purifier photocatalytic surface: A case study](#)

Kreyling WG, Holzwarth U, Haberl N, Kozempel J, Hirn S, Wenk A, Schleh C, Schaffler M, Lipka J, Semmler-Behnke M and Gibson N, (2017a). Quantitative biokinetics of titanium dioxide nanoparticles after intravenous injection in rats: part 1. Nanotoxicology., 11, 434–442.

Kreyling WG, Holzwarth U, Schleh C, Kozempel J, Wenk A, Haberl N, Hirn S, Schaffler M, Lipka J, Semmler-Behnke M and Gibson N, (2017b). Quantitative biokinetics of titanium dioxide nanoparticles after oral application in rats: part 2. Nanotoxicology, 11, 443–453.

Lamas, B., Martins Breyner, N. and Houdeau, E., 2020. Impacts of foodborne inorganic nanoparticles on the gut microbiota-immune axis: potential consequences for host health. **Particle and fibre toxicology**, 17(1), pp.1-22.

Lee et al. 2019. Titanium dioxide nanoparticles oral exposure to pregnant rats and its distribution. *Particle and Fibre Toxicology*, 16, 31.

MacNicoll, A., Kelly, M., Aksoy, H., Kramer, E., Bouwmeester, H. and Chaudhry, Q. (2015) A study of the uptake and biodistribution of nano-titanium dioxide using in vitro and in vivo models of oral intake. *Journal of Nanoparticle Research*, 17(2) pp.1-20.

Mortensen

National Cancer Institute. (1979). Bioassay of Titanium Dioxide for Possible Carcinogenicity. *Carcinogenicity*. 97, [Online]. Available at: [Bioassay of Titanium Dioxide for Possible Carcinogenicity](#)

Netherlands Food and Consumer Product Safety Authority (2019): Opinion of BuRO on possible health effects of the food additive titanium dioxide (E171).

OECD Test Guidelines for Chemical Testing [OECD Test Guidelines for Chemicals - OECD](#)

Pele LC, Thoree V, Bruggaber S, Koller D, Thompson RP, Lomer MC and Powell JJ, (2015). Pharmaceutical/food grade titanium dioxide particles are absorbed into the bloodstream of human volunteers. *Particle and Fibre Toxicology*, 12, 26.

Peters RJB, Oomen AG, van Bommel G, van Vliet L, Undas AK, Munniks S, Bleys RLAW, Tromp PC, Brand W and van der Lee M, (2020). Silicon dioxide and titanium dioxide particles found in human tissues. *Nanotoxicology*, 14,420–432.

Phue, W.H., Xu, K. and George, S. (2022) Inorganic food additives nanomaterials alter the allergenicity of milk proteins. *Food and Chemical Toxicology* pp.112874.

Pinget, G., Tan, J., Janac, B., Kaakoush, N.O., Angelatos, A.S., O'Sullivan, J., Koay, Y.C., Sierro, F., Davis, J., Divakarla, S.K. and Khanal, D. (2019) Impact of the food additive titanium dioxide (E171) on gut microbiota-host interaction. *Frontiers in nutrition*, 6 p.57.

Proquin, H., Jetten, M.J., Jonkhout, M.C., Garduño-Balderas, L.G., Briedé, J.J., de Kok, T.M., van Loveren, H. and Chirino, Y.I. (2018b) Transcriptomics analysis reveals new insights in E171-induced molecular alterations in a mouse model of

colon cancer. Scientific reports, 8(1) pp.1-16.

Riedle et al. (2020) A murine oral-exposure model for nano- and micro-particulates: demonstrating human relevance with food-grade titanium dioxide. Nano-Micro Small, 16, 2000486.

Roberts, C.; Steer, T.; Maplethorpe, N.; Cox, L.; Meadows, S.; Page, P.; Nicholson, S.; Swan, G. (2018) National Diet and Nutrition Survey Results from Years 7 and 8 (combined) of the Rolling Programme (2014/2015 – 2015/2016) Available at: [National Diet and Nutrition Survey](https://publishing.service.gov.uk) (publishing.service.gov.uk)

Rodríguez-Escamilla, J.C., Medina-Reyes, E.I., Rodríguez-Ibarra, C., Déciga-Alcaraz, A., Flores-Flores, J.O., Ganem-Rondero, A., Rodríguez-Sosa, M., Terrazas, L.I., Delgado-Buenrostro, N.L. and Chirino, Y.I. (2019) Food-grade titanium dioxide (E171) by solid or liquid matrix administration induces inflammation, germ cells sloughing in seminiferous tubules and blood-testis barrier disruption in mice. Journal of Applied Toxicology, 39(11) pp.1586- 1605.

Rompelberg, C., Heringa, M.B., van Donkersgoed, G., Drijvers, J., Roos, A., Westenbrink, S., Peters, R., van Bemmelen, G., Brand, W. and Oomen, A.G., 2016. Oral intake of added titanium dioxide and its nanofraction from food products, food supplements and toothpaste by the Dutch population. *Nanotoxicology*, 10 (10), pp.1404-1414.

Ropers, M.H., Terrisse, H., Mercier-Bonin, M. and Humbert, B. (2017) Titanium dioxide as food additive. Rijeka InTech, pp. 1-22.

Sadiq R, Bhalli JA, Yan J, Woodruff RS, Pearce MG, Li Y, Mustafa T, Watanabe F, Pack LM, Biris AS, Khan QM, Chen T. (2012). Genotoxicity of TiO₂ anatase nanoparticles in B6C3F1 male mice evaluated using Pig-a and flow cytometric micronucleus assays. *Mutat Res.*, 745(1-2):65-72.

SCCNFP (Scientific Committee on Cosmetics and Non-Food Products), 2000. Opinion of the Scientific Committee on Cosmetic Products and Non-Food Products Intended for Consumers concerning titanium dioxide. [Opinion concerning Titanium Dioxide, Colipa n° S75 adopted by the SCCNFP during the 14th plenary meeting of 24 October 2000 \(europa.eu\)](#)

SCCS (Scientific Committee on Consumer Safety), 2017. Opinion Titanium Dioxide (nano form) coated with Cetyl Phosphate, Manganese Dioxide or Triethoxycaprylylsilane as UV-filter in dermally applied cosmetic. Adopted on 7 March 2017 (corrigendum 22 June 2018). [sccs_o_202_0.pdf \(europa.eu\)](#)

SCCS (Scientific Committee on Consumer Safety), 2020. Opinion on Titanium dioxide (TiO₂) used in cosmetic products that lead to exposure by inhalation. Adopted on 6 October 2020. [Opinion on Titanium dioxide \(TiO₂\) used in cosmetic products that lead to exposure by inhalation \(europa.eu\)](https://ec.europa.eu/food/scs/scs_opinion_on_titanium_dioxide_en)

SCF (Scientific Committee for Food) (1977): Reports of the Scientific Committee for Food: Fourth Series, p. 27.

Shakeel M., Jabeen F., Shabbir S., Asghar M.S., Khan M.S., Chaudhry A.S, 2016. Toxicity of nano-titanium dioxide (TiO₂-NP) through various routes of exposure: A review. Biol. Trace. Elem. Res. 2016;172:1–36. doi: 10.1007/s12011-015-0550-x.

Shelby MD, Erexson GL, Hook GJ and Tice RR. (1993) Evaluation of a three-exposure mouse bone marrow micronucleus protocol: results with 49 chemicals. Environmental and Molecular Mutagenesis. 21, pp 160–179.

Shelby MD and Witt KL. (1995) Comparison of results from mouse bone marrow aberration and micronucleus test. Environmental and Molecular Mutagenesis. 25 pp 302–313.

Shwter AN, Abdullah NA, Alshawsh MA, El-Seedi HR, Al-Henhena NA, Khalifa SA and Abdulla MA, (2016) Chemopreventive effect of Phaleria macrocarpa on colorectal cancer aberrant crypt foci in vivo. Journal of Ethnopharmacology. 193 pp 195–206.

Sofranko, A., Wahle, T., Heusinkveld, H.J., Stahlmecke, B., Dronov, M., Pijnenburg, D., Hilhorst, R., Lamann, K., Albrecht, C. and Schins, R.P., (2021) Evaluation of the neurotoxic effects of engineered nanomaterials in C57BL/6J mice in 28-day oral exposure studies. Neurotoxicology, 84, pp.155-171.

Stocco A, Di Bucchianico S, Coppedè F, Ponti J, Ubaldi C, Blosi M, Delpivo C, Ortelli S, Costa AL, Migliore L. (2017). Multiple endpoints to evaluate pristine and remediated titanium dioxide nanoparticles genotoxicity in lung epithelial A549 cells. Toxicol Lett., 276:48-61.

Swiatkowska, I., Martin, N. and Hart, A.J. (2019) Blood titanium level as a biomarker of orthopaedic implant wear. Journal of Trace Elements in Medicine and Biology, 53, pp.120-128.

Talamini, L., Gimondi, S., Violatto, M.B., Fiordaliso, F., Pedica, F., Tran, N.L., Sitia, G., Aureli, F., Raggi, A., Nelissen, I. and Cubadda, F. (2019) Repeated administration of the food additive E171 to mice results in accumulation in

intestine and liver and promotes an inflammatory status. *Nanotoxicology*, 13(8), pp.1087-1101.

Tassinari et al., 2014. Oral, short-term exposure to titanium dioxide nanoparticles in Sprague-Dawley rat: focus on reproductive and endocrine systems and spleen. ***Nanotoxicology***, 8, 654– 662.

TDMA (2020). Extended One-Generation Reproductive Toxicity study of titanium dioxide E171 in rats by oral administration via the diet.

Ünal, F., Demirtaş Korkmaz, F., Suludere, Z., Erol, Ö. (2021) Genotoxicity of Two Nanoparticles: Titanium Dioxide and Zinc Oxide. *Gazi University Journal of Science*, 34(4),pp 948-958.

Verleysen (2020). Physico-chemical characterization of the pristine E171 food additive by standardized and validated methods. *Nanomaterials (Basel)*, 10, pp 592.

Verleysen E, Waegeneers N, De Vos S, Brassinne F, Ledecq M, Van Steen F, Andjelkovic M, Janssens R, Mathioudaki S, Delfosse L, Machiels R, Cheyns K and Mast J, (2021) Physicochemical characterization of nanoparticles in food additives in the context of risk identification. EFSA supporting publication 2021 31 pp.

Wang, J., Zhou, G., Chen, C., Yu, H., Wang, T., Ma, Y., Jia, G., Gao, Y., Li, B., Sun, J. and Li, Y. (2007) Acute toxicity and biodistribution of different sized titanium dioxide particles in mice after oral administration. *Toxicology letters*, 168(2), pp.176-185.

Warheit et al. (2015b). Developmental toxicity studies with 6 forms of titanium dioxide test materials (3 pigment-different grade & 3 nanoscale) demonstrate an absence of effects in orally-exposed rats. *Regulatory Toxicology and Pharmacology*, 73, 887–896.

Warheit DB, Brown SC, Donner EM. (2015) Acute and subchronic oral toxicity studies in rats with nanoscale and pigment grade titanium dioxide particles. *Food Chemical Toxicology* 84, 208-24.

Weir, A., Westerhoff, P., Fabricius, L., Hristovski, K., & Von Goetz, N. (2012). Titanium dioxide nanoparticles in food and personal care products. *Environmental science & technology*, 46(4), 2242-2250.

West, B. & Wyzan, H. (1963) Unpublished report by American Cyanamid Company (cited by JECFA 1969).

Winkler HC, Notter T, Meyer U, Naegeli H. (2018). Critical review of the safety assessment of titanium dioxide additives in food. *J Nanobiotechnology*. 16(1):51.

Yang, X. H., Fu, H. T., Wang, X. C., Yang, J. L., Jiang, X. C., & Yu, A. B. (2014). Synthesis of silver-titanium dioxide nanocomposites for antimicrobial applications. *Journal of nanoparticle research*, 16 pp 1-13.