

Deriving a health-based guidance value for antimony to support development of UK Drinking Water Standards

# Health Canada

## In this guide

### [In this guide](#)

1. [Introduction and Background](#)
2. [Properties of antimony](#)
3. [Toxicokinetics and Toxicity](#)
4. [Summary of the Poon et al. \(1998\) study](#)
5. [Lynch et al. \(1999\) interpretation](#)
6. [Response from Valli et al. \(2000\)](#)
7. [HBGV's established by the WHO, ATSDR and Health Canada](#)
8. [Agency for Toxic Substances and Disease Registry \(ATSDR\)](#)
9. [Health Canada](#)
10. [Differences between WHO, ATSDR and Health Canada](#)
11. [Additional Toxicology Studies](#)
12. [Summary and Questions for the Committee](#)
13. [List of abbreviations and their full meanings](#)
14. [References- Deriving a health-based guidance value for antimony to support development of UK Drinking Water Standards](#)

## Health Canada

35. In 2024, Health Canada selected a NOAEL of 60 µg Sb /kg bw/day from the study by Poon et al., (1998), based on observed histopathological changes in the liver (anisokaryosis) and alterations in serum biochemistry indicative of liver effects. Using this NOAEL, a tolerable daily intake (TDI) was derived by applying an uncertainty factor (UF) of 300 to account for interspecies variation (×10), intraspecies variation (×10) and the use of a subchronic study (×3), resulting in a TDI of 0.2 µg/kg body weight per day (Health Canada,2024).

36. The Health Canada health-based drinking water value of 3 µg/L for total antimony in drinking water was then derived from this TDI using an average adult

body weight of 74 kg, a drinking water allocation factor of 0.3 (based on the upper bound of estimated intake for drinking water) and a drinking water intake rate of 1.53 L/day (Health Canada,2024).