

Overall Conclusion

In this guide

[In this guide](#)

1. [Executive Summary - Annex 1 to TOX/2025/23](#)
2. [Background and scope of discussion - Annex 1 to TOX/2025/23](#)
3. [Properties of antimony and sources in drinking water - Annex 1 to TOX/2025/23](#)
4. [Oral toxicity data for antimony - Annex 1 to TOX/2025/23](#)
5. [HBGVs established by WHO, ATSDR and Health Canada - Annex 1 to TOX/2025/23](#)
6. [Discussion - Annex 1 to TOX/2025/23](#)
7. [Overall Conclusion - Annex 1 to TOX/2025/23](#)
8. [List of abbreviations and their full meanings - Annex 1 to TOX/2025/23](#)
9. [References - Annex 1 to TOX/2025/23](#)
10. [Annex A - Annex 1 to TOX/2025/23](#)
11. [Annex A References - Annex 1 to TOX/2025/23](#)

This is a draft position statement for discussion. This does not represent the views of the Committee and should not be cited.

35. Overall, the COT concluded that the NOAEL of 6,000 µg Sb/kg bw/day, from the Poon et al. (1998) study based on decreased body weight gain and reduced food and water consumption in adult rats, was the appropriate point of departure to use as the basis of a HBGV for antimony.

36. The Committee also highlighted that the pentavalent form of antimony, which is predominant in drinking water, exhibits lower toxicity compared to the trivalent form. As Poon et al. (1998) utilized the trivalent form of antimony (antimony potassium tartrate) in their study, a HBGV derived from the NOAEL of 6,000 µg Sb/kg bw/day was considered a sufficiently protective for antimony in drinking water.

37. The Committee recommended a UF of 300, comprising a factor of 10 for interspecies variation, 10 for intraspecies variation, and 3 for subchronic to chronic extrapolation. This results in a TDI of 20 µg Sb/kg bw/day.

COT Month 2025

Statement 2025/XX