Annex B to TOX/2025/26 - First draft statement on the risk for T-2 and HT-2 mycotoxins in food

Table 10

In this guide

In this guide

- 1. Supplementary information Annex B to TOX/2025/26
- 2. <u>Table 1 Annex B to TOX/2025/26</u>
- 3. Table 2 Annex B to TOX/2025/26
- 4. Table 3 Annex B to TOX/2025/26
- 5. Table 4 Annex B to TOX/2025/26
- 6. Table 5 Annex B to TOX/2025/26
- 7. Table 6 Annex B to TOX/2025/26
- 8. Table 7 Annex B to TOX/2025/26
- 9. Table 8 Annex B to TOX/2025/26
- 10. <u>Table 9 Annex B to TOX/2025/26</u>
- 11. <u>Table 10 Annex B to TOX/2025/26</u>
- 12. Table 11 Annex B to TOX/2025/26
- 13. Table 12 Annex B to TOX/2025/26
- 14. Table 13 Annex B to TOX/2025/26
- 15. Table 14 Annex B to TOX/2025/26
- 16. Table 15 Annex B to TOX/2025/26
- 17. Table 16 Annex B to TOX/2025/26

This is a paper for discussion. This does not represent the views of the Committee and should not be cited.

Table 10: Acute risk

Acute risk to sum of T-2 and HT-2 from some cereal grains based on DNSYIC and NDNS years 1-11 consumption data and occurrence data collected from the FSA call, expressed as estimated x-fold of chronic exposure as a percentage of the ARfD of 0.3 μ g/kg bw/day.

		Infants (4-18 months) (n=2683)	(1.5-3 years)	Toddlers (1.5-3 years) (n=1157)	(4-10 years)	(4-10 years)	Older Children (11-18 years) (n=2657)
Food groups	Mean*	P97.5*	Mean *	P97.5*	Mean*	P97.5*	Mean*
Unprocessed oats	0.3	2	0.3	2	0.2	1	0.1
Unprocessed oats with reduction factor	0-0.1	0.3	0-0.1	0.2-0.3	0	0.1-0.2	0
Oats combined***	0-0.1	0.3	0-0.1	0.2-0.3	0	0.2	0
All grains sum (unprocessed)	0-0.4	0-2	0-0.4	0-2	0-0.3	0-1	0-0.1
All grains sum (processed)	0.1	0.3	0-0.1	0.2-0.3	0.1	0.2	0

*Estimates have been rounded to 1 significant figure; except in cases where the exceedance is <1.

+ Oat grains with reduction factor - unprocessed oat grains with reduction factor (85%) applied.

++ Oats combined - combination of processed oat grains and unprocessed oat grains with reduction factor (85%) applied.