Annex B to TOX/2025/26 - First draft statement on the risk for T-2 and HT-2 mycotoxins in food

Table 11

In this guide

In this guide

- 1. Supplementary information Annex B to TOX/2025/26
- 2. Table 1 Annex B to TOX/2025/26
- 3. Table 2 Annex B to TOX/2025/26
- 4. Table 3 Annex B to TOX/2025/26
- 5. Table 4 Annex B to TOX/2025/26
- 6. Table 5 Annex B to TOX/2025/26
- 7. Table 6 Annex B to TOX/2025/26
- 8. Table 7 Annex B to TOX/2025/26
- 9. Table 8 Annex B to TOX/2025/26
- 10. Table 9 Annex B to TOX/2025/26
- 11. Table 10 Annex B to TOX/2025/26
- 12. Table 11 Annex B to TOX/2025/26
- 13. Table 12 Annex B to TOX/2025/26
- 14. Table 13 Annex B to TOX/2025/26
- 15. Table 14 Annex B to TOX/2025/26
- 16. Table 15 Annex B to TOX/2025/26
- 17. Table 16 Annex B to TOX/2025/26

This is a paper for discussion. This does not represent the views of the Committee and should not be cited.

Table 11: Chronic exposure

Chronic exposure to sum of T-2 and HT-2 from RTE products from DNSIYC and NDNS years 1-11 consumption data and occurrence data collected from the FSA call.

Infants (4-18 months) (n=2683)

Food Groups	Count of T2 HT2 Data Call Samples (n)	Mean T2 Concentration (μg/kg)	Max T2 Concentration (μg/kg)	Consumers (n)	Mean	P97.5	Mean	P97.5
Infants'	2	110.6	219	1128	0.36	1.5	0.71	2.9

Toddlers (1.5-3 years) (n=1157)

Food Groups	Count of T2 HT2 Data Call	Mean T2 Concentration (μg/kg)	Max T2 Concentration (μg/kg)	Consumers (n)	Mean	P97.5	Mean	P97.
	Samples (n)							
Infants' cereals	2	110.6	219	34	0.22	0.70	0.44	1.4

Exposure estimates have been rounded to 2 significant figures; concentrations have been rounded to one decimal place.

^{*}Mean/Max concentration.