

Table 12 -

In this guide

[In this guide](#)

1. [Supplementary information - Annex B to TOX/2025/26](#)
2. [Table 1 - Annex B to TOX/2025/26](#)
3. [Table 2 - Annex B to TOX/2025/26](#)
4. [Table 3 - Annex B to TOX/2025/26](#)
5. [Table 4 - Annex B to TOX/2025/26](#)
6. [Table 5 - Annex B to TOX/2025/26](#)
7. [Table 6 - Annex B to TOX/2025/26](#)
8. [Table 7 - Annex B to TOX/2025/26](#)
9. [Table 8 - Annex B to TOX/2025/26](#)
10. [Table 9 - Annex B to TOX/2025/26](#)
11. [Table 10 - Annex B to TOX/2025/26](#)
12. [Table 11 - Annex B to TOX/2025/26](#)
13. [Table 12 - Annex B to TOX/2025/26](#)
14. [Table 13 - Annex B to TOX/2025/26](#)
15. [Table 14 - Annex B to TOX/2025/26](#)
16. [Table 15 - Annex B to TOX/2025/26](#)
17. [Table 16 - Annex B to TOX/2025/26](#)

This is a paper for discussion. This does not represent the views of the Committee and should not be cited.

Table 12: Acute exposure

Acute exposure to sum of T-2 and HT 2 from RTE products from DNSIYC and NDNS years 1-11 consumption data and occurrence data collected from the FSA call.

Infants (4-18 months) (n=2683)

Food Groups	Count of T2							
	HT2 Data Call Samples (n)	Mean T2 Concentration (µg/kg)	Max T2 Concentration (µg/kg)	Consumers (n)	Mean P97.5		Mean P97.5	
Infants' cereals	2	110.6	219	1128	0.36	1.5	0.71	2.9

Toddlers (1.5-3 years) (n=1157)

Food Groups	Count of T2 HT2 Data Call	Mean T2 Concentration (µg/kg)	Max T2 Concentration (µg/kg)	Consumers (n)	Mean P97.5			
	Samples (n)							
Infants' cereals	2	110.6	219	34	0.22	0.70	0.44	1.4

Exposure estimates have been rounded to 2 significant figures; concentrations have been rounded to one decimal place.

*Mean/Max concentration.