

Mode of action

In this guide

[In this guide](#)

1. [Introduction and Background - EFSA draft scientific opinion on risks for human health of plant lectins in food](#)
2. [Summary of 2025 EFSA draft evaluation - EFSA draft scientific opinion on risks for human health of plant lectins in food](#)
3. [Acute toxicity studies - EFSA draft scientific opinion on risks for human health of plant lectins in food](#)
4. [Repeat dose toxicity studies - EFSA draft scientific opinion on risks for human health of plant lectins in food](#)
5. [Observations in Humans - EFSA draft scientific opinion on risks for human health of plant lectins in food](#)
6. [Mode of action - EFSA draft scientific opinion on risks for human health of plant lectins in food](#)
7. [Occurrence data and dietary exposure assessment for the European population - EFSA draft scientific opinion on risks for human health of plant lectins in food](#)
8. [Risk characterisation - EFSA draft scientific opinion on risks for human health of plant lectins in food](#)
9. [Uncertainty analysis - EFSA draft scientific opinion on risks for human health of plant lectins in food](#)
10. [Recommendations - EFSA draft scientific opinion on risks for human health of plant lectins in food](#)
11. [Questions on which the views of the Committee are sought - EFSA draft scientific opinion on risks for human health of plant lectins in food](#)
12. [List of Abbreviations - EFSA draft scientific opinion on risks for human health of plant lectins in food](#)
13. [References - EFSA draft scientific opinion on risks for human health of plant lectins in food](#)

This is a paper for discussion. This does not represent the views of the Committee and should not be cited.

24. The adverse effects of lectins are dependent on a) structure and carbohydrate binding activity, b) number of carbohydrate binding sites, c) overall lectin content, and d) arrangement of sub-units and their interaction with glycans. By binding to carbohydrate moieties within the gut epithelial cells, lectins can affect the microvilli, villi and crypts. It has further been suggested that as a result of lectin consumption, the exocrine pancreas may produce digestive juice and grow due to the release of cholecystokinin from enteroendocrine cells into the bloodstream.

25. Lectins have demonstrated that they are capable of binding to the gut epithelial cells. Lectins have also been thought to be involved in the induction of autoimmune and allergic responses as a result of their effect on the gut barrier and microbiota. The animal species and the amount of lectin within the circulation would thereby affect the type of immune response.

26. EFSA considered that lectins could modulate intracellular/intercellular signalling pathways by binding to gut receptors which may affect viability.