

Table 10

In this guide

[In this guide](#)

1. [Table 1 - Annex B to TOX/2025/40](#)
2. [Table 2 - Annex B to TOX/2025/40](#)
3. [Table 3 - Annex B to TOX/2025/40](#)
4. [Table 4 - Annex B to TOX/2025/40](#)
5. [Table 5 - Annex B to TOX/2025/40](#)
6. [Table 6 - Annex B to TOX/2025/40](#)
7. [Table 7 - Annex B to TOX/2025/40](#)
8. [Table 8 - Annex B to TOX/2025/40](#)
9. [Table 9 - Annex B to TOX/2025/40](#)
10. [Table 10 - Annex B to TOX/2025/40](#)
11. [Table 11 - Annex B to TOX/2025/40](#)
12. [Table 12- Annex B to TOX/2025/40](#)
13. [Table 13- Annex B to TOX/2025/40](#)
14. [Table 14- Annex B to TOX/2025/40](#)
15. [Table 15- Annex B to TOX/2025/40](#)

This is a draft statement for discussion. It does not reflect the final views of the Committee and should not be cited.

Table 10. Chronic exposure to sum of T-2 and HT-2 from RTE products from DNSIYC and NDNS years 1-11 consumption data and occurrence data collected from the FSA call.

Infants (4-18 months) (n=2683)

Food groups	Count of T-2		T-2 Concentration $\mu\text{g/kg}$ (mean)	T-2 Concentration $\mu\text{g/kg}$ (max)	Consumers (n)	Mean*; Mean*; Max*; Ma			
	HT-2 data call samples (n)					mean	P97.5	mean	P9
Infants' cereals	2		110.6	219	1128	0.36	1.5	0.71	2.9

Toddlers (1.5-3 years) (n=1157)

Food groups	Count of T-2		T-2 Concentration $\mu\text{g/kg}$ (mean)	T-2 Concentration $\mu\text{g/kg}$ (max)	Consumers (n)	Mean*; Mean*; Max*; Ma			
	HT-2 data call samples (n)					mean	P97.5	mean	P9
Infants' cereals	2		110.6	219	34	0.22	0.70	0.44	1.4

Exposure estimates have been rounded to 2 significant figures; concentrations have been rounded to one decimal place.

*Mean/Max concentration.