

Table 12

In this guide

[In this guide](#)

1. [Table 1 - Annex B to TOX/2025/40](#)
2. [Table 2 - Annex B to TOX/2025/40](#)
3. [Table 3 - Annex B to TOX/2025/40](#)
4. [Table 4 - Annex B to TOX/2025/40](#)
5. [Table 5 - Annex B to TOX/2025/40](#)
6. [Table 6 - Annex B to TOX/2025/40](#)
7. [Table 7 - Annex B to TOX/2025/40](#)
8. [Table 8 - Annex B to TOX/2025/40](#)
9. [Table 9 - Annex B to TOX/2025/40](#)
10. [Table 10 - Annex B to TOX/2025/40](#)
11. [Table 11 - Annex B to TOX/2025/40](#)
12. [Table 12- Annex B to TOX/2025/40](#)
13. [Table 13- Annex B to TOX/2025/40](#)
14. [Table 14- Annex B to TOX/2025/40](#)
15. [Table 15- Annex B to TOX/2025/40](#)

This is a draft statement for discussion. It does not reflect the final views of the Committee and should not be cited.

Table 12: Chronic exposure to T-2 from RTE products from DNSIYC and NDNS years 1-11 consumption data and occurrence data collected from the FSA call.

Infants (4-18 months) (n=2683)

Food groups	Count of T-2 data call samples (n)	T-2 concentration µg/kg (mean)	T-2 concentration µg/kg (max)	Consumers (n)	*Mean; Mean	*Mean; P97.5	*Max; Mean
Plain muesli	3	4.0	5.9	91	0.00030	0.0010	0.0004
Biscuits, oatmeal	4	8.4	12.5	1441	0.0059	0.020	0.0088
Multigrain bread rolls	2	3.1	3.1	98	0.0027	0.0081	0.0027
Wheat bread rolls	2	3.1	3.1	435	0.0050	0.018	0.0050
Dried pasta	4	3.1	3.1	1140	0.0056	0.020	0.0056
Mixed breakfast cereals	3	5.0	8.9	1691	0.0095	0.036	0.017
Extruded breakfast cereals	4	5.0	10	338	0.0029	0.010	0.0057
Oat porridge	24	11.7	36.1	389	0.033	0.17	0.10

Plain cereal bars	2	5.9	8.7	5	0.0038	0.0056	0.0056
-------------------	---	-----	-----	---	--------	--------	--------

Toddlers (1.5-3 years) (n=1157)

Food groups	Count of T-2 data call samples (n)	T-2 concentration $\mu\text{g/kg}$ (mean)	T-2 concentration $\mu\text{g/kg}$ (max)	Consumers (n)	*Mean; Mean	*Mean; P97.5	*Max; Mean
Plain muesli	3	4.0	5.9	14	0.0043	0.0076	0.0062
Biscuits, oatmeal	4	8.4	12.5	566	0.0058	0.018	0.0086
Multigrain bread rolls	2	3.1	3.1	75	0.0053	0.019	0.0053
Wheat bread rolls	2	3.1	3.1	125	0.0063	0.021	0.0063
Dried pasta	4	3.1	3.1	602	0.0055	0.016	0.0055
Mixed breakfast cereals	3	5.0	8.9	983	0.0077	0.033	0.014

Extruded breakfast cereals	4	5.0	10	379	0.0032	0.010	0.0064
Oat porridge	24	11.7	36.1	70	0.038	0.14	0.12
Plain cereal bars	2	5.9	8.7	24	0.0032	0.0066	0.0046

Adults (19-64 years) (n=5094)

Food groups	Count of T-2 data call samples (n)	T-2 concentration µg/kg (mean)	T-2 concentration µg/kg (max)	Consumers (n)	*Mean; Mean	*Mean; *Max; P97.5	*Mean; *Max; Mean
Plain muesli	3	4.0	5.9	211	0.0015	0.0049	0.0022
Biscuits, oatmeal	4	8.4	12.5	1472	0.0017	0.0059	0.0025
Multigrain bread rolls	2	3.1	3.1	530	0.0018	0.0051	0.0018
Wheat bread rolls	2	3.1	3.1	293	0.0017	0.0055	0.0017

Dried pasta	4	3.1	3.1	1677	0.0027	0.0075	0.0027
Mixed breakfast cereals	3	5.0	8.9	2935	0.0021	0.0078	0.0037
Extruded breakfast cereals	4	5.0	10	278	0.0011	0.0031	0.0022
Oat porridge	24	11.7	36.1	222	0.012	0.036	0.038
Plain cereal bars	2	5.9	8.7	135	0.00098	0.0031	0.0014

Adult Vegetarians / Vegans (19-64 years) (n=170)

Food groups	Count of T-2 data call samples (n)	T-2 concentration µg/kg (mean)	T-2 concentration µg/kg (max)	Consumers (n)	*Mean; Mean	*Mean; P97.5	*Max; Mean
Plain muesli	3	4.0	5.9	11	0.0016	0.0032	0.0023
Biscuits, oatmeal	4	8.4	12.5	55	0.0018	0.0041	0.0026

Multigrain bread rolls	2	3.1	3.1	33	0.0020	0.0046	0.0020
Wheat bread rolls	2	3.1	3.1	10	0.0013	0.0033	0.0013
Dried pasta	4	3.1	3.1	64	0.0035	0.010	0.0035
Mixed breakfast cereals	3	5.0	8.9	98	0.0023	0.0058	0.0041
Extruded breakfast cereals	4	5.0	10	14	0.0010	0.0021	0.0020
Oat porridge	24	11.7	36.1	6	0.016	0.052	0.050
Plain cereal bars	2	5.9	8.7	2	0.0015	0.0016	0.0022

Exposure estimates have been rounded to 2 significant figures; concentrations have been rounded to one decimal place.

*Mean/Max concentration.