

Table 13

In this guide

[In this guide](#)

1. [Table 1 - Annex B to TOX/2025/40](#)
2. [Table 2 - Annex B to TOX/2025/40](#)
3. [Table 3 - Annex B to TOX/2025/40](#)
4. [Table 4 - Annex B to TOX/2025/40](#)
5. [Table 5 - Annex B to TOX/2025/40](#)
6. [Table 6 - Annex B to TOX/2025/40](#)
7. [Table 7 - Annex B to TOX/2025/40](#)
8. [Table 8 - Annex B to TOX/2025/40](#)
9. [Table 9 - Annex B to TOX/2025/40](#)
10. [Table 10 - Annex B to TOX/2025/40](#)
11. [Table 11 - Annex B to TOX/2025/40](#)
12. [Table 12- Annex B to TOX/2025/40](#)
13. [Table 13- Annex B to TOX/2025/40](#)
14. [Table 14- Annex B to TOX/2025/40](#)
15. [Table 15- Annex B to TOX/2025/40](#)

This is a draft statement for discussion. It does not reflect the final views of the Committee and should not be cited.

Table 13: Acute exposure to T-2 from RTE products from DNSIYC and NDNS years 1-11 consumption data and occurrence data collected from the FSA call.

Infants (4-18 months) (n=2683)

Food groups	Count of T-2 data call samples (n)	T-2 concentration µg/kg (mean)	T-2 concentration µg/kg (max)	Consumers (n)	*Mean; Mean	*Mean; P97.5	*Max; Mean
Plain muesli	3	4.0	5.9	91	0.00073	0.0025	0.0011
Biscuits, oatmeal	4	8.4	12.5	1441	0.014	0.037	0.020
Multigrain bread rolls	2	3.1	3.1	98	0.0075	0.019	0.0075
Wheat bread rolls	2	3.1	3.1	435	0.011	0.032	0.011
Dried pasta	4	3.1	3.1	1140	0.018	0.056	0.018
Mixed breakfast cereals	3	5.0	8.9	1691	0.017	0.084	0.030
Extruded breakfast cereals	4	5.0	10	338	0.0069	0.022	0.014
Oat porridge	24	11.7	36.1	389	0.074	0.27	0.23

Plain cereal bars	2	5.9	8.7	5	0.011	0.021	0.017	0.017
-------------------	---	-----	-----	---	-------	-------	-------	-------

Toddlers (1.5-3 years) (n=1157)

Food groups	Count of T-2 data call samples (n)	T-2 concentration $\mu\text{g/kg}$ (mean)	T-2 concentration $\mu\text{g/kg}$ (max)	Consumers (n)	*Mean; Mean	*Mean; P97.5	*Max; *Mean P	
Plain muesli	3	4.0	5.9	14	0.0093	0.019	0.014	0.014
Biscuits, oatmeal	4	8.4	12.5	566	0.016	0.039	0.023	0.023
Multigrain bread rolls	2	3.1	3.1	75	0.012	0.032	0.012	0.012
Wheat bread rolls	2	3.1	3.1	125	0.014	0.035	0.014	0.014
Dried pasta	4	3.1	3.1	602	0.017	0.046	0.017	0.017
Mixed breakfast cereals	3	5.0	8.9	983	0.014	0.057	0.025	0.025

Extruded breakfast cereals	4	5.0	10	379	0.0072	0.020	0.014	0
Oat porridge	24	11.7	36.1	70	0.11	0.27	0.34	0
Plain cereal bars	2	5.9	8.7	24	0.011	0.023	0.016	0

Adults (19-64 years) (n=5094)

Food groups	Count of T-2 data call samples (n)	T-2 concentration µg/kg (mean)	T-2 concentration µg/kg (max)	Consumers (n)	*Mean; Mean	*Mean; *Max; P97.5	*Mean; *Max; Mean	*Mean; *Max; Mean
Plain muesli	3	4.0	5.9	211	0.0028	0.0070	0.0041	0
Biscuits, oatmeal	4	8.4	12.5	1472	0.0046	0.014	0.0068	0
Multigrain bread rolls	2	3.1	3.1	530	0.0040	0.010	0.0040	0
Wheat bread rolls	2	3.1	3.1	293	0.0040	0.0093	0.0040	0

Dried pasta	4	3.1	3.1	1677	0.0087	0.019	0.0087	0
Mixed breakfast cereals	3	5.0	8.9	2935	0.0038	0.013	0.0067	0
Extruded breakfast cereals	4	5.0	10	278	0.0025	0.0052	0.0051	0
Oat porridge	24	11.7	36.1	222	0.026	0.050	0.081	0
Plain cereal bars	2	5.9	8.7	135	0.0025	0.0080	0.0037	0

Adult Vegetarians / Vegans (19-64 years) (n=170)

Food groups	Count of T-2 data call samples (n)	T-2 concentration $\mu\text{g/kg}$ (mean)	T-2 concentration $\mu\text{g/kg}$ (max)	Consumers (n)	*Mean; Mean	*Mean; P97.5	*Max; Mean	
Plain muesli	3	4.0	5.9	11	0.0034	0.0057	0.0050	
Biscuits, oatmeal	4	8.4	12.5	55	0.0052	0.011	0.0077	

Multigrain bread rolls	2	3.1	3.1	33	0.0040	0.0068	0.0040
Wheat bread rolls	2	3.1	3.1	10	0.0039	0.0092	0.0039
Dried pasta	4	3.1	3.1	64	0.010	0.020	0.010
Mixed breakfast cereals	3	5.0	8.9	98	0.0045	0.015	0.0079
Extruded breakfast cereals	4	5.0	10	14	0.0025	0.0045	0.0050
Oat porridge	24	11.7	36.1	6	0.039	0.10	0.12
Plain cereal bars	2	5.9	8.7	2	0.0040	0.0053	0.0060

Exposure estimates have been rounded to 2 significant figures; concentrations have been rounded to one decimal place.

*Mean/Max concentration.