## **Chemical composition**

## In this guide

## In this guide

- 1. Background and Introduction Garcinia cambogia oral consumption
- 2. Data from authoritative bodies Garcinia cambogia oral consumption
- 3. Chemical composition Garcinia cambogia oral consumption
- 4. ANSES Garcinia cambogia oral consumption
- 5. AESAN Garcinia cambogia oral consumption
- 6. Australian TGA Garcinia cambogia oral consumption
- 7. <u>BfR (German Federal Institute for Risk Assessment) Garcinia cambogia oral</u> consumption
- 8. EFSA Garcinia cambogia oral consumption
- 9. Health Canada Garcinia cambogia oral consumption
- 10. NCCIH Garcinia cambogia oral consumption
- 11. Data from literature search Garcinia cambogia oral consumption
- 12. Human data Garcinia cambogia oral consumption
- 13. Hepatotoxic case reports Garcinia cambogia oral consumption
- 14. Animal studies Garcinia cambogia oral consumption
- 15. Exposure Garcinia cambogia oral consumption
- 16. Uncertainties Garcinia cambogia oral consumption
- 17. Risk Characterisation Garcinia cambogia oral consumption
- 18. Questions to the Committee Garcinia cambogia oral consumption
- 19. Abbreviations Garcinia cambogia oral consumption
- 20. References Garcinia cambogia oral consumption

## This is a discussion paper. It does not reflect the views of the Committee. It should not be cited.

11. The bioactive compounds which are extracted and isolated from *G. cambogia* are shown in Table 1 and Figure 1.

**Table 1** - list of bioactive compounds from plant parts of G. cambogia and their related bioactivities (reproduced from Espirito Santo et al., 2020).


Xanthones	Plant part	Activity
Garbogiol	Roots	Inhibition of α-glucosid
Rheedia xanthone A	Peel	Not applicable*
Oxy-guttiferone i	Fruits	Not applicable*
Oxy-guttiferone k	Fruits	Not applicable*
Oxy-guttiferone k2	Fruits	Not applicable*
Oxy-guttiferone m	Fruits	Not applicable*
Benzophenones	Not applicable*	Not applicable*
Garcinol	Peel	Anticancer, anti-inflammatory, antiparasitic, action on nervous system
Isogarcinol	Peel	Anticancer, anti-inflammatory, antiparasitic, action on nervous system
Guttiferone i	Fruits	Not applicable*
Guttiferone n	Fruits	Not applicable*
Guttiferone j	Fruits	Not applicable*

Guttiferone k	Fruits	Topoisomerase II inhibitor
Guttiferone m	Fruits	Topoisomerase II inhibitor
Organic acids	Not applicable*	Not applicable*
Heterocyclic amines	Fruits	Antiobesity
Tartaric acid	Fruits	Not applicable*
Citric acid	Fruits	Not applicable*
Malic acid	Fruits	Antimicrobial
Garcinialactone	Fruits	Not applicable*

This figure shows 9 chemical structures of the xanthone and benzophenone classes of bioactive compounds from G. cambogia (reproduced from Espirito Santo et al., 2020).

**Figure 1** - chemical structures of the xanthone and benzophenone classes of bioactive compounds from G. cambogia. Xanthones: 40) oxy-guttiferone-i; 41) Oxy-guttiferone k; 42) Oxy-guttiferone k2 and 43) Oxy-guttiferone k. Benzophenones: 44) guttiferone-I; 45) guttiferone-j; 46) guttiferone-k; 47) guttiferone-n and 48) guttiferone-m (reproduced from Espirito Santo et al., 2020).

12. The HCA present in G. cambogia is a "potent" and "competent" inhibitor of adenosinetriphosphate (ATP) citrate lyase, which is a key enzyme in the synthesis of fatty acids, cholesterol, and triglycerides. It also regulates the level of serotonin, which has been associated with satiety, increased oxidation of fat, and decreased gluconeogenesis (Semwal et al., 2015; Preuss et al., 2004). HCA compromises a citric acid with a hydroxyl group at the second carbon. HCA has two diastereomers as there are two chiral centres, as such there are four stereoisomers of HCA, compromising two pairs of enantiomers (see Figure 2). Each of the stereoisomers can form a  $\gamma$ -lactone ring and in general solution, HCA is a mixture of non-lactone and lactone forms (Yamada et al., 2007).



This figure shows 4 structures of HCA derived from Garcinia ssp. It shows the non-lactone and lactone forms (reproduced from Yamada et al., 2007).

**Figure 2** - Structures of HCA derived from Garcinia ssp. Upper and bottom structures show the non-lactone and lactone forms, respectively. (2S,3S)-HCA (a mixture of 1a and 1b) is found in G. cambogia. The other isomers (2R,3R)-HCA (a mixture of 2a and 2b) and (2R,3S)-HCA have not been isolated from natural sources (reproduced from Yamada et al., 2007).