AESAN

In this guide

In this guide

- 1. Background and Introduction Garcinia cambogia oral consumption
- 2. Data from authoritative bodies Garcinia cambogia oral consumption
- 3. Chemical composition Garcinia cambogia oral consumption
- 4. ANSES Garcinia cambogia oral consumption
- 5. AESAN Garcinia cambogia oral consumption
- 6. Australian TGA Garcinia cambogia oral consumption
- 7. <u>BfR (German Federal Institute for Risk Assessment) Garcinia cambogia oral consumption</u>
- 8. EFSA Garcinia cambogia oral consumption
- 9. Health Canada Garcinia cambogia oral consumption
- 10. NCCIH Garcinia cambogia oral consumption
- 11. Data from literature search Garcinia cambogia oral consumption
- 12. Human data Garcinia cambogia oral consumption
- 13. Hepatotoxic case reports Garcinia cambogia oral consumption
- 14. Animal studies Garcinia cambogia oral consumption
- 15. Exposure Garcinia cambogia oral consumption
- 16. Uncertainties Garcinia cambogia oral consumption
- 17. Risk Characterisation Garcinia cambogia oral consumption
- 18. Questions to the Committee Garcinia cambogia oral consumption
- 19. Abbreviations Garcinia cambogia oral consumption
- 20. References Garcinia cambogia oral consumption

This is a discussion paper. It does not reflect the views of the Committee. It should not be cited.

93. In 2019, the Scientific Committee of the Spanish Agency for Food Safety and Nutrition (AESAN) reviewed the risk associated with the consumption of food supplements that contain *G. gummi-gutta*.

- 94. They state that it is recommended to not exceed a daily dose of 3,000 mg of standardised extract at 50-60% HCA (dose equivalent to 1,500 1,800 mg of HCA), administered orally in three "shots", 30-60 minutes before three main meals.
- 95. It was found that different supplements marked in Spain containing *Garcinia* and/or HCA were in the form of tablets, capsules, sachets or vials with differing compositions of *G. cambogia* and/or HCA content. The range of recommended daily allowances in these products was 30 2,070 mg of HCA.
- AESAN noted that consumption of supplements containing *Garcinia* and/or HCA has been linked to toxic effects including hepatotoxicity, rhabdomyolysis, nephropathy, cardiovascular toxicity, hypomania and serotonin toxicity and psychosis; however, it has not been confirmed if *Garcinia* is the main toxicant, as it is often found in products with other ingredients.
- 97. AESAN concluded that there was "sufficient clinical evidence to establish a causal association between the consumption of garcinia and the duration of treatment, and the development of acute liver injury, with an obvious improvement in liver function after the withdrawal from the garcinia food supplement." This conclusion was based on the literature reviewed by AESAN including Lunsford et al., (2016) (see paragraphs 153-155), Crescioli et al., (2018) (see paragraphs 143-149, 181), and Sharma et al., (2018) (see paragraphs 150-152), among others.
- 98. Further to this, AESAN highlighted the importance of regulatory authorities to develop systems for post-market surveillance and for healthcare professionals, researchers and citizens to report adverse effects.