

# Previous COT evaluation

## In this guide

### [In this guide](#)

1. [TOX/2025/46 Annex 1 - Executive Summary](#)
2. [TOX/2025/46 Annex 1 - Background and scope of discussion](#)
3. [TOX/2025/46 Annex 1 - Properties of boron and sources in drinking water](#)
4. [TOX/2025/46 Annex 1 - Toxicity data for boron](#)
5. [TOX/2025/46 Annex 1 - Previous COT evaluation](#)
6. [TOX/2025/46 Annex 1 - Previous COT evaluation - Evaluations by other authoritative bodies](#)
7. [TOX/2025/46 Annex 1 - Discussion](#)
8. [TOX/2025/46 Annex 1 - Overall Conclusion](#)
9. [TOX/2025/46 Annex 1 - List of abbreviations](#)
10. [TOX/2025/46 Annex 1 - References](#)

**This is a paper for discussion. This does not represent the views of the Committee and should not be cited.**

58. Following publication of a new WHO drinking water guideline value in 1993, which was lower than the UK regulatory limit for drinking water at the time, the COT was asked to consider the toxicity data. In 1994, the COT provisionally agreed with the WHO (1993) NOAEL of 8.8 mg/kg bw/day for boron and acknowledged that the testis was a target organ for boron toxicity. However, the COT requested further information before finalising its conclusions.

59. In 1995, the COT considered a review of borates by the European Centre for Ecotoxicology and Toxicology of Chemicals (ECETOC) and a US National Toxicology Program (NTP) follow up to the Heindel et al., 1992 developmental toxicity study in rats (that would later be published by Price et al., 1996). The COT noted reported reduction in relative testis weight (without histological changes) following a dose of 4.4 mg/kg bw/day of boron in a 90-day oral study of boric acid in dogs (Weir and Fisher 1972). However, this was not seen in a 2-year study in dogs at higher doses where no convincing evidence of testicular toxicity was

observed at boron doses of 8 to 9 mg/kg bw/day (Weir and Fisher 1972). The COT considered that the dog studies reported by Weir and Fisher 1972 were unreliable due to small animal numbers in the experimental groups and inadequacies in the histopathological descriptions. The COT agreed that the NTP follow up developmental study in rats was of good quality and demonstrated a NOAEL of 9.6 mg born/kg bw/day for adverse effects of boron. Overall, the COT concluded that a NOAEL of around 10 mg/kg bw/day could be set based on the rat developmental studies.