

## Table 11

### In this guide

#### [In this guide](#)

1. [Annex A: Supplementary information - Table 1](#)
2. [Annex A: Supplementary information - Table 2](#)
3. [Annex A: Supplementary information - Table 3](#)
4. [Annex A: Supplementary information - Table 4](#)
5. [Annex A: Supplementary information - Table 5](#)
6. [Annex A: Supplementary information - Table 6](#)
7. [Annex A: Supplementary information - Table 7](#)
8. [Annex A: Supplementary information - Table 8](#)
9. [Annex A: Supplementary information - Table 9](#)
10. [Annex A: Supplementary information - Table 10](#)
11. [Annex A: Supplementary information - Table 11](#)
12. [Annex A: Supplementary information - Table 12](#)
13. [Annex A: Supplementary information - Table 13](#)
14. [Annex A: Supplementary information - Table 14](#)
15. [Annex A: Supplementary information - Table 15](#)
16. [Annex A: Supplementary information - Table 16](#)

**Table 11. Chronic exposure to sum of T-2 and HT-2 from RTE foods from DNSIYC and NDNS years 1-11 consumption data and occurrence data collected from the FSA call.**

**Infants (4-18 months) (n=2683)**

Food groups	Count of T-2		T-2 Concentration $\mu\text{g}/\text{kg}$ (mean)	T-2 Concentration $\mu\text{g}/\text{kg}$ (max)	Consumers (n)	Mean*; Mean*; Max*; Ma			
	HT-2 data call samples (n)	T-2				mean	P97.5	mean	P9
Infants' cereals	2		110.6	219	1128	0.36	1.5	0.71	2.9

### Toddlers (1.5-3 years) (n=1157)

Food groups	Count of T-2		T-2 Concentration $\mu\text{g}/\text{kg}$ (mean)	T-2 Concentration $\mu\text{g}/\text{kg}$ (max)	Consumers (n)	Mean*; Mean*; Max*; Ma			
	HT-2 data call samples (n)	T-2				mean	P97.5	mean	P9
Infants' cereals	2		110.6	219	34	0.22	0.70	0.44	1.4

Exposure estimates have been rounded to 2 significant figures; concentrations have been rounded to one decimal place.

\*Mean/Max concentration.