

# **Second draft statement on the risk assessment of cow's milk in children aged 1 to 5 years, in the context of plant-based drinks evaluations**

**This is a paper for discussion.**

**This does not represent the views of the Committee and should not be cited.**

## **Introduction**

1. The draft statement attached at Annex 1 to this paper consists of risk assessments of contaminants and chemicals within cow's milk which were reviewed as part of the work being conducted by the SACN/COT joint working group on plant based drinks. This will allow a comparison to be made between milk and the chemical risk assessment performed for plant based drinks performed in 2021 (COT, 2021a) This will facilitate the use of milk as a comparator against plant based drinks. It contains the risk characterisation and where applicable exposure assessments for a range of chemicals at the levels identified within cows' milk for children aged 6 months to 5 years.
2. This draft statement has been amended to reflect Members' comments after being discussed by the Committee at the February meeting. This included:
  - I. Clarification within the sections of veterinary medicines and pesticides sections involving good veterinary and agricultural practice, included within the main statement.
  - II. Corrections to inconsistencies between the text and summary tables for BPA, endogenous oestrogens and isoflavones, included in the main statement.

- III. Addition of a statement in the main text that the COT are currently reevaluating the safety of dioxins. Rewording of the conclusions for polycyclic aromatic hydrocarbons has been included in the main statement and Annex A.
- IV. Elaboration on the contributions of cow's milk to lead and cadmium exposure, included within the main text.
- V. Addition of a statement that the exposures discussed in this assessment may not cover individuals who consume large amounts of milk for protein supplementation, included in the main text.
- VI. Alterations to the conclusions for mercury that the Committee does not consider levels of milk in mercury to be of concern, included within the main statement.
- VII. Inclusion of EFSA's HBGV's within the exposure assessment of iodine, included in the main statement.
- VIII. Mention of the COT's 2000 statement on iodine in cow's milk before the COT's later 2017 statement, included within the main statement.
- IX. Clarification of the nature of exceedances of the health based guidance values for iodine, included in the main statement.
- X. A discussion of the natural levels of IGF-1 within milk, included in Annex A.
- XI. Editing of the introductory paragraph for oestrogens included in the main statement.
- XII. Alterations to the content within the endogenous oestrogens section to reaffirm that the COT does not consider 17 $\beta$ -oestradiol to be a direct genotoxin, included in the main statement and Annex A.
- XIII. Inclusion of a comparison of natural levels of circulating oestrogens and levels within milk, this has been included as a comparison between production rates of oestrogens within humans and exposures to oestrogens from cow's milk.
- XIV. An inclusion of a piece of text in the main statement discussing the conclusions for other mycotoxins besides aflatoxins.
- XV. Inclusion of a comment that in the cases where AFB1 has been found in milk these were outside of the EU and due to high levels of feed contamination. And this is unlikely to occur within the EU / UK.

XVI. Alterations to the captions for the tables and text displaying and describing EFSA's assessment for aflatoxins to emphasise the displayed MOEs are derived from total dietary exposure, included in the main statement.

XVII. Alteration of the phrase 'no risk of harm' to 'no risk to health' within the aflatoxins concluding paragraph.

XVIII. Inclusion of the conclusions of Kowalczyk et al., (2013) found in the Journal of Agricultural Food Chemistry and Hill, Becanova and Lohmann, (2021) found in the journal Analytical and Bioanalytical chemistry within the main statement.

XIX. A rephrasing of the conclusions for PBDEs and TBPPA within the main statement.

XX. Alteration of inconsistencies between the microplastics text and summary tables, in addition to alterations to the microplastics text, changes were made in the main statement and Annex A.

3. The majority of chemicals did not present any risk to health to children aged 6 months to 5 years at the levels of exposure identified within cows' milk.

4. Iodine was identified as being of low concern. This is in line with previous COT statements in 2000 and 2017 as being due to the close proximity of recommended dietary levels and HBGVs, uncertainty in HBGV calculation and the small nature of the HBGV exceedance.

5. Aflatoxin M1 was identified as being of low concern. There is a low concern for total aflatoxins within milk, however, this is driven by levels of AFM1 within cows' milk.

6. Risk relating to isoflavones is uncertain due to a lack of HBGVs for young children, therefore the significance of the occurrence values in milk. Any risk however is lower than that in soya milk which contains higher levels of isoflavones.

7. The draft statement is attached as Annex 1 to this paper. In order to ensure the statement is not excessively long, detailed supporting information for chemicals deemed to have minimal risk is attached as Annex A to the main statement.

## **Questions for the Committee**

8. The Committee are asked to consider:
- a) Does the Committee have any comments on the general structure and content of this statement and its Annex?
  - b) Does the Committee concur with the conclusions presented within this statement?
  - c) Does the Committee have any other comments on this statement or its Annex?

**Secretariat**  
**March 2022.**

## **Annex 1 to TOX/2022/04**

### **Background**

1. Plant-based drinks have become increasingly popular in the United Kingdom (UK) both for individuals with an allergy to cows' milk or lactose intolerance and those who wish to avoid dairy products for other ethical or cultural reasons.
2. Current UK Government advice regarding the use of plant-based drinks for infants and young children is that unsweetened calcium-fortified plant-based drinks, such as soya, oat and almond drinks, can be given to children from the age of 12 months as part of a healthy balanced diet; rice drinks should not be given due to the levels of arsenic in these products (NHS, 2018). The Committee reviewed three of the drinks, with a statement being published last year at the request of the Department of Health and Social Care (DHSC) (COT, 2021a). The Scientific Advisory Committee on Nutrition (SACN) have also been considering the nutritional aspects of plant based drinks and in order to bring together the nutritional and chemical risk assessments of these drinks, a joint working group of SACN and the Committee on Toxicity of Chemicals in Food, Consumer Products and the Environment (COT) has been established.
3. DHSC is in the process of conducting an Equalities Analysis covering both the Nursery Milk Scheme and the Healthy Start Scheme which considers equalities issues posed by the current legislation as it pertains both to plant-based drinks, and also to animal milks other than cow's milk. DHSC is keen to ensure that this Equalities Analysis reflects the most up-to-date advice on safety and toxicity issues from COT, and on nutritional issues from the SACN. However, this process is currently on hold whilst the joint Working Group considers plant-based drinks.

4. The Committee was asked to consider the potential for adverse effects arising from the consumption of plant drinks by young children (aged 6 months- 5 years) who were following a plant-based diet. The drinks considered were soya, oat and almond; rice drinks were not reviewed since there is existing advice that these should not be given to young children due to their arsenic content. The three drinks were selected as they were the most popular alternatives to dairy at the time of review. The statement setting out the views and conclusions of the Committee was published in January 2021 (COT, 2021a).

5. The Committee agreed during their meeting in July 2021 that the main comparator for plant-based drinks should be cow's milk and that a discussion paper should be produced looking at the potential chemical risks in the consumption of this over the identical population group of interest, children aged 6 months to 5 years

6. Most of the fresh cow's milk available in the UK is UK derived, thus the risks and relevant chemical exposures for this paper are European Union (EU) or UK focused and it is assumed that EU farming practices are similar to the UK.

7. This statement follows two discussion papers presented over the course of 2021 (TOX/2021/53 and TOX/2021/58) which presented exposure assessments and subsequent risk characterisation for a large range of chemical compounds that could potentially occur in milk. The compounds discussed were:

Part 1 (TOX/2021/53):

- I. Veterinary medicines
- II. Pesticides
- III. Nitrate and Nitrite
- IV. Bisphenol A (BPA)
- V. Phthalates
- VI. Dioxins and Dioxin-Like Polychlorinated Biphenyls (DL-PCBs)
- VII. Non-Dioxin-Like Polychlorinated Biphenyls (NDL-PCBs)
- VIII. Polycyclic Aromatic Hydrocarbons (PAHs)
- IX. Isoflavones: Genistein (GEN), Daidzein (DAI), Equol (EQU, metabolite of DAI), Formononetin (FOR) and Biochanin A (BIO)

Part 2 (TOX/2021/58):

- X. Heavy metals: Lead (Pb), Arsenic (As), Mercury (Hg) and Cadmium (Cd)
- XI. Iodine

- XII. Perchlorate and Chlorate
- XIII. Mycotoxins: Aflatoxins (AFB1 and AFM1) and others including Deoxynivalenol (DON)
- XIV. Hormones – Oestrogens, Insulin-Like Growth Factor 1 (IGF-1)
- XV. Per- and polyfluoroalkyl substances (PFAS)
- XVI. Brominated Flame Retardants (BFRs)
- XVII. Microplastics

8. The Committee considered the compounds in cows' milk to be of minimal risk where the evidence suggested that there was no exceedance of health based guidance values from consumption of cow's milk. In these cases supplementary information, including the discussion of health-based guidance values (HBGVs), detailed exposure assessments and, where relevant, risk characterisation are included in Annex A to this statement.

9. It is acknowledged from scrutiny of the historical EU RASFF (Rapid Alert System for Food and Feed) data and FSA's alert tools that occasional incidents of contamination of cow's milk with chemicals not included in the discussion papers have occurred; this has involved chemicals such as mineral oils (Montgomery, Haughey and Elliott, 2020), other plant toxins from feed contamination, other agricultural contaminants (e.g. urease inhibitors) (Byrne et al., 2020) and other industrial contaminants (e.g. Parabens). As 'one-off' incidents these are acknowledged but not discussed or evaluated in this statement as the overall risks are deemed minimal.

10. Members discussed comparing the levels of particular contaminants within selected plant-based drinks and cow's milk. Due to the independent nature of the cow's milk assessment, compounds present in cows' milk may not be present at significant levels in plant-based drinks and vice versa. Where compounds are shared between drink types, comparisons have been made.

## **Consumption data**

11. The National Diet and Nutrition Survey (NDNS) rolling programme and Diet and Nutrition Survey of Infants and Young Children (DNSIYC) data were used to undertake any chronic exposure assessments in this statement, required for assessing the safety of milk from a chemical contaminant perspective, in young children aged 6 months to 5 years (Department of Health, 2011; Bates et al., 2014; Roberts et al., 2018). The data presented in Table 1 includes consumption data for cow's milk consumed as a drink and when used in recipes. Consumption

data for children aged 6 – 12 months are derived from milk used in recipes only, as cow’s milk is not recommended by the NHS as a main drink for infants in this age range (NHS, 2018). Table 2 presents consumption data for milk as a drink only. As these values are only slightly lower, all exposure assessments have been undertaken using the highest consumption estimates from Table 1 only.

Table 1. Estimated chronic consumption of cow’s milk in consumers (as a drink and with recipes)

| Age (months) | Number of Consumers | (g/person/day) |                   | (g/kg bw/day) |                   |
|--------------|---------------------|----------------|-------------------|---------------|-------------------|
|              |                     | Mean           | 97.5th percentile | Mean          | 97.5th percentile |
| 6 - <12      | 1257                | 120            | 460               | 13            | 48                |
| 12 - <18     | 1275                | 350            | 790               | 32            | 75                |
| 18 - <24     | 157                 | 350            | 840               | 29            | 79                |
| 24 - <48     | 351                 | 320            | 770               | 23            | 59                |
| 48 - <60     | 618                 | 290            | 780               | 17            | 46                |

Table 2. Estimated chronic consumption of cow’s milk in consumers (as a drink without recipes)

| Age (months) | Number of Consumers | (g/kg bw/day) Mean | (g/kg bw/day) 97.5th percentile |
|--------------|---------------------|--------------------|---------------------------------|
| 12 - <18     | 1148                | 30                 | 71                              |
| 18 - <24     | 147                 | 28                 | 73                              |

|          |     |    |    |
|----------|-----|----|----|
| 24 - <48 | 337 | 21 | 54 |
| 48 - <60 | 585 | 15 | 42 |

12. Exposure assessments utilising this data cover the general population at both mean and high levels of consumption and may not include individuals who drink large amounts of milk daily for reasons such as protein supplementation.

## **Chemicals assessed**

### **Veterinary medicines**

13. Veterinary medicines, for example antibiotics, are used in animal husbandry to alleviate suffering and disease. UK farmers should follow the Veterinary Medicines Directorate (VMD) recommended guidance on responsible use (VMD, 2014). These include accurate record keeping, purchasing from authorised sources, correct administration and observing relevant withdrawal periods (the length of time any subsequent animal products must not enter the food chain) after administration.

14. Veterinary medicines can be present below the maximum residue limits (MRLs) as a result of good veterinary practice and this does not constitute a risk to health. However, they can on occasion be present in animal derived products above these MRLs when procedures are not followed correctly. Cow's milk is routinely monitored through ongoing surveys with the UK National Reference Laboratory (NRL).

15. Between 2015 and the end of 2020, 21,574 analyses of cow's milk samples were undertaken as part of the VMD survey covering, anthelmintics, avermectins, cephalosporins, chloramphenicol, dapsone, florfenicol, Non-Steroidal Anti-inflammatory drugs (NSAIDS), and other antimicrobials (as a screening method) (VMD, 2015, 2016, 2017, 2018, 2019, 2020). From the analysis over this 6-year period only 0.12% (24) returned a positive result. Positive results were considered instances where medicines were above the maximum residue limit (MRL) for milk. From these only two residues, penicillin G and triclabendazole both in 2017, resulted in a subsequent risk assessment concluding the milk samples represented a potential food safety risk to the consumer, and this was before taking any dilution effect into account, e.g. from bulk tanks at dairies.

16. Based on the last 6 years UK statutory survey the COT concluded that it appears that the risk of veterinary medicine exposure after isolated incidents from drinking cow's milk is negligible.

## **Pesticides**

17. Pesticides primarily enter the dairy food chain via consumption of contaminated feed or water by cattle. They are routinely monitored through ongoing statutory surveillance with the UK National Reference Laboratory. Compounds detected below the regulated limits due to good agricultural practice are deemed to have a minimal safety risk, However, when above these levels they can potentially present a risk to health.

18. Between 2015 and the end of 2020, 1,723 cow's milk samples were analysed and reported by The Expert Committee on Pesticide Residues in Food (PRiF), 2015, 2016, 2017, 2018, 2019, 2020). From all the samples analysed over this 6-year period only 1 returned a positive result above the Maximum Residue Limit (MRL). This residue, in 2019, was a persistent quaternary ammonium compound at 0.3 mg/kg, likely a contaminant from a cleaning product.

19. Based on the last 6 UK statutory survey results the COT concluded that the risk of pesticide exposure from drinking cow's milk is negligible.

## **Nitrate and nitrite**

20. Nitrate and nitrite are naturally occurring chemicals that form part of the nitrogen cycle. They act as oxidising agents that can cause methemoglobinemia in animals and humans after high consumption. They occur naturally in vegetables but are also used widely as meat preservatives, in agricultural waste streams e.g. from fertiliser use, and as chemical contaminants from industrial processes and materials.

21. Nitrates are widely consumed by animals and humans, although nitrite is regulated as an undesirable substance in animal feed (EU 574/ 2011). In animals the largest potential exposure of nitrite is from the in-vivo transformation of nitrate to nitrite. Feed and contaminated water can have high levels of nitrate and represent the main contributor to nitrite exposure for food-producing animals (Cockburn et al., 2013).

22. An exposure assessment has been undertaken for nitrate within Annex A using UK consumption data (Table 1 above). This is presented alongside a

discussion of EFSA's 2009 opinion on nitrite. Nitrate exposure was found to be below 1% of the ADI. EFSA's 2009 opinion concluded that nitrite is present at extremely low levels in fresh animal products and therefore not of human health concern (EFSA, 2009).

23. In light of the very low percentages of the recommended ADI for nitrate that would occur through consumption of cow's milk in young children, along with the EFSA (2009) opinion's conclusion, the COT concluded that nitrite and nitrate contamination pose a minimal risk in the daily consumption of cow's milk.

## **Bisphenol A**

24. Bisphenol A (BPA) is a compound used as a monomer in the production of many plastics and resins, particularly polycarbonate materials employed in the manufacture of food contact materials and food storage containers such as cans. It is known to potentially migrate from plastic containers, or resins from coatings, into food and drinks. It is also widely used in the production of non-food related products such as surface coatings, resin-based paints, flame retardants and medical devices. For cow's milk, potential BPA contamination may come from the mechanical milking apparatus and subsequent storage vessels in the dairy chain such as cooling tanks.

25. BPA is an endocrine disrupter in that it potentially interferes with the regulation of hormones in the endocrine system. It is therefore assumed to have toxic effects on metabolism, growth, sexual development, stress response, insulin production, gender behaviour, reproduction, and foetal development (Cirillo et al., 2015). It is also considered a contributing factor in the onset of metabolic disorders, including diabetes and obesity, and immune dysfunction (Bansal, Henao-Mejia and Simmons, 2018).

26. EFSA's 2015 opinion on BPA, discussed in Annex A, advises a reduced temporary tolerable daily intake (t-TDI) and concluded that BPA poses no health concern for consumers at any age group (EFSA, 2015b).

27. In 2019, COT was asked to review the risk of toxicity of chemicals in the diets of infants and young children aged 0-5 years, in support of a review by SACN of Government recommendations on complementary and young child feeding (COT, 2019c, 2020) and BPA was considered as part of that. For BPA, COT's current position is that they are awaiting EFSA's new updated scientific opinion (currently ongoing) to conclude if a new COT evaluation is required. Based on the 2015 opinion, the COT do not currently consider that BPA within cows' milk

presents a risk to health for children aged 6 months to 5 years of age.

## **Phthalates**

28. Phthalates are esters of the aromatic dicarboxylic acid phthalic acid that have a long history of use as additives to plastics to improve their flexibility but also have wide applicability across industry, for example in pharmaceutical coatings, paints, cosmetics and food contact materials.

29. Phthalates do not form covalent bonds with the material into which they are incorporated, therefore can readily migrate into food from packaging materials. The extensive and historic use of phthalates has led to their being widely distributed in the environment and the food chain. The general population is exposed to phthalates via food (including migration from food contact materials) and drinking water, but also through inhalation and dermal exposure (Heudorf, Mersch-Sundermann and Angerer, 2007).

30. In 2005, EFSA performed risk assessments on a small range of the most widely used phthalates, namely, di-butylphthalate (DBP), butyl-benzyl-phthalate (BBP), bis(2-ethylhexyl)phthalate (DEHP), di-isononylphthalate (DINP) and diisodecylphthalate (DIDP) and derived TDIs for them (EFSA, 2005b, 2005c, 2005d, 2005e, 2005f).

31. In Annex A, EFSA's 2005 and 2019 risk assessments of phthalates are discussed. In 2019 the group phthalates (DEHP+ DBP+ BBP+ DINP expressed as DEHP equivalents) contributed up to 14% of the recommended group TDI whilst for 95th percentile consumers a maximum of 23% (EFSA, 2019). For DIDP both mean and 95th percentile consumers were exposed to well below the TDI.

32. In May 2011, COT produced a statement COT, (2011) on dietary exposure to the phthalates DBP, BBP, DEHP, DINP, DIDP and diethyl phthalate (DEP) using data from the UK Total Diet Study (TDS), and concluded that the levels of phthalates that were found in samples from the 2007 TDS did not indicate a risk to human health from dietary exposure, either when the compounds were assessed alone or in combination.

33. In the recent COT review of EFSA's public consultation on the EFSA Opinion "Draft update of the risk assessment of dibutylphthalate (DBP), butyl-benzyl-phthalate (BBP), bis(2-ethylhexyl)phthalate (DEHP), di-isononylphthalate (DINP) and diisodecylphthalate (DIDP) for use in food contact materials", the COT were content that for DBP, BBP, DEHP and DINP the exposures estimated by EFSA did

not indicate a health concern using the group TDI (COT, 2019).

34. From this information the COT concluded that phthalates within cows' milk do not present a risk to health for children aged 6 months to 5 years of age.

### **Dioxins and Dioxin-Like polychlorinated biphenyls (DL-PCBs)**

35. Formed as by-products of a number of industrial processes, polychlorinated dibenzodioxins (PCDDs) and polychlorinated dibenzofurans (PCDFs) are two groups of tricyclic planar compounds that are formed by combustion of organochlorine compounds or of non-chlorine compounds in the presence of chlorine. Of these, 75 PCDD and 135 PCDF "congeners" are known, with structures varying in the number of chlorine atoms and their positions in the rings. Only 17 of these are relatively persistent in animals and humans and therefore considered relevant (EFSA, 2018).

36. HBGVs have been generated by multiple authorities and these are discussed within Annex A.

37. An exposure assessment has been undertaken for cow's milk using consumption data from Table 1 and is presented within Annex A using occurrence levels from EFSA's 2018 opinion paper (EFSA, 2018), compared against the recommended TDI of 2 pg WHO-TEQ/kg bw per day from COT in 2001 (COT, 2001). Utilising the upper bound (UB) mean occurrence levels led to exceedances of the TDI in two age groups. Factors including the worst-case assumption of a 3.5% fat content of milk and using the upper bound of the mean occurrence concentrations suggest that realistic exposure will be below the levels estimated in this exposure assessment.

38. At the 95th percentile occurrence value, exceedances of the TDI occurred for both mean and high level consumers, however, this scenario is considered to be highly conservative and unrealistic.

39. As noted in the recent COT review of chemicals in the diets of infants and young children, the COT is currently reviewing the current guidance values for dioxins and dioxin like PCBs however does not consider it necessary to update their advice until this work has been completed (COT, 2021c).

40. The COT has concluded from the exposure assessments conducted in annex A that dioxins within cows' milk do not currently present a risk to health for children aged 6 months to 5 years of age.

## **Non-dioxin-like PCBs**

41. Some PCBs do not share the same toxic endpoints as the dioxins and have different effects, for example oestrogenic and anti-oestrogenic effects, and are therefore regarded as a separate group of persistent organic chemicals that are present in the environment and food.

42. Dietary exposure assessments by EFSA, (2005a) and JECFA, (2016) are discussed within Annex A, these surveys suggest that dietary exposure is within safe levels for young children,

43. The COT concluded, based on the above evidence, that there was no risk to health from NDL-PCBs within cows' milk for children aged 6 months to 5 years of age.

## **Polycyclic Aromatic Hydrocarbons (PAHs)**

44. PAHs (polycyclic aromatic hydrocarbons) are organic compounds characterised by the presence of 2 or more fused aromatic rings, many of which are known carcinogens. Although naphthalene, with 2 fused rings, would technically be part of this group of compounds it is usually not regarded as a member. PAHs are common products of combustion and are widely distributed in the environment as the result of vehicle exhaust and industrial processes and in the diet in cooked food and cooking by-products such as oils vaporised from frying pans and smoke from barbecues. Production of PAHs by cooking is greater when fat expressed from the food drips directly onto the heating element or hot coals.

45. An exposure assessment for benzo[a]pyrene (BaP) and a separate assessment for PAH4 (sum of BaP, benz[a]anthracene (BaA), benzo[b]fluoranthene (BbF) and chrysene (ChR), was undertaken. These are presented within Annex A utilising consumption data in Table 1 and the UK TDS from 2012 (Fernandes et al., 2012).

46. The subsequent MOEs presented for the exposure to PAH4 are all above 10,000 for both average and high-level consumers across all age ranges of young children. These high MOE's indicate there is no concern to health from these chemicals from drinking cow's milk.

47. In the recent COT review with SACN on the risk of toxicity of chemicals in the diets of infants and young children, the COT concluded the intakes of PAHs

(BaP and PAH4) from human breast milk and food are of no concern to health in children aged 12 to 60 months (COT, 2020).

## **Isoflavones**

48. Phytoestrogens are chemicals of plant origin that have been shown to influence biological processes mainly through their structural similarities to oestrogens, and their ability to bind to oestrogen receptors (ERs). They can therefore potentially cause unfavourable effects such as disruptions in sexual behaviour and brain sexual differentiation, changes in hormone levels, and increases in breast cancer risk (Xiao, 2008; Socas-Rodríguez et al., 2015). The largest group of phytoestrogens are flavonoids, which can be further divided into three subclasses, coumestans, prenylated flavonoids and isoflavones.

49. Isoflavones can be found in many plants, including barley, sunflower, clover, lentils, alfalfa sprout, broccoli and cauliflower. However, the richest sources of isoflavones in the human diet are foods and dietary supplements made from soya bean and soya protein (McCarver et al., 2011). Soya isoflavones in foods occur mainly as carbohydrate conjugates (glycosides), the major group being the glucose conjugates (glucosides), e.g. genistein (GEN) and daidzein (DAI). The other most commonly considered isoflavones include formononetin (FOR), biochanin A (BIO) and a metabolite of DAI, equol (EQU).

50. The phenolic and hydroxyl moieties (and the distance between them) are key structural similarities between isoflavones and  $17\beta$ -oestradiol which allow them to bind to ERs. Numerous studies have indicated that GEN is the isoflavone with the greatest oestrogenic activity (McCarver et al., 2011).

51. Animal studies performed before 2003 indicated that intake of isoflavones in early life can produce oestrogenic effects, affect thyroid function, alter protein concentrations and structures in the brain, and alter some parameters of immune function, as well as sexual development in older animals. Although some animal studies indicated possible risks to humans, overall, the results of animal studies were inconsistent. The COT 2003 report noted that human data were limited, and that most of the relevant scientific information was derived from experimental studies in animals, mainly rodents. The extrapolation of such studies to humans was difficult because of inter-species differences in ADME (absorption, distribution, metabolism, and excretion), sexual development and reproductive function, and the use of relatively high doses or non-oral routes of administration.

52. In vitro experiments reviewed in the 2003 COT report (COT, 2003) showed that phytoestrogens could modulate the levels of sex hormone binding globulin (SHBG), inhibit enzymes involved in oestrogen biosynthesis and metabolism to modulate concentrations of endogenous oestrogens, and inhibit thyroid peroxidase activity to reduce the concentrations of thyroid hormones. GEN was found to interact with topoisomerase II and protein kinases (enzymes involved in cellular proliferation and differentiation) and to inhibit human T-cell proliferation and interleukin-2 production.

53. The 2003 COT report concludes that it is not possible to propose HBGVs for infants (COT, 2003). Reasons for this include the difficulty in extrapolation from animals to humans because of differences in toxicokinetics, uncertainty about differences between adults and infants (particularly those arising from development of the gut microflora), and the lack of dose-response data and the possibility of bias and chance effects in the available human studies. In a more recent 2013 COT report COT, (2013a) assessing literature since 2003, the same conclusions were drawn, in that it is not possible to propose HBGVs due to limitations in the available data.

54. Other health authorities have proposed HBGVs such as the Nordic Council in 2020 (Nordic Council of Ministers, 2020). For children they proposed 'a rounded value of 0.07 mg/kg bw per day of GEN. This corresponds to 2.1 mg genistein per day for a person weighing 30 kg.' This value was derived from the Li et al., (2014) rat study taking the LOAEL of 20 mg/kg bw and applying a further uncertainty factor of 3 on top of the factors of 10 for inter species differences and intraspecies variation.

55. Isoflavones are known to be transferred to cow's milk after digestion of plant-based feed stuffs (Bláhová et al., 2016). Occurrence in the milk is dependent on the feed. Milk phytoestrogen concentration is strongly influenced by silage plant composition. Feed with either deliberate addition of, or inadvertently contaminated with red clover for example will have greatly increased concentrations of isoflavones (Höjer et al., 2012).

## **Risk Characterisation**

56. To obtain published concentrations for Isoflavones in cow's milk a literature search was undertaken using the keywords Isoflavone AND Cow AND Milk AND Risk in both PubMed and Science Direct. A large number of results with very varied isoflavone concentrations were returned from European countries. The UK

data only are summarised below (Table 3) from (Nørskov et al., 2019).

Table 3. Summary of mean isoflavone concentrations (all µg/kg) GEN, EQU, FOR and DAI from differing cow's milk types in the UK and a comparison with mean soya milk concentrations (µg/kg).

| Milk Type       | Number of samples | GEN  | EQU  | FOR  | DAI  | Sum   |
|-----------------|-------------------|------|------|------|------|-------|
| Conventional    | 48                | 0.83 | 63.6 | 0.08 | 0.95 | 67.7  |
| Organic         | 48                | 2.32 | 411  | 1.10 | 2.69 | 417   |
| Free range      | 24                | 0.85 | 66.4 | 0.09 | 0.96 | 70.4  |
| Low fat soya* 1 |                   | 875  | -    | -    | 567  | 1,442 |

57. As noted above, COT have not established a HBGV for isoflavones for young children and the significance of the concentrations summarised in Table 3 is uncertain. However, isoflavone concentrations from cow's milk are considerably lower than those from soya alternatives, suggesting that any associated risk will also be lower.

## Lead

58. Lead is a well-known heavy metal and pollutant which can cause multiple negative health effects in humans, its impact on the health of infants was evaluated by the COT, (2013) in their statement on the potential risks from lead in the infant diet and their addendum (COT, 2016a).

59. EFSA's 2012 opinion on lead and the COT's 2013 and 2016 statements on lead exposure in the diets of infants and children have been discussed (Annex A). Whilst exceedances of the benchmark dose lower confidence limit (BMDL)01 of 0.5 µg/kg bw/day were observed in EFSA's total dietary exposure estimate for infants aged <1 year (0.83 and 0.91 µg/kg bw/day in two surveys), toddlers aged 1-<3 years (1.32 µg/kg bw/day) and other children aged 3-<10 years (1.03 µg/kg bw/day), the contribution of cows' milks to total middle bound lead exposure did not exceed 2% for infants, 5% for toddlers and 4% for other children, therefore lead within cows' milk is not a concern (EFSA, 2012b). The COT's statement found

diet contributed little to lead exposure with other sources of exposure being the most significant (COT, 2013b, 2016a).

60. Based on the information provided in EFSA (2012b) and the evaluation by the COT in 2013 and 2016 the COT concludes that it is unlikely that lead in cow's milk would pose a risk to infants and children from the ages of 6 months to 5 years.

## **Arsenic**

61. Inorganic arsenic is the focus of this evaluation as it with the previous COT statement (COT, 2016b).

62. The COT's 2016 statement and EFSA's 2021 evaluation have been discussed in Annex A. The COT's 2016 risk assessment suggest that at mean levels of consumption, for infants aged 4 months to 5 years the MOE's were below 10, therefore a risk to health may exist from dietary exposure. However, in EFSA's recent 2021 evaluation cow's milk was shown to contain minimal amounts of iAs (EFSA, 2021a).

63. The COT concluded from the above information that inorganic arsenic in cows' milk does not present a risk to health to children aged 6 months to 5 years of age.

## **Mercury**

64. Mercury is a metal released from both anthropogenic and natural sources. It is found as elemental mercury (Hg<sup>0</sup>), inorganic mercury (mercurous and mercuric cations (Hg<sup>+</sup> and Hg<sup>2+</sup> respectively) and organic mercury. Methylmercury is the most abundant organic mercury compound in the food chain (COT, 2018c).

65. The toxicity of mercury varies depending on whether the mercury is in an organic or inorganic form. The focus of this paper is inorganic mercury as in EFSA's 2012 report it was assumed the majority of mercury within milk was inorganic in nature (EFSA, 2012c).

66. The COT's 2018 statement on methylmercury in the diets of infants and children and EFSA's 2012 opinion have been discussed in Annex A. EFSA did not consider total dietary exposure to inorganic mercury to be a risk for the European population. For all age groups, excepting toddlers, the TWI for inorganic mercury was not exceeded. Cows' milk contributed a maximum of 15% to this total

exposure. The COT in 2018 found no exceedances of the inorganic mercury TWI using either TDS or infant metals survey data for the assessment.

67. From the above information the COT concluded that there is no risk to health for infants and children aged 6 months – 5 years from exposure to inorganic mercury in cows' milk is low.

## **Cadmium**

68. Cadmium (Cd) is a soft, silver-white or blue-white metal existing in various mineral forms and is present throughout the environment. It is used in many processes such as electroplating, alloy production, paints and pigments and is found in a wide range of industrial and consumer products. Environmental cadmium concentrations are reflective of natural sources such as volcanic activity as well as anthropogenic sources for example non-ferrous metal smelting.

69. Exposure assessments performed by (EFSA, 2012a) and the COT have been discussed, this can be found within Annex A. Whilst exceedances of the TWI were observed in both (COT, 2018b) and EFSA, (2012a) exposure assessments the relative contribution of cow's milk in both of these assessments was low.

70. In the EFSA exposure assessment (EFSA (2012a) collected surveys were merged and weighted to reflect the years individuals spent within the age brackets from a 77 year average lifespan producing mean average upper bound lifetime exposures. For infants aged <1 year this was 3.50 µg/kg bw/week, for toddlers aged 3-< 10 years this was 5.9 µg/kg bw/week and for other children aged 3-<10 years this was 4.69 µg/kg bw/week. These all exceed the(EFSA, 2011c) EFSA 2011 TWI of 2.5 µg/kg bw/week. For infants liquid milk contributed 1.59% to total dietary cadmium exposures whilst for toddlers it contributed 1.78% and 2.28% for other children.

71. In COT (2018b) it was reported that exposures up to 260% of the EFSA 2011 TWI were calculated but cow's milk was not identified as one of the key contributing food groups.

72. The COT concluded from the above information that cadmium in cow's milk presents a low risk to the health of infants and children aged between 6 months and 5 years.

## **Iodine**

73. Iodine is an essential micronutrient necessary to produce thyroid hormones. The COT released a statement (COT, 2017b) discussing in depth the potential risks of excess iodine in the diets of infants and children aged 0-5 years. Milk is a considerable source of iodine in the diet, this may be due to fortification of animal feed with iodine compounds and teat dipping with sterilising compounds prior to milking.

74. Iodine excess is well tolerated by healthy individuals. For some it may cause hypothyroidism, hyperthyroidism, goitre and/or thyroid autoimmunity. Individuals with prior exposure to iodine deficiency or pre-existing thyroid disease may be more vulnerable to iodine excess induced thyroid disorders (Farebrother, Zimmermann and Andersson, 2019).

75. In 1989 the Joint Expert Committee on Food Additives (JECFA) established a provisional Maximum Tolerable Daily Intake (PMTDI) for iodine of 17 µg/kg bw/day from all sources, based on the same longer term studies in adults used by the European Scientific Committee on Food (SCF) in 2002 in support of their TUL, recorded in EFSA, (2006). No safety factors were used as these studies encompassed a relatively large number of subjects (FAO/WHO, 1989). This PMTDI was utilised to perform an exposure assessment in the COT's 2000 statement on iodine in cow's milk (COT, 2000).

76. The COT (2017b) stated "Excess iodine has considerably varied effects between individuals. The adult thyroid gland secretes about 80 µg thyroxine per day which requires a dietary intake of between 100 and 150 µg/day of iodine. Humans have a number of mechanisms by which they can counter an excess of iodine. These include the sodium-iodide symporter which blocks the transport of iodine into the thyroid cells and the Wolff-Chaikoff effect, more details of which can be found in the review by Bürgi, (2010). Most people can tolerate a chronic excess of iodine of up to 2 g of iodine per day but there will be some individuals who experience effects at much lower levels, close to the upper recommended limit for intake (Bürgi, 2010)."

77. In the COT's 2017 statement on the risks of excess iodine exposure to infants and young children they assessed three HBGVs, This assessment is paraphrased below.

78. The Expert Group on Vitamins and Minerals (EVM) looked in detail at the metabolism of iodine and the effects of excess iodine in 2003 (EVM, 2003). The EVM concluded that there were insufficient data to set a Safe Upper Level (SUL) for iodine. For guidance purposes, they indicated that a level of 0.5 mg/day of

supplemental iodine in addition to the background intake of 0.43 mg/day would be unlikely to cause adverse effects in adults based on slight alterations in serum thyroid hormone levels at supplemental doses of 0 – 2 mg/day in a range of human studies. From this data the EVM proceeded to set a guidance level for iodine at 15 µg/kg bw/day for adults. This value is utilised in an exposure assessment in this paper due to its conservative nature.

79. In 2002, the SCF published an opinion on the tolerable upper intake levels of vitamins and minerals, recorded in EFSA, (2006). For iodine, they set a tolerable upper level (TUL) of 600 µg/day for adults, reduced on a body surface area (body weight<sup>0.75</sup>) basis for children to 200 µg/day for ages 1-3 years and 250 µg/day for ages 4-6 years. This TUL was based on dose-response studies of short duration in humans, which showed changes in serum thyroid hormone levels at dose levels of 1800 µg/day and was supported by longer term studies with approximately similar doses that did not show adverse effects, but lacked detailed iodine intake data. An uncertainty factor of 3 was used. These values were endorsed by EFSA (2006).

80. In 2017 the COT calculated HBGVs based on the EFSA (2006) endorsed TULs in their statement assessing the risks of excess iodine in the diet of infants and young children. This used differing TULs for separate age groups based on different mean bodyweights. This approach has been followed below in Table 4 using mean bodyweight data supplied by the FSA’s exposure assessment team to produce derivative TULs for the selected age ranges (EFSA, 2006; Department of Health, 2013; Bates et al., 2014, 2016, 2020; Roberts et al., 2018).

Table 4: Table displaying the age adjusted TULs from EFSA 2006, mean bodyweight for each age group supplied from NDNS data and the TULs derived from this data (EFSA, 2006; Department of Health, 2013; Bates et al., 2014, 2016, 2020; Roberts et al., 2018).

| Age group                  | 0-<12 months  | 12-<18 months | 18-<24 months | 24-<48 months | 48 -<60 months |
|----------------------------|---|---------------|---------------|---------------|----------------|
| EFSA adjusted TUL (µg/day) | No tolerable upper limit (TUL) specified for this group | 200           | 200           | 200           | 250            |

|                                  |     |      |      |      |      |
|----------------------------------|-----|------|------|------|------|
| Average<br>bodyweight (kg<br>bw) | N/A | 10.9 | 12.2 | 15.3 | 19.4 |
| Derived TUL<br>µg/kg bw/day      | N/A | 18.4 | 16.4 | 13.1 | 12.9 |

## Exposure Assessment and Risk Characterisation

81. The 2016 infant metals survey provided comprehensive occurrence information for iodine in UK milk. Iodine was found to be present at a mean level of 271 µg/kg (FSA, 2016).

82. In addition to the infant metal survey, occurrence levels were found through an interrogation of the PubMed database using the terms “iodine AND cows AND milk” and “ iodine AND excess AND milk” with search results limited to 2001-2021.

83. A review article by Reijden et al., collated iodine occurrence data from 30 European and 1 United States (US) study including 2 from the UK in 2012 and 2016 (Reijden, Zimmermann and Galetti, 2017). The 2012 UK study presented a median iodine level in conventional milk at 250 µg/kg from 80 samples whilst the 2016 study presented a mean value of 458 µg/kg from 24 samples (Bath, Button and Rayman, 2012; Payling et al., 2015).

84. Bath et al (2017) also documented iodine at median levels of 438 µg/kg in conventional (non-organic) milk. Sample numbers were restricted to 5 samples, taken at a single time in winter, due to the seasonal variation in iodine levels this may have increased levels of iodine in samples as winter milk is often recorded as having higher iodine levels (Bath et al., 2017; Reijden, Zimmermann and Galetti, 2017).

85. A study by O’Kane et al. investigating seasonal variation in iodine and selenium concentration in milk found mean ( $\pm$  SD) (standard deviation) iodine levels were 475.9 ( $\pm$  63.5) µg/kg in pasteurised Northern Irish milk (O’Kane et al., 2018). This mean was obtained from the analysis of 36 samples. 95th percentile or maximum occurrence data were not presented in this study. The highest recorded mean concentration was 543.3 ( $\pm$  53.7) µg/kg from 9 samples of milk collected in spring.

86. The highest UK mean iodine concentration reported was found in O’Kane et al. (475.9 µg/kg). Using this occurrence data, the JECFA 1989 PMTDI of 17 µg/kg bw/day and consumption rates in Table 1 an exposure assessment was undertaken which is presented in Table 5.

Table 5: Exposure assessment from cows’ milk consumption using mean iodine occurrence in O’Kane et al. (2018), consumption data from the NDNS (Table 1) and JECFA’s PMTDI (FAO/WHO, 1989).

| Age (months) | Estimated exposure mean µg/kg bw/day | Estimated exposure 97.5th percentile µg/kg bw/day | Mean % guidance value | 97.5th percentile % guidance value |
|--------------|--------------------------------------|---|-----------------------|------------------------------------|
| 6 - <12      | 6.19                                 | 22.8  | 36.4                  | 134                                |
| 12 - <18     | 15.2                                 | 35.7  | 89.6                  | 210                                |
| 18 - <24     | 13.8                                 | 37.6  | 81.2                  | 221                                |
| 24 - <48     | 10.9                                 | 28.1  | 64.4                  | 165                                |
| 48 - <60     | 8.09                                 | 21.9  | 47.6                  | 128                                |

Comparing exposure to the JECFA PMTDI as in Table 5 there were no exceedances at mean levels of consumption in any of the selected populations. For the 97.5th percentile of consumption there were exceedances in all populations.

87. Using the consumption rates in Table 1, occurrence data from O’Kane et al. (475.9 µg/kg) and the EVM, (2003) guidance value of 15 µg/kg bw /day, an exposure assessment was undertaken which is presented in Table 6.

Table 6. Exposure assessment from cow's’ milk consumption using mean iodine occurrence in O’Kane et al. (2018), consumption data from the NDNS (Table 1) and the EVM 2003 guidance value (EVM, 2003).

| Age (months) | Estimated exposure mean $\mu\text{g}/\text{kg bw}/\text{day}$ | Estimated exposure 97.5th percentile $\mu\text{g}/\text{kg bw}/\text{day}$ | Mean % guidance value | 97.5th percentile % guidance value |
|--------------|---|--|-----------------------|------------------------------------|
| 6 - <12      | 6.19  | 22.8   | 41.2                  | 152                                |
| 12 - <18     | 15.2  | 35.7   | 102                   | 238                                |
| 18 - <24     | 13.8  | 37.6   | 92.0                  | 251                                |
| 24 - <48     | 10.9  | 28.1   | 73.0                  | 187                                |
| 48 - <60     | 8.09  | 21.9   | 54.0                  | 146                                |

88. Average consumers in the age group 12 - < 18 months slightly exceed the guidance value of 15  $\mu\text{g}/\text{kg bw}/\text{day}$  set by the EVM in 2003. High consumer exposures exceed the guidance value for all age groups.

89. Using the derived SCF/EFSA TULs presented in Table 4, occurrence data from O’Kane et al. (475.9  $\mu\text{g}/\text{kg}$ ) and consumption rates in Table 1 an exposure assessment was undertaken which is presented in Table 7.

Table 7: Exposure assessment from cows’ milk consumption using mean iodine occurrence in O’Kane et al. (2018), consumption data from the NDNS (Table 1) and TUL’s derived from EFSA’s 2006 values and UK average data from NDNS data (EFSA, 2006; Department of Health, 2013; Bates et al., 2014, 2016, 2020; Roberts et al., 2018).

| Age (months) | Estimated exposure mean $\mu\text{g}/\text{kg bw}/\text{day}$ | Estimated exposure 97.5th percentile $\mu\text{g}/\text{kg bw}/\text{day}$ | Mean % guidance value | 97.5th percentile % guidance value |
|--------------|---|--|-----------------------|------------------------------------|
| 6 - <12      | No TUL  | No TUL   | No TUL                | No TUL                             |

|          |      |      |       |     |
|----------|------|------|-------|-----|
| 12 - <18 | 15.2 | 35.7 | 83.00 | 195 |
| 18 - <24 | 13.8 | 37.6 | 84.2  | 229 |
| 24 - <48 | 10.9 | 28.1 | 83.7  | 215 |
| 48 - <60 | 8.09 | 21.9 | 62.8  | 170 |

90. Comparing consumption to the SCF/EFSA TULs, at mean levels of consumption none of the selected populations exceeded derived TULs whilst at the 97.5th percentile of consumption all populations exceeded the TULs.

91. In the COT's 2000 paper, a survey of UK cows' milk from 1998-9 was discussed which identified the overall mean iodine concentration in cow's milk to be 311 µg/kg with a lowered mean concentration in summer (200 µg/kg). These values were used to generate exposure data and their safety assessed against guidance values calculated from the JECFA PMTDI of 0.017 mg/kg bw/day (17 µg/kg bw day) which was available at the time. At mean levels of consumption of the total diet, exceedance of the guidance values was observed for the age group 1½ - 2½ years at 221 µg/day. For the age groups 2½ - 3½, and 3½ - 4 years iodine exposure approached the guidance level at 215 and 204 µg/day respectively. For high level consumers, exceedances for the 3 age groups 1½ - 2½, 2½ - 3½, and 3½ - 4½ years at 362, 379 and 330 µg/day respectively were observed. For milk consumption alone, exceedances of the guidance values calculated from the previously adopted PTWI were present in high level consumers (97.5th percentile) for the groups aged 1½ - 2½, 2½ - 3½ years. The COT concluded that iodine in cows' milk was unlikely to pose a risk to health even in children who are high level consumers (COT, 2000).

92. The COT's 2000 conclusion was reaffirmed in the COT 2017b paper on the risk of excess iodine in the diets of infants and young children arguing:

'These HBGVs are based on limited data. In all cases the relevant studies on which the HBGV was established did not allow an accurate estimation of dietary intakes. The response to high iodine intakes can be highly variable between individuals and will depend on iodine status. Individuals with a low iodine status who are suddenly exposed to high iodine levels are more likely to experience adverse effects than those with an adequate iodine status.'

For many of the parameters of thyroid function normally assessed, it is difficult to distinguish between adverse effects and normal homeostatic changes due to iodine. Further, the RNI and the guidance levels/tolerable daily intakes are of a similar order of magnitude. These two factors, together with the fact that the relevant available studies are all in adult populations, make it difficult to identify a safe upper level which is applicable for all infants and children.'

93. In the COT paper of 2000 on iodine in cow's milk, exceedances were identified for 97.5th percentile consumers. This was mirrored in the exposure assessments produced in this paper with high level consumers of milk exceeding the EVM TDI, TULs derived from EFSA and the JECFA PMTDI. For mean level consumers however, iodine exposure approached the 2003 EVM's 15 µg/kg bw/day TDI for the group 12- <18 months. COT's 2000 and 2017 statements stated that iodine levels in cow's milk were seen to pose no toxicological concern due to the close proximity of the exposures to HBGVs and reasons discussed above. With similar results in this exposure assessment the COT concluded that the risk to health from iodine in cows' milk is likely to be low.

## **Perchlorate**

94. Perchlorate (ClO<sub>4</sub><sup>-</sup>) has both natural and anthropogenic sources. Previous biomonitoring studies have suggested it is most likely to be a ubiquitous compound. It is present in the environment due to Chilean fertilisers and industrial emissions such as ammonium perchlorate in solid rocket fuel propellants and formation of perchlorate from degradation of chlorine-based cleaning products. Within the EU likely sources include Chilean nitrate (fertiliser) leading to accumulation in plants. Plant protection products and water disinfection could slightly increase exposure (EFSA, 2014).

95. Perchlorate acts on the thyroid, inhibiting iodine uptake via the sodium-iodide symporter protein. This leads to depletion in levels of thyroid hormones leading to hypothyroid effects in individuals with a moderate iodine deficiency; this was discussed in a discussion paper in 2018 by the COT (COT, 2018a).

96. An exposure assessment is presented within Annex A using occurrence data for liquid milk from EFSA's 2017 exposure assessment and NDNS consumption data. For mean occurrence at the UB concentration, there were no exceedances of the TDI of 0.3 µg/kg bw/day (from EFSA, 2014) at mean levels of consumption for any age group. For the age range 12-<48 months there were exceedances ranging from 110-140% of the ADI.. Using the 95th percentile UB occurrence

value there was an exceedance of 107% at mean consumption levels for the age group 12-<18 months for all assessed age groups at high level (97.5th percentile) consumers ranging from 153-263%. This, however, is an extremely conservative assessment due to the use of upper bound occurrence values in addition to high consumption levels (97.5th percentile).

97. The COT (2019c) discussed EFSA's assessments (EFSA, 2014, 2017) in 2019. The COT concluded that in both long and short term exposure scenarios for all age groups, there was potential concern, particularly in the case of individuals with mild to moderate iodine deficiency.

98. Based on the exposure assessment presented in Annex A which showed that the TDI was unlikely to be exceeded in a realistic scenario, and, their previous conclusions, the COT concluded that perchlorate levels in cows' milk do not represent a significant health risk to children aged 6 months to 5 years. However, milk is a significant contributor to total perchlorate exposure levels.

## **Chlorate**

99. Chlorate is formed as a by-product of chlorine, chlorine dioxide or hypochlorite usage in disinfecting drinking water, water for plant production and food surface contacts. Chlorine washing of animal derived products is illegal within the EU however plant derived foods can be washed.

100. The EFSA CONTAM panel concluded in their 2015 opinion that the majority of chlorate enters the food chain by washing of food and food contact surfaces. Chlorate is likely to enter milk by cleaned surfaces and sterilised containers (EFSA , 2015a).

101. COT's previous statement on the infant diet (2019c) which included chlorate, discussed EFSA's 2015 opinion and stated that chlorate levels in the total diet were of potential concern for high consumers particularly for individuals with iodine deficiency.

102. In EFSA's 2015 scientific opinion on the risks of chlorate, the mean occurrence of chlorate in liquid milk was calculated at 10 -17 µg/kg (LB-UB) from 38 samples. There was no higher or maximum occurrence value provided. The COT considered that this number of samples was low.

103. An exposure assessment was performed using the mean UB occurrence of chlorate in liquid milk from EFSA (2015a) and is presented in Annex A. No exceedances of the TDI of 3 µg/kg bw day were observed for any age group for

both mean and higher level (95th percentile) consumers.

104. From this information the COT concluded that chlorate in cow's milk is unlikely to pose a risk to health to infants and children aged 6 months – 5 years.

### **Insulin-like Growth Factor (IGF-1)**

105. IGF-1 is a hormone naturally present in both cow's milk and humans. Through treatment with bovine somatotropin (BST), IGF-1 levels in cows can be artificially increased to improve milk production through treatment of cows with bovine somatotropin (BST). BST treatment of cows is illegal within the UK and EU, however milk from BST treated cows can be legally imported. IGF-1 in the diet has been discussed in the scientific literature due to concern over its potential links to cancer.

106. Using data from DEFRA (DEFRA, 2021), liquid drinking milk from BST treated cows is unlikely to enter circulation into the UK in significant amounts. These details are discussed further in Annex A.

107. The COC's 2018 'Statement on possible carcinogenic hazard to consumers from insulin growth factor-1 (IGF-1) in the diet. (COC, 2018) is discussed in annex A. Naturally occurring levels in milk are discussed with the COC stating that the levels in milk consumed by humans are unlikely to exceed 100 ng/ml. The COC concluded there was 'insufficient risk evidence to draw any firm conclusions as to whether exposure to dietary IGF-1 is associated with an increased risk of cancer in consumers. However the data indicate that the levels of IGF-1 consumed are likely to be low and that IGF-1 is likely to be broken down in the gut and not absorbed to any significant extent. Thus, the risk, if any, is likely low.'

108. From this information above and that presented in annex A the COT concluded that it is unlikely that IGF-1 within cows' milk poses a risk to health for children aged 6 months to 5 years of age.

### **Naturally occurring oestrogens in cow's milk**

109. Endogenous oestrogens are oestrogens that are naturally present within cows and humans as well as other animals. They are naturally present within cow's milk. Supplementation of cows with oestrogens is illegal within the UK. There is discussion in the literature over the potential effects of ingested oestrogens on the hypothalamic-pituitary-gonadal axis (HPG axis).

110. Regarding 17 $\beta$ -oestradiol, opinions from the Veterinary Products Committee (Veterinary Products Committee, (2006), JECFA (FAO/WHO, 2000) and

the European Scientific Committee (SCVPH, 2002) have been discussed within annex A. There are varied regulatory opinions on the genotoxicity of 17 $\beta$ -oestradiol however, the COT considers that any genotoxic effect is due to an indirect mechanism.

111. Within annex A is a comparison of exposure to oestrogens naturally found in cow's milk to endogenous production rates of oestrogens in prepubescent boys and girls under the age of 8, presented by JECFA in 2000 (FAO/WHO, 2000). Exposure values were generated from occurrence data from Malekinejad, Scherpenisse and Bergwerff (2006) and NDNS consumption data for children aged 6 months to 5 years of age (Table 1) This was to assess the common claim within the literature that levels of oestrogens within cow's milk would be markedly below the levels produced endogenously within children. The significant uncertainties regarding the endogenous daily production rates in prepubertal children are discussed. From this assessment it has been shown that whilst levels within cow's milk are potentially lower than the total daily production of oestrogens it is unclear how much lower.

112. In addition, an exposure assessment has been performed and is presented within Annex A which compares exposures to the JECFA (2000) ADI of 0.05  $\mu\text{g}/\text{kg}$  bw/day, based on hormonal effects for 17 $\beta$ -oestradiol. No exceedance of the ADI was seen in any population group.

113. From the above information and that discussed further in Annex A, the COT concluded that there is no exceedance of the JECFA 2000 ADI from exposure to oestrogens in cow's milk, and the levels of oestrogens within cow's milk do not present a risk to health for children aged 6 months to 5 years of age.

## **Mycotoxins**

114. Mycotoxins are a highly toxic group of fungi derived compounds. Cow's milk can be contaminated with multiple mycotoxins. A large wealth of information exists regarding occurrence of the aflatoxin M1 in milk. Regarding other mycotoxins, contamination studies have shown variation in the transfer of fumonisins, zearalenone, ochratoxin and trichothecenes from feed to dairy cows and then subsequently into milk. The scientific literature contains far less information on these other mycotoxins and their occurrence in milk.

115. Other mycotoxins are discussed within Annex A with the literature currently suggesting that they are unlikely to transfer into cow's milk from feed and do not present a risk to health for children aged 6 months to 5 years of age.

## Aflatoxins

116. Aflatoxins can enter cow's milk through feed contaminated with fungi such as *Aspergillus flavus* and *Aspergillus parasiticus*. The aflatoxin AFB1 is a common aflatoxin in feed. This is converted within the bovine liver via cytochrome P450 hydroxylation to form the major metabolite AFM1. AFM1 is the most commonly reported and researched mycotoxin within milk, however, AFB1 has also been detected albeit in non-European countries after cattle were exposed to very high levels of AFB1, a scenario which is highly unlikely in UK and EU dairy cattle (Scaglioni et al., 2014; Becker-Algeri et al., 2016). Other aflatoxins include aflatoxins B2, G1, G2 and M2 (AFB2, AFG1, AFG2 and AFM2) and these have also been detected in milk, however, far less information is available on these compounds (EFSA, 2020a).

117. Chronic aflatoxin exposure can lead to immunotoxic effects due to impaired DNA duplication in bone marrow resulting in low leukocyte levels and immunodeficiency, as well as carcinogenic and mutagenic effects. Non-specific cell multiplication inhibition can also affect other cell types with effects particularly apparent within the gastrointestinal tract. The liver is the primary target for aflatoxin exposure. This results in bile duct proliferation, hepatic lesions, centrilobular necrosis and fatty acid infiltration. This often results in liver cancer (Ráduly et al., 2020).

118. Aflatoxins have been reviewed by the SCF in 1996, and EFSA in 1996, 2007 and 2020. They have also been evaluated by JECFA in 1998, 2001 and AFM1 was also reviewed in 2018. EFSA's most recent risk assessment produced by the CONTAM panel concluded that the chronic endpoint of liver carcinogenicity in rats was the most relevant endpoint (EFSA, 2020a). They considered the Wogan et al, study of 1974 to be the most satisfactory for dose response modelling (Wogan, Paglialunga and Newberne, 1974). This value was also used in the COT's 2021 statement on plant-based drinks (see below).

119. The COT's (2021a) overarching statement on consumption of plant-based drinks in children aged 6 months to 5 years of age describes the Wogan, et al. (1974) study as follows: "Groups of male Fisher rats were administered diets containing 0, 1, 5, 15, 50, or 100 µg/kg diet of AFB1 (purity >95%) until clinical deterioration of animals was observed, at which time all survivors in that treatment group were killed. EFSA converted the dietary concentrations of AFB1 into daily intakes assuming that an average adult male rat consumed 40 g diet per kg body using weight per day. EFSA also adjusted the daily intake to 104 weeks in order to compensate for the shorter study duration in some of the AFB1

groups. In the modelling of the results from the Wogan et al. (1974) study the highest dose was omitted because this dose resulted in a 100% tumour incidence. Using model averaging, the BMDL10 for AFB1 was 0.4 µg/kg bw per day.

### **Risk Characterisation**

120. EFSA calculated the contributions of individual food categories in the collected surveys using the LB mean occurrence value in their 2020 risk assessment. It was reported that 'milk and dairy products' were the most substantial contributor to AFM1 exposure for all age groups. For the other children ( $\geq 36$  months to  $< 10$  years old), liquid milk was found to account for up to 89% of exposure to AFM1. Liquid milk also contributed up to 49% of total exposure for infants  $< 12$  months old and up to 74% of total exposure for toddlers ( $\geq 36$  months to  $< 10$  years old). In addition to this, in situations of high exposure liquid milk could contribute up to 89% of total exposure to AFM1. Liquid milk is therefore a significant contributor to AFM1 exposure levels.

121. Analysing the information within EFSA's 2020 risk assessment 'milk and dairy products' contributed  $< 1\%$  of total AFB1 exposure in all surveys. This suggests that there is no risk to health from AFB1 exposure from milk to children aged 6 months to 5 years of age.

122. EFSA also concluded that liquid milk was an important source of exposure of AFM1 + AFT (the sum of AFB1, AFB2, AFG1 and AFG2) for infants, toddlers and children. However, this is driven by high AFM1 contributions.

123. In 2020 EFSA utilised both an animal derived BMDL10 and human epidemiological data to perform 2 risk characterisations, the animal derived BMDL10 approach and the subsequent total dietary exposure assessment are discussed below.

124. In (EFSA, 2020a), for AFM1 a 0.1 potency factor was applied to account for the fact that in a study on Fischer rats, AFM1 was found to induce liver cancer at a rate of 0.1 of that of AFB1. This produced a value of 4.0 µg/kg bw/day for the assessment of AFM1 using a MOE approach (EFSA, 2020a). For mean total dietary AFM1 exposure, MOE values were below 10,000 for infants ( $< 12$  months old) in median and maximum exposure groups, all exposure groups for toddlers ( $\geq 12$  months to  $< 36$  months old) and median UB exposure values and maximum exposure for other children ( $\geq 36$  months to  $< 10$  years old). For the 95th percentile of total dietary exposure all populations within relevant groups ('infants', 'toddlers' and 'other children') exhibited MOE values below 10,000. EFSA commented that this is a health concern however it was noted that high

levels of milk exposure may only occur for a short period in a child's life. For total AFT +AFM1 dietary exposure all age groups and exposure levels exhibited MOEs below 10,000 suggesting there is a health concern from the total diet. MOEs for AFM1 exposure for total dietary exposure are presented below in Tables 9 through to 12. MOEs for AFT + AFM1 for total dietary exposure are presented below in Tables 13 through to 16.

Table 9. MOEs at the lower bound of the minimum, median and maximum mean exposure levels in the total diet to AFM1 from (EFSA, 2020a).

| Age group      | Minimum MOE | Median MOE | Maximum MOE |
|----------------|-------------|------------|-------------|
| Infants        | 28571       | 7018       | 2564        |
| Toddlers       | 8889        | 5882       | 2817        |
| Other Children | 22222       | 11429      | 5128        |

Table 10. MOEs at the upper bound of the minimum, median and maximum at mean exposure levels to AFM1 in the total diet from EFSA (EFSA, 2020a).

| Age group      | Minimum MOE | Median MOE | Maximum MOE |
|----------------|-------------|------------|-------------|
| Infants        | 19048       | 4938       | 2020        |
| Toddlers       | 6250        | 3810       | 2210        |
| Other Children | 14286       | 7692       | 4000        |

Table 11. MOEs at the lower bound of the minimum, median and maximum at 95th percentile exposure levels to AFM1 in the total diet from (EFSA, 2020a).

| Age group | Minimum MOE | Median MOE | Maximum MOE |
|-----------|-------------|------------|-------------|
|-----------|-------------|------------|-------------|

|                |      |      |      |
|----------------|------|------|------|
| Infants        | 6061 | 2703 | 642  |
| Toddlers       | 3810 | 2721 | 1053 |
| Other Children | 9302 | 5000 | 1852 |

Table 12. MOEs at the upper bound of the minimum, median and maximum at 95th percentile exposure levels to AFM1 in the total diet from (EFSA, 2020a).

| Age group      | Minimum MOE | Median MOE | Maximum MOE |
|----------------|-------------|------------|-------------|
| Infants        | 4082        | 1942       | 508         |
| Toddlers       | 2685        | 1835       | 825         |
| Other Children | 6452        | 3175       | 1465        |

Table 13. MOEs at the lower bound of the minimum, median and maximum at mean exposure levels to AFT + AFM1 in the total diet from (EFSA, 2020a).

| Age group      | Minimum MOE | Median MOE | Maximum MOE |
|----------------|-------------|------------|-------------|
| Infants        | 2222        | 952        | 396         |
| Toddlers       | 541         | 325        | 195         |
| Other children | 460         | 328        | 208         |

Table 14. MOEs at the upper bound of the minimum, median and maximum at mean exposure levels to AFT + AFM1 in the total diet from (EFSA, 2020a).

| Age group | Minimum MOE | Median MOE | Maximum MOE |
|-----------|-------------|------------|-------------|
|-----------|-------------|------------|-------------|

|                |     |     |    |
|----------------|-----|-----|----|
| Infants        | 455 | 155 | 40 |
| Toddlers       | 79  | 44  | 32 |
| Other children | 75  | 46  | 32 |

Table 15. MOEs at the lower bound of the minimum, median and maximum at 95th percentile exposure levels to AFT + AFM1 in the total diet from EFSA (EFSA, 2020a).

| Age group      | Minimum MOE | Median MOE | Maximum MOE |
|----------------|-------------|------------|-------------|
| Infants        | 615         | 345        | 122         |
| Toddlers       | 310         | 172        | 90          |
| Other children | 235         | 174        | 91          |

Table 16. MOEs at the upper bound of the minimum, median and maximum at 95th percentile exposure levels to AFT + AFM1 in the total diet from (EFSA, 2020a).

| Age group      | Minimum MOE | Median MOE | Maximum MOE |
|----------------|-------------|------------|-------------|
| Infants        | 99          | 54         | 14          |
| Toddlers       | 48          | 26         | 15          |
| Other children | 53          | 25         | 17          |

125. In light of EFSA's latest risk assessment it is unlikely that AFB1 in liquid milk presents a risk to human health. Cow's milk was, however, found to be a significant contributor (up to 89%) to total dietary exposure of AFM1 and AFM1 +

AFT in 'infants', 'toddlers' and 'other children'. As total dietary exposures to AFM1 and AFM1 + AFT produced MOEs below 10,000 in these populations at a mean exposure level with cow's milk contributing up to 89% to this exposure, a risk to human health cannot be excluded for infants and children aged 6 months to 5 years.

126. In the overarching statement on plant-based drinks it was noted that the margins of exposure for estimated exposure to aflatoxins from almond drink or from the general diet in children 6 months to < 10 years were in general below 10,000, the indicative value for low concern from exposure to a genotoxic carcinogen. However, the exposure estimates were very uncertain, and while exposure would have been overestimated, it was not possible to determine by how much (COT, 2021a).

127. From the above information, the COT concluded that Aflatoxin M1 was identified as being of low concern for children aged 6 months to 5 years of age. There is also a low concern for total aflatoxins within milk however the low MOEs present are largely driven by levels of AFM1 within cow's milk. Other aflatoxins did reduce the MOE further.

## **Per- and polyfluoroalkyl substances (PFAS)**

128. PFAS are a range of synthetic compounds that contain multiple fluorine atoms. They possess excellent surfactant properties and are widely used in consumer products such as paints, polishes and stain repellents. The Organisation for Economic Co-operation and Development (The Organisation for Economic Co-operation and Development OECD, (2021) define PFAS as:

'fluorinated substances that contain at least one fully fluorinated methyl or methylene carbon atom (without any H/Cl/Br/I atom attached to it), i.e. with a few noted exceptions, any chemical with at least a perfluorinated (-CF<sub>3</sub>) or a perfluorinated (-CF<sub>2</sub>-) is a PFAS.'

129. The 2 main classes of PFAS are perfluoroalkyl carboxylic acids (PFCAs) and perfluoroalkane sulfonic acids (PFSAs). In 2020 EFSA undertook a risk assessment related to human health related to the presence of perfluoroalkyl substances in food focussing on 4 of the PFAS. These were two PFCAs: Perfluorooctanoic acid (PFOA), Perfluorononanoic acid (PFNA) and two PFSAs: Perfluorohexane sulfonic acid (PFHxS) and Perfluorooctane sulfonic acid (PFOS) (EFSA, 2020b).

130. Further information on HBGV derivation and a risk characterisation have been discussed within Annex A. Within EFSA's 2020 dietary exposure evaluation no positive samples for the 4 PFAS compounds were detected above the analytical method reporting levels.

131. In an article by Hill, Becanova and Lohmann, (2021) in 13 milk samples taken from US cattle in areas of concern. These were farms which reported biosolid amendments on cropland or were located within proximity to aqueous film forming foam (AFFF) contaminated soils. No perfluoroalkyl acids (PFAAs) were detected with the authors considering that uptake of PFAAs into dairy milk within the U.S. is low.

132. Kowalczyk et al., (2013) in their study of absorption, distribution, metabolism and excretion (ADME) of PFAS contaminated feed in dairy cows concluded that the kinetics of PFOA were similar to PFBS and differed from PFHxS and PFOS. Low levels of PFBS were detected in milk, plasma and trace amounts were detected in the kidneys and liver. Coupled with a high urinary excretion the authors concluded that PFBS does not accumulate in dairy cows.

133. Considering the lack of reported quantifiable amounts of PFHxS, PFOS, PFOA and PFNA in all liquid milk sample data presented by EFSA (2020c) plus the conclusions from Kowalczyk et al. (2013) and Hill, Becanova and Lohmann (2021) , the COT concluded that PFAS exposures via cow's milk are unlikely to be of current health concern to infants and children aged 6 months to 5 years

## **Brominated flame retardants (BFRs)**

134. Brominated flame-retardants (BFRs) are structurally diverse chemicals used in plastics, textiles and other materials to enhance their flame-retardant properties. There are 5 main classes of BFRs:

- i) Hexabromocyclododecanes (HBCDDs), example uses include thermal insulation
- ii) Polybrominated biphenyls (PBBs), example uses include in consumer appliances, textiles and plastic foams
- iii) Polybrominated diphenyl ethers (PBDEs), example uses include in electronic circuitry, casings and textiles
- iv) Tetrabromobisphenol A (TBBPA) and other phenols, example uses include in electronic circuitry and within thermoplastics in TV sets
- v) Other brominated flame retardants.

135. Some BFRs, including polybrominated diphenyl ethers (PBDEs) and hexabromocyclododecane (HBCD) are mixed into polymers rather than being chemically bound to them and can leach out of the products/materials in which they are used and into the environment.

The use of many of the BFRs are restricted or prohibited within the EU, nevertheless due to their persistent nature they are widely distributed in the environment such as within water systems, air and soil. BFRs can therefore readily enter the food chain primarily through animal products such as milk and meat

### **Hexabromocyclododecanes (HBCDDs)**

136. HBCDDs are non-aromatic, brominated cyclic alkanes used primarily as additive flame retardant in materials such as styrene resins. The commercial product consists of three diastereoisomers  $\alpha$ ,  $\beta$  and  $\gamma$ -HBCD. Although technical HBCD typically consists primarily of  $\gamma$ -HBCD, the relative proportions of the isomers varies depending on product application.

137. A discussion of the MOE approach taken by EFSA, COT's 2015 opinion on this work in addition to additional work by EFSA is presented within Annex A (EFSA, 2011a, 2021b; COT, 2015c).

138. Regarding risk characterisation, in EFSA's 2021 assessment the mean LB concentration of HBCDDs within milk was  $< 0.01 \mu\text{g}/\text{kg}$ . The COT concluded that the MOEs by dietary intake of breast milk, infant formula, commercial infant food, fish oil and food in general are at least 400 and not a cause for concern for any age group, as they are considerably greater than 8 (A factor of 2.5 to cover inter-species differences and a factor of 3.2 to cover uncertainties in the elimination half-life in humans were multiplied. MOEs. This produces a value of 8. For MOEs above this level there is adequate reassurance that there is no health concern.)

139. In light of the (EFSA, 2021b) and (COT, 2015c) conclusions (see Annex A) the COT concluded that HBCDDs in cow's milk do not pose a health risk to infants and children aged 6 months to 5 years.

### **Polybrominated biphenyls (PBB)s**

140. (PBBs) are brominated hydrocarbons formerly used as additive flame retardants. As such these substances were added, rather than chemically bound to plastics used in a variety of consumer products, such as computer monitors,

television, textiles and plastic foams, and were able to leave the plastic and enter the environment. They are structurally similar compounds in which 2-10 bromine atoms are attached to the biphenyl molecular structure. In total, as with the structurally similar Polychlorinated Biphenyls (PCBs), 209 different PBB congeners are possible.

141. EFSA concluded that 'the risk to the European population from exposure to PBBs through the diet is of no concern.' Levels in milk were obtained for BB-52 and BB-101 at the levels of 0.55 to 6.83 ng/kg fat (LB and UB) and 0.64 to 6.92 pg/g fat (LB and UB) for BB-52 and BB-101 respectively (EFSA, 2010). This is discussed further in Annex A.

142. In 2015 the COT concluded that a reliable estimation of infants' exposures to PBBs was not possible due to limitations within data sources such as the number of congeners covered and a lack of UK data. In spite of this they considered it a low priority due to the restriction of their use (COT, 2015a). Within the literature (discussed in Annex A), minimal levels of PBBs have been reported in milk.

143. In light of the EFSA, (2010) conclusion, the COT 2015 statement and evidence from the literature the COT concluded that PBBs in cow's milk do not pose a health risk to infants and children aged 6 months to 5 years.

## **PBDEs**

144. PBDEs are produced by direct bromination of diphenyl ether. There are 209 individual PBDE congeners, each of which is identifiable by a unique congener number. Three commercial PBDE flame-retardants, pentabromodiphenyl ether (pentaBDE), octabromodiphenyl ether (octaBDE) and decabromodiphenyl ether (decaBDE) have been available in the UK. The commercial PBDEs are not pure products but a mixture of various diphenyl ethers with varying degrees of bromination.

145. EFSA's 2011 exposure assessment (discussed further in Annex A) determined that the only safety concern was for young children aged 1- < 3. Milk contributed a low percentage to this total dietary exposure

146. A review of the literature for occurrence of PBDEs in milk did not show a concern for health due to low concentrations (discussed in Annex A)

147. The COT concluded in 2015 that there was a possible concern with respect to exposure of infants to BDE-99 and (to a lesser extent) BDE-153 from food,

other than commercial infant food (COT, 2015b). An addendum to this statement was produced in 2017 that further concluded that 'The current analysis indicated that exposure of young children aged 1-5 years to these congeners from such food was unlikely to be a health concern' (COT, 2017a).

148. Reviewing the EFSA (2011b) and COT (2015b; 2017a) conclusions in addition to the evidence from the literature that cow's milk does not contain levels of concern, the COT concluded that PBDEs in cow's milk do not pose a risk to health for infants and children aged 6 months to 5 years.

## **Tetrabromobisphenol A (TBBPA)**

149. Worldwide, TBBPA is the most widely used BFR and approximately 90% of TBBPA, manufactured by bromination of bisphenol A, is used as a reactive intermediate in the manufacture of epoxy and polycarbonate resins. In this case it is covalently bound to the polymer and is unlikely to escape into the environment. The remaining 10% is used as an additive flame retardant, where it does not react chemically with the other components of the polymer and may therefore leach out of the matrix into the environment.

150. (EFSA, 2011b) and the COT (2019d) concluded that there was no risk to health from TBBPA. EFSA found that dietary exposures produced large MOEs with cow's milk not containing any TBBPA above reporting levels. The work by COT in 2019 produced MOEs larger than those generated in EFSA's 2011 evaluation. The COT concluded that with respect to cow's milk exposure the MOEs were adequately protective. This is further discussed in Annex A.

151. In light of the EFSA (2011b) and COT (2019c) conclusions and evidence from the literature (further discussed in Annex A) that cow's milk does not contain levels of concern, the COT concluded that TBBPA in cow's milk does not pose a risk to health for infants and children aged 6 months to 5 years

## **Microplastics**

152. Plastic pollution has been widely recognised as a global environmental problem (Villarrubia-Gómez, Cornell and Fabres, 2018). The adverse effects of plastic litter have been widely documented for marine animals (e.g. entanglement, ingestion and lacerations); however, the potential risks from exposure to smaller plastic particles i.e. micro- and nanoplastics in humans are yet to be fully understood.

153. Due to their widespread presence in the environment, microplastics also occur in food (e.g. seafoods, beer, salt and honey, tea, vegetables) and drinks (e.g. bottled water, milk, soft drinks) (Toussaint et al., 2019). The occurrence of microplastics in milk will likely be due to contamination from dairy machinery and / or packaging rather than the cow itself.

154. The ECHA in 2019 listed their four major concerns posed by the presence of microplastics in the environment, listed in Annex A (ECHA, 2019).

155. (COT, 2021b) stated that a full risk assessment on the potential toxic effect(s) of microplastics could not be carried out. This was due to the lack of toxicokinetic and toxicity data in general, the paucity of currently available data for microplastics in different food types and the difficulty of performing an accurate exposure assessment. Whilst risks from microplastics cannot be quantified currently, microplastic contamination in milk is likely to be lower than other foodstuffs.

156. The COT concluded from the above information and that included in Annex A that microplastics in milk currently do not represent a risk to health for children aged 6 months to 5 years of age.

## **Summary**

157. To aid in the risk assessment of the chemicals described in this statement three summary tables are provided (Tables 17, 18 and 19) providing a summary of the conclusions and where appropriate to this paper, the HBGV for each substance and highest age range estimated exposure via the diet, based on the mean consumption data.

## **Conclusions**

158. The COT reviewed an extensive of chemical compounds that could be present as contaminants in cow's milk to allow comparison with plant based dairy alternatives.

159. As can be seen in the summary tables, the vast majority of these potential contaminants present no risk of adverse health effects at the levels currently observed within cow's milk.

160. Based on the levels found in cow's milk, iodine, AFM1 specifically and total aflatoxins due to AFM1 levels, represent a low risk to health. Risk to health

relating to isoflavones is uncertain due to no HBGVs being set for young children and therefore a lack of knowledge of the significance of the levels in milk. Any risk from isoflavones in cow's milk will be lower than that in soya milk due to lower occurrence levels.

Table 17. Summary of risk assessment conclusions for selected compounds and their occurrence levels within cows' milk based on previous authority opinions.

| <b>Compound (s)</b>                               | <b>HBGV, (endpoint)</b>                | <b>Effect (s)</b>   | <b>Authority</b> | <b>Conclusion: Health risk from cow's milk</b>  |
|---|--|---|------------------|---|
| Nitrite   | n/a                                    | Methemoglobinemia   | EFSA             | No health concern   |
| Bisphenol A                                       | 4 µg (Increase in mouse kidney weight) | Endocrine disrupter affecting metabolism, growth, sexual development, stress response, insulin production, gender behaviour, reproduction, and foetal development | EFSA             | No health concern (However currently awaiting EFSA's updated position on BPA expected 2022) |
| DBP, BBP, DEHP, DINP (Summed as DEHP equivalents) | 0.05 mg (reproductive effects in rats) | Reproductive effects, hepatic effects   | EFSA / COT       | No health concern   |

|                                      |  |  |                |  |
|--------------------------------------|--|--|----------------|--|
| DEP                                  | 5 mg (increased liver and prostate weights, decreased epididymal sperm concentration of the F1 generation in mice) | Organ weight changes   | WHO / COT      | No health concern  |
| NDL-PCBs                             | n/a  | Neurological, endocrine, immunological and carcinogenic effects  | JECFA          | No health concern  |
| Isoflavones<br>GEN, EQU,<br>FOR, DAI | 0.07 mg (GEN only)   | Endocrine disrupter (oestrogenic effects) effecting thyroid and immune function and sexual development | Nordic Council | Any risk to health is uncertain as for the majority of isoflavones an HBGV has not been set for young children. Any concern is lower than that in soya milk. |
| Lead                                 | None, BMDL01 of 0.5 µg/kg bw/day (development of intellectual function)  | Multiple toxic effects   | EFSA/COT       | No health concern  |
| Inorganic Arsenic                    | None. BMDL0.5 of 3 µg/kg bw/day JECFA / COT (lung cancer)  | Multiple toxic effects including carcinogenicity   | EFSA/COT       | No health concern  |

|                   |   |   |            |  |
|-------------------|---|---|------------|--|
| Inorganic Mercury | TWI - 4 µg/kg bw/week (kidney weight change in rats)  | Multiple toxic effects including renal, haematological, hepatic and gastrointestinal effects.   | EFSA / COT | No health concern  |
| Cadmium           | TWI - 2.5 µg/kg bw/week (urinary β-2-microglobulin (B2M) as a marker for kidney damage)   | Multiple toxic effects including renal toxicity, hepatotoxicity, osteoporosis and osteomalacia. | EFSA / COT | No health concern  |
| AFM1              | None. Guidance value of 4 µg/kg bw/day derived from a BMDL10 based on tumour incidence for AFB1 in rats with a 0.1 potency factor applied | Multiple effects such as immunotoxicity, carcinogenicity and mutagenicity                       | EFSA / COT | Low- - concern   |
| AFB1              | None. BMDL10 of 0.4 µg/kg bw/day based on tumour incidence in rats after AFB1 exposure.   | Multiple effects such as immunotoxicity, carcinogenicity and mutagenicity                       | EFSA / COT | No health concern  |
| Total aflatoxins  | None. BMDL10 of 0.4 µg/kg bw/day based on tumour incidence in rats after AFB1 exposure.   | Multiple effects such as immunotoxicity, carcinogenicity and mutagenicity                       | EFSA / COT | Low concern, contributions driven by AFM1 milk occurrence. |

|  |  |   |      |                      |
|--|--|---|------|----------------------|
| PFAS<br>(PFHxS,<br>PFOS, PFOA<br>and PFNA) | TWI of 4.4 µg/kg<br>bw/day (reduced<br>antibody levels<br>against diphtheria<br>vaccine in 1-year old<br>children)     | increased relative<br>liver weight, effects<br>on the immune<br>system  | EFSA | No health<br>concern |
| HBCDDs                                     | None. From a LOAEL<br>(neurodevelopmental<br>effects in mice)<br>maximum chronic<br>intake of 2.35 µg/kg<br>bw per day | Neurodevelopmental,<br>immune system<br>effects, reproductive<br>system effects, liver<br>effects and thyroid<br>hormone<br>homeostasis | EFSA | No health<br>concern |
| PBBs                                       | None. NOEL of 0.15<br>mg/kg bw (hepatic<br>carcinogenicity)  | Multiple effects<br>(dioxin like) such as<br>altered vitamin A<br>homeostasis,<br>chloracne and body<br>weight changes                  | EFSA | No health<br>concern |
| PBDEs                                      | None. Range of<br>BMDL10 s between<br>12 and 1,700 µg/kg<br>bw<br>(neurodevelopmental<br>effects)                      | Neurodevelopmental,<br>immune system<br>effects, reproductive<br>system effects, liver<br>effects and thyroid<br>hormone<br>homeostasis | EFSA | No health<br>concern |
| TBBPA                                      | None. BMDL10 of 16<br>mg/kg bw (thyroid<br>hormone<br>homeostasis)   | Thyroid hormone<br>regulation   | EFSA | No health<br>concern |

Table 18. Summary table displaying a comparison of highest estimated mean exposures (occurrence and consumption) to potential chemical contaminants of cow's milk with their health-based guidance values, from exposure assessments presented in this paper and its annex.

| <b>Compound (s)</b>                 | <b>HBGV, (endpoint)</b>                               | <b>Authority</b> | <b>Highest Exposure (mean consumption), kg bw/day</b> | <b>% HBGV or MOE</b> | <b>Highest exposure age range (months)</b> | <b>Effect</b>   |
|-------------------------------------|---|------------------|---|----------------------|--|---|
| Nitrate                             | 3.7 mg (growth retardation in dogs and rats)          | EFSA             | 0.00416 mg  | 0.112                | 12 - <18                                   | Methemoglobinemia   |
| Dioxins plus DL-PCBs                | 2 pg WHO-TEQ, (reproductive effects in rats)          | EFSA             | 1.024 pg  | 51.2                 | 12 - <18                                   | Range of effects including chloracne and reproductive effects |
| Benzo[a]pyrene (BaP)                | None, BMDL10 of 70 µg (total tumour-bearing animals)  | EFSA             | 0.00128 µg  | 54,688 (MOE)         | 12 - <18                                   | Carcinogenic  |
| Sum of BaP, BbF, ChR and BaA (PAH4) | None, BMDL10 of 340 µg (total tumour-bearing animals) | EFSA             | 0.0032 µg   | 106,250 (MOE)        | 12 - <18                                   | Carcinogenic  |

|             |   |                          |          |       |                      |          |  |
|-------------|---|--------------------------|----------|-------|----------------------|----------|--|
| Iodine      | EVM: Guidance level of 15 µg/kg bw/day<br><br>(Alterations in serum thyroid hormone levels from human studies)      | COT / EVM / EFSA / JECFA | 15.2 µg  | 102   | (EVM guidance value) | 12 - <18 | Varied effect depending on previous iodine intake                |
|             | EFSA: TUL of 200-250 µg/day   |                          |          |       |                      |          |  |
|             | JECFA: PMTDI 17 µg/kg bw/day  |                          |          |       |                      |          |  |
| Perchlorate | TDI of 0.3 µg/kg bw/day<br><br>(inhibition of radiolabelled iodine uptake by the thyroid)                           | EFSA                     | 0.179 µg | 59.6  |                      | 12 - <18 | Inhibition of iodine uptake, decreased thyroid hormone synthesis |
|             | TDI of 3 µg/kg bw/day   |                          |          |       |                      |          |  |
| Chlorate    | (Carried over from perchlorate with a 0.1 potency factor, inhibition of radiolabelled iodine uptake by the thyroid) | EFSA                     | 0.544 µg | 18.1% |                      | 12 - <18 | Inhibition of iodine uptake, decreased thyroid hormone synthesis |

|  |   |           |       |          |  |
|--|---|-----------|-------|----------|--|
| Naturally occurring oestrogens within cows' milk | <p>ADI - 0.05 µg/kg bw/day for 17β-oestradiol (NOEL based off of multiple hormone dependent parameters in postmenopausal JECFA women. To protect sensitive population subgroups an uncertainty factor of 10 was applied.)</p> | 0.0875 µg | 17.5% | 12 - <18 | <p>Suggeste<br/>in childre<br/>developm<br/>effects in<br/>urogenita<br/>hormona<br/>central n<br/>systems<br/>mamma</p> <p>The COT<br/>any geno<br/>17β-oest<br/>indirect i</p> |
|--|---|-----------|-------|----------|--|

Table 19. A summary of information for compounds within milk where a satisfactory standard risk assessment of the compounds within cows' milk could not be performed.

| <b>Compound (s)</b>  | <b>Literature evaluation</b>  | <b>Effect</b>   | <b>Conclusion: Health risk from cow's milk</b> |
|----------------------|---|-----------------|--|
| Veterinary Medicines | Incidents where veterinary medicines are found within UK milk are uncommon. isolated incidents. | Various effects | No health concern                              |

|                  |  |  |                   |
|------------------|--|--|-------------------|
| Pesticides       | Between 2015 and the end of 2020 only 1 cows' milk sample of 1,723 returned a positive result (above the maximum residue level). The risk of pesticides form cow's milk is minimal   | Various effects  | No health concern |
| IGF-1            | IGF-1 supplementation is unlikely to generate a risk to consumer health. In addition milk from IGF-1 treated cow's is unlikely to enter the UK as fresh milk in significant quantities.  | No substantiated carcinogenic effects                              | No health concern |
| Other mycotoxins | Milk is considered unlikely to contain significant amounts of other mycotoxins   | Effects including immunotoxicity, carcinogenicity and mutagenicity | No health concern |
| Microplastics    | A lack of toxicokinetic and toxicity data in general, the paucity of currently available data for microplastics in different food types and difficulties in performing an accurate exposure assessment, however levels of microplastics in milk are lower than in other areas of the diet. | Various, depending on type   | No health concern |

## **Annex B to TOX/2022/23**

### **Abbreviations and Technical Information**

|           |  |
|-----------|--|
| ADI       | Acceptable Daily Intake                            |
| 15-Ac-DON | 15-Acetyldeoxynivalenol                            |
| 3-Ac-DON  | 3-Acetyldeoxynivalenol                             |
| ADME      | Absorption, Distribution, Metabolism and Excretion |
| AFB1      | <i>Aflatoxin B1</i>                                |
| AFB1      | Aflatoxin B1                                       |
| AFB2      | Aflatoxin B2                                       |
| AFFF      | Aqueous Film Forming Foam                          |
| AFG1      | Aflatoxin G1                                       |
| AFM1      | <i>Aflatoxin M1</i>                                |
| AFM1      | Aflatoxin M1                                       |
| AFM2      | Aflatoxin M2                                       |
| AFT       | Sum of AFB1, AFB2, AFG1 and AFG2                   |
| AhR       | Aryl Hydrocarbon Receptor                          |
| As        | Arsenic  |
| BaA       | Benz[a]anthracene                                  |

|      |   |
|------|---|
| BaP  | Benzo[a]pyrene  |
| BbF  | Benzo[b]fluoranthene  |
| BBP  | Butyl-benzyl-phthalate  |
| BFR  | Brominated Flame Retardants                                       |
| BIO  | Biochanin A   |
| BMDL | Benchmark Dose Lower Confidence Limit                             |
| BPA  | Bisphenol A   |
| Br   | Bromine   |
| BST  | Bovine Somatotropin   |
| bw   | Body Weight   |
| CAR  | Constitutive androstane receptor                                  |
| Cd   | Cadmium   |
| CEP  | EFSA Panel on Food Contact Materials, Enzymes and Processing Aids |
| CF2  | Perfluorinated <a href="#">Methylene Group</a>                    |
| CF3  | Perfluorinated <a href="#">Methyl Group</a>                       |
| ChR  | Chrysene  |

|         |   |
|---------|---|
| Cl      | Chlorine  |
| COC     | The Committee on Carcinogenicity Food, Consumer Products and the Environment      |
| CONTAM  | EFSA Panel on Contaminants in the Food Chain                                      |
| COT     | Committee on Toxicity of Chemicals in Food, Consumer Products and the Environment |
| DAI     | Daidzein  |
| DBP     | Di-butylphthalate   |
| DecaBDE | Decabromodiphenyl ether   |
| DEFRA   | Department for Environment, Food and Rural Affairs                                |
| DEHP    | Bis(2- ethylhexyl)phthalate   |
| DHSC    | Department of Health and Social Care  |
| DIDP    | Di-isodecylphthalate  |
| DINP    | Di-isononylphthalate  |
| DL-PCBs | Dioxin-Like Polychlorinated Biphenyls   |
| DL-PCBs | Dioxins and Dioxin-Like Polychlorinated   |
| DNSIYC  | Diet and Nutrition Survey of Infants and Young Children                           |

|                 |                                       |
|-----------------|---------------------------------------|
| DON             | Deoxynivalenol                        |
| DON-3-glucoside | Deoxynivalenol-3-Glucoside            |
| E1              | Oestrone                              |
| E2              | 17 $\beta$ -Oestradiol                |
| EC              | European Commission                   |
| ECHA            | European Chemical Agency              |
| EFSA            | European Food Safety Authority        |
| EFSA            | European Food Safety Authority        |
| EHDI            | Estimated Human Daily Intakes         |
| EQU             | Equol                                 |
| EQU             | Equol (metabolite of DAI)             |
| ERs             | Oestrogen Receptors                   |
| EU              | European Union                        |
| EVM             | Expert Group on Vitamins and Minerals |
| FAO             | Food and Agriculture Organisation     |

|                  |                              |
|------------------|------------------------------|
| FDA              | Food and Drug Administration |
| FOR              | Formononetin                 |
| FSA              | Food Standards Agency        |
| FSH              | Follicle Stimulating Hormone |
| FTOHs            | Fluorotelomer alcohols       |
| GEN              | Genistein                    |
| GH               | Growth Hormone               |
| GI               | Gastrointestinal             |
| H                | Hydrogen                     |
| HBCD             | Hexabromocyclodecane         |
| HBGV             | Health Based Guidance Value  |
| HED              | Human Equivalent Dose        |
| Hg               | Mercury                      |
| Hg <sup>+</sup>  | Mercurous cation             |
| Hg <sup>0</sup>  | Elemental mercury            |
| Hg <sup>2+</sup> | Mercuric cation              |

|          |   |
|----------|---|
| HPG axis | Hypothalamic-Pituitary-Gonadal Axis               |
| I        | Iodine  |
| IARC     | International Agency for Research on Cancer       |
| iAS      | Inorganic Arsenic                                 |
| ICES- 6  | Indicator PCBs: 28, 52, 101, 138, 153 and 180     |
| IGF-1    | Insulin-like Growth Factor 1                      |
| IGFBP-3  | Insulin Growth Promoting Factor Binding Protein 3 |
| IQ       | Intelligence quotient                             |
| JECFA    | Joint FAO/WHO Expert Committee on Food Additives  |

Lower Bound- - Lower bound and upper bound approaches are utilised in order to assess left censored data (Occurrence values below the limits of detection or quantification).

|     |   |
|-----|---|
| LB  | The lower bound refers to situations where a zero value has been assigned to occurrence values below the limit of detection or limit of quantification. |
| LH  | Luteinising Hormone   |
| LOD | Limit of Detection  |

|          |   |
|----------|---|
| MB       | Middle Bound - The middle bound is an approach for assessing left censored data. Any values below the limit of detection (LOD) or limit of quantification (LOQ) are assigned the value LOD/2 or LOQ/2 respectively. |
| mg       | Milligram   |
| mm       | Millimetre  |
| MoBB     | Margin of Body Burdens  |
| MOE      | Margin Of Exposure  |
| MRL      | Maximum Residue Limit   |
| MT       | Metallothionein   |
| NDL-PCBs | Non-Dioxin-Like Polychlorinated Biphenyls   |
| ng       | Nanogram  |
| NHS      | National Health Service   |
| NIS      | Na <sup>+</sup> /I <sup>-</sup> symporter   |
| nm       | Nanometre   |
| NOAELs   | No-Observed-Adverse-Effect Levels   |
| NOEL     | No Observed Effect Level  |

|           |  |
|-----------|--|
| NRL       | National Reference Laboratory                              |
| NSAIDS    | Non-Steroidal Anti-inflammatory drugs                      |
| OctaBDE   | Octabromodiphenyl Ether                                    |
| OECD      | The Organisation for Economic Co-operation and Development |
| OTA       | Ochratoxin A   |
| PAHs      | Polycyclic Aromatic Hydrocarbons                           |
| PAPs      | Polyfluorinated Phosphate Esters                           |
| Pb        | Lead   |
| PBB-169   | 3,3',4,4',5,5'-hexaBB                                      |
| PBBs      | Polybrominated Biphenyls                                   |
| PBDEs     | Polybrominated Diphenyl Ethers                             |
| PCBs      | Polychlorinated Biphenyls                                  |
| PCDDs     | Polychlorinated Dibenzodioxins                             |
| PCDFs     | Polychlorinated Dibenzofurans                              |
| PE        | Polyethene   |
| PentaPBDE | Pentabromodiphenyl Ether                                   |

|       |  |
|-------|--|
| PFAAs | Perfluoroalkyl Acids                       |
| PFAS  | Per- and polyfluoroalkyl substances        |
| PFBS  | Perfluorobutanesulfonic Acid               |
| PFCAs | Perfluoroalkyl Carboxylic Acids            |
| PFHxS | Perfluorohexane sulfonic acid              |
| PFNA  | Perfluorononanoic Acid                     |
| PFOA  | Perfluorooctanoic Acid                     |
| PFOS  | Perfluorooctane sulfonic acid              |
| PFSA  | Perfluoroalkane Sulfonic Acids             |
| pg    | picograms                                  |
| PHE   | Public Health England                      |
| PMTDI | Provisional Maximum Tolerable Daily Intake |
| PP    | Polypropene                                |
| PTMI  | Provisional tolerable Monthly Intake       |
| PTWI  | Provisional Tolerable Weekly Intake        |
| RASFF | Rapid Alert System for Food and Feed       |

|       |   |
|-------|---|
| SACN  | Scientific Advisory Committee on Nutrition                            |
| SCF   | Scientific Committee on Food  |
| SCF   | European Scientific Committee on Food                                 |
| SCVPH | Scientific Committee on Veterinary measures relating to Public Health |
| SD    | Standard Deviation  |
| SUL   | Safe Upper Level  |
| TBBPA | Tribromobisphenol A   |
| TCDD  | 2,3,7,8-Tetrachlorodibenzyl Dioxin                                    |
| TDI   | Tolerable Daily Intake  |
| TDS   | UK Total Diet Study   |
| TEF   | Toxicity Equivalency Factor   |
| TEQ   | Toxic Equivalent Value  |
| TSH   | Thyroid-Stimulating Hormone   |
| TUL   | Tolerable Upper Level   |
| TWI   | Tolerable Weekly Intake   |

|        |  |
|--------|--|
| UB     | Upper Bound - Lower bound and upper bound approaches are utilised in order to assess left censored data (Occurrence values below the limits of detection or quantification). In the upper bound approach any occurrence levels below the limit of detection or limit of quantification (left censored data) are assigned the value of the limit of detection or the limit of quantification. |
| U-Cd   | Urinary Cadmium  |
| UK     | United Kingdom   |
| US     | United States  |
| US-EPA | United States Environmental Protection Agency  |
| VMD    | Veterinary Medicines Directorate   |
| VPC    | Veterinary Products Committee  |
| WHO    | World Health Organisation  |
| β2M    | β-2-microglobulin  |
| μg     | microgram  |

## References

Bansal, A., Henao-Mejia, J. and Simmons, R.A. (2018) 'Immune System: An Emerging Player in Mediating Effects of Endocrine Disruptors on Metabolic Health', *Endocrinology*, 159(1), pp. 32–45. doi:10.1210/en.2017-00882. [Immune System: An Emerging Player in Mediating Effects of Endocrine Disruptors on Metabolic Health. - Abstract - Europe PMC](#)

Bates, B. et al. (2014) National Diet and Nutrition Survey Results from Years 1, 2, 3 and 4 (combined) of the Rolling Programme (2008/2009 – 2011/2012). [Main heading \(publishing.service.gov.uk\)](#)

Bates, B. et al. (2016) 'National Diet and Nutrition Survey Results from Years 5 and 6 (combined) of the Rolling Programme (2012/2013 – 2013/2014)'. [Main heading \(publishing.service.gov.uk\)](#)

Bates, B. et al. (2020) 'National Diet and Nutrition Survey Results from Years 1, 2, 3 and 4 (combined) of the Rolling Programme (2016/2017 – 2018/2019)'. [National Diet and Nutrition Survey \(publishing.service.gov.uk\)](#)

Bath, S.C., Button, S. and Rayman, M.P. (2012) 'Iodine concentration of organic and conventional milk: implications for iodine intake', British Journal of Nutrition. 2011/07/05 edn, 107(7), pp. 935–940. doi:10.1017/S0007114511003059. [Iodine concentration of organic and conventional milk: implications for iodine intake - PubMed \(nih.gov\)](#)

Becker-Algeri, T.A. et al. (2016) 'Mycotoxins in Bovine Milk and Dairy Products: A Review', Journal of Food Science, 81(3), pp. R544–R552. doi:10.1111/1750-3841.13204. [Mycotoxins in Bovine Milk and Dairy Products: A Review - Becker-Algeri - 2016 - Journal of Food Science - Wiley Online Library](#)

Bláhová, L. et al. (2016) 'Phytoestrogens in milk: Overestimations caused by contamination of the hydrolytic enzyme used during sample extraction', Journal of Dairy Science, 99(9), pp. 6973–6982. doi:10.3168/jds.2016-10926. [Phytoestrogens in milk: Overestimations caused by contamination of the hydrolytic enzyme used during sample extraction - PubMed \(nih.gov\)](#)

Bürgi, H. (2010) 'Iodine excess', Best Practice & Research Clinical Endocrinology & Metabolism, 24(1), pp. 107–115. doi:10.1016/j.beem.2009.08.010. [Iodine excess - PubMed \(nih.gov\)](#)

Byrne, M.P. et al. (2020) 'Urease and Nitrification Inhibitors—As Mitigation Tools for Greenhouse Gas Emissions in Sustainable Dairy Systems: A Review', Sustainability, 12(15), p. 6018. doi:10.3390/su12156018. [Sustainability | Free Full-Text | Urease and Nitrification Inhibitors—As Mitigation Tools for Greenhouse Gas Emissions in Sustainable Dairy Systems: A Review \(mdpi.com\)](#)

Cirillo, T. et al. (2015) 'Exposure to Di-2-Ethylhexyl Phthalate, Di-N-Butyl Phthalate and Bisphenol A through Infant Formulas', Journal of Agricultural and Food Chemistry, 63(12), pp. 3303–3310. doi:10.1021/jf505563k. [Exposure to Di-2-](#)

[Ethylhexyl Phthalate, Di-N-Butyl Phthalate and Bisphenol A through Infant Formulas | Journal of Agricultural and Food Chemistry \(acs.org\)](#)

COC (2018) Statement on possible carcinogenic hazard to consumers from insulin-like growth factor-1 (IGF-1) in the diet. [COC statement on IGF-1 and cancer risk \(publishing.service.gov.uk\)](#)

Cockburn, A. et al. (2013) 'Nitrite in feed: From Animal health to human health', Toxicology and Applied Pharmacology, 270(3), pp. 209–217. doi:10.1016/j.taap.2010.11.008. [Nitrite in feed: from animal health to human health - PubMed \(nih.gov\)](#)

COT (2000) Statement on iodine in cows' milk. [iodin2 \(food.gov.uk\)](#)

COT (2001) Statement on the tolerable daily intake for dioxins and dioxin-like polychlorinated biphenyls. [\[ARCHIVED CONTENT\] UK Government Web Archive - The National Archives](#)

COT (2003) 'Phytoestrogens and Health'. [phytoreport0503.pdf \(food.gov.uk\)](#)

COT (2011) 'Statement on dietary exposure to phthalates - data from the total diet study'. Available at: [Microsoft Word - phthalates statement 04-11.docx \(food.gov.uk\)](#)

COT (2013a) 'Statement on the potential risks from high levels of soyaphytoestrogens in the infant diet'. [\[ARCHIVED CONTENT\] COT Statement on Phytoestrogens | Food Standards Agency \(nationalarchives.gov.uk\)](#)

COT (2013b) Statement on the potential risks from lead in the infant diet. [\[ARCHIVED CONTENT\] UK Government Web Archive - The National Archives](#)

COT (2015a) 'Statement on polybrominated biphenyls (PBBs) in the infant diet'. [TOX -2015-06 PBB 2nd Draft Statement.pdf \(food.gov.uk\)](#)

COT (2015b) 'Statement on potential risks from polybrominated diphenyl ethers (PBDEs) in the infant diet: potential risks from polybrominated diphenyl ethers (PBDEs) in the diets of infants and young children'. [PBDEstatementfinal.pdf \(food.gov.uk\)](#)

COT (2015c) 'Statement on the potential risks from hexabromocyclododecanes (HBCDDs) in the infant diet'. [\[ARCHIVED CONTENT\] COT Statement on hexabromocyclododecanes \(HBCDDs\) | Food Standards Agency \(nationalarchives.gov.uk\)](#)

COT (2016a) Addendum to the 2013 COT statement on potential risks from lead in the infant diet. [\[ARCHIVED CONTENT\] Addendum to the 2013 COT statement on lead | Food Standards Agency \(nationalarchives.gov.uk\)](#)

COT (2016b) Statement on potential risks from arsenic in the diet of infants aged 0 to 12 months and children aged 1 to 5 years. [\[ARCHIVED CONTENT\] COT Statement on Arsenic | Food Standards Agency \(nationalarchives.gov.uk\)](#)

COT (2017a) Addendum to the 2015 COT statement on potential risks from polybrominated diphenyl ethers (PBDEs) in the infant diet: potential risks from polybrominated diphenyl ethers (PBDEs) in the diets of infants and young children. [Tox 2017-3 PBDEs final \(food.gov.uk\)](#)

COT (2017b) Statement on the potential risks from excess iodine in the diets of infants aged 0-12 months and children aged 1 to 5 years. [\[ARCHIVED CONTENT\] Statement on the potential risks from excess iodine | Food Standards Agency \(nationalarchives.gov.uk\)](#)

COT (2018a) Review of potential risks from contaminants in the diet of infants aged 0 to 12 months and children aged 1 to 5 years. [tox2018-31.pdf \(food.gov.uk\)](#)

COT (2018b) Statement on potential risks from cadmium in the diet of infants aged 0 to 12 months and children aged 1 to 5 years. [The potential risks from cadmium in the infant diet \(food.gov.uk\)](#)

COT (2018c) Statement on potential risks from methylmercury in the diet of infants aged 0 to 12 months and children aged 1 to 5 years. [\[ARCHIVED CONTENT\] COT Statemet on Methylmercury | Food Standards Agency \(nationalarchives.gov.uk\)](#)

COT (2019a) 'Discussion paper on soya drink consumption in children aged 6 months to 5 years of age'. [TOX 2019-65 Discussion paper on soya drink \(food.gov.uk\)](#)

COT (2019b) Minutes of the meeting held on Tuesday, 19th of March 2019 in Broadway House Conference Centre, Tothill St, London SW1H 9NQ. [COT Final Minutes March 2019 \(food.gov.uk\)](#)

COT (2019c) 'Overarching statement on the potential risks from contaminants in the diet of infants aged 0 to 12 months and children aged 1 to 5 year'. [TOX-2018-48 Overarching statement 0-5 year olds \(food.gov.uk\)](#)

COT (2019d) Review of potential risks from tetrabromobisphenol A (TBBPA) in the diet of infants aged 0 to 12 months and children aged 1 to 5 years. [Potential risks from TBBPA in infant diet 0-5 years \(food.gov.uk\)](#)

COT (2020) Conclusions of the Overarching Statement and Addendum on the potential risks from contaminants in the diet of infants aged 12 to 60 months – Summary tables for SACN. [TOX 2020 36 SACN Report 1 to 5\\_COT chapter \(food.gov.uk\)](#)

COT (2021a) Overarching statement on consumption of plant-based drinks in children aged 6 months to 5 years of age. [Final Plant based drinks statement 2021 \(food.gov.uk\)](#)

COT (2021b) 'Overarching statement on the potential risks from exposure to microplastics'. [COT Microplastics Overarching Statement 2021 \(food.gov.uk\)](#)

COT (2021c) 'Position paper on dioxins'. [Dioxin Interim Position Statement \(food.gov.uk\)](#)

DEFRA (2021) Agriculture in the United Kingdom 2020, p. 90. [Agriculture in the United Kingdom 2020 - GOV.UK \(www.gov.uk\)](#)

Department of Health (2013) Diet and Nutrition Survey of Infants and Young Children (DNSIYC), 2011. [DNSIYC UK report ALL chapters DH V10.0.pdf \(publishing.service.gov.uk\)](#)

EFSA (2005a) 'Opinion of the Scientific Panel on contaminants in the food chain [CONTAM] related to the presence of non dioxin-like polychlorinated biphenyls (PCB) in feed and food', EFSA Journal, 3(11), p. 284. doi:10.2903/j.efsa.2005.284. [Opinion of the Scientific Panel on contaminants in the food chain \[CONTAM\] related to the presence of non dioxin-like polychlorinated biphenyls \(PCB\) in feed and food - - 2005 - EFSA Journal - Wiley Online Library](#)

EFSA (2005b) 'Opinion of the Scientific Panel on food additives, flavourings, processing aids and materials in contact with food (AFC) related to Bis(2-ethylhexyl)phthalate (DEHP) for use in food contact materials', EFSA Journal [Preprint], (EFSA Journal). doi:10.2903/j.efsa.2005.243. [Opinion of the Scientific Panel on food additives, flavourings, processing aids and materials in contact with food \(AFC\) related to Bis\(2-ethylhexyl\)phthalate \(DEHP\) for use in food contact materials - - 2005 - EFSA Journal - Wiley Online Library](#)

EFSA (2005c) 'Opinion of the Scientific Panel on food additives, flavourings, processing aids and materials in contact with food (AFC) related to Butylbenzylphthalate (BBP) for use in food contact materials', EFSA Journal [Preprint], (EFSA Journal). doi:10.2903/j.efsa.2005.241. [Opinion of the Scientific Panel on food additives, flavourings, processing aids and materials in contact with food \(AFC\) related to Butylbenzylphthalate \(BBP\) for use in food contact materials - - 2005 - EFSA Journal - Wiley Online Library](#)

EFSA (2005d) 'Opinion of the Scientific Panel on food additives, flavourings, processing aids and materials in contact with food (AFC) related to di-Butylphthalate (DBP) for use in food contact materials', EFSA Journal [Preprint], (EFSA Journal). doi:10.2903/j.efsa.2005.242. [Opinion of the Scientific Panel on food additives, flavourings, processing aids and materials in contact with food \(AFC\) related to di-Butylphthalate \(DBP\) for use in food contact materials - - 2005 - EFSA Journal - Wiley Online Library](#)

EFSA (2005e) 'Opinion of the Scientific Panel on food additives, flavourings, processing aids and materials in contact with food (AFC) related to Di-isodecylphthalate (DIDP) for use in food contact materials', EFSA Journal [Preprint], (EFSA Journal). doi:10.2903/j.efsa.2005.245. [Opinion of the Scientific Panel on food additives, flavourings, processing aids and materials in contact with food \(AFC\) related to Di-isodecylphthalate \(DIDP\) for use in food contact materials - - 2005 - EFSA Journal - Wiley Online Library](#)

EFSA (2005f) 'Opinion of the Scientific Panel on food additives, flavourings, processing aids and materials in contact with food (AFC) related to Di-isononylphthalate (DINP) for use in food contact materials', EFSA Journal [Preprint], (EFSA Journal). doi:10.2903/j.efsa.2005.244. [Opinion of the Scientific Panel on food additives, flavourings, processing aids and materials in contact with food \(AFC\) related to Di-isononylphthalate \(DINP\) for use in food contact materials - - 2005 - EFSA Journal - Wiley Online Library](#)

EFSA (2006) in TOLERABLE UPPER INTAKE LEVELS FOR VITAMINS AND MINERALS: Scientific Committee on Food, Scientific Panel on Dietetic Products, Nutrition and Allergies, pp. 135–150. [complet\\_chapitres.indd \(europa.eu\)](#)

EFSA (2009) 'Nitrite as undesirable substances in animal feed - Scientific Opinion of the Panel on Contaminants in the Food Chain', EFSA Journal, 7(4). doi:10.2903/j.efsa.2009.1017. [Nitrite as undesirable substances in animal feed - Scientific Opinion of the Panel on Contaminants in the Food Chain - - 2009 - EFSA Journal - Wiley Online Library](#)

EFSA (2010) 'Scientific Opinion on Polybrominated Biphenyls (PBBs) in Food'. [Scientific Opinion on Polybrominated Biphenyls \(PBBs\) in Food - - 2010 - EFSA Journal - Wiley Online Library](#)

EFSA (2011a) 'Scientific Opinion on Hexabromocyclododecanes (HBCDDs) in Food', EFSA Journal, 9(7), p. 2296. doi:10.2903/j.efsa.2011.2296. [Scientific Opinion on Hexabromocyclododecanes \(HBCDDs\) in Food - - 2011 - EFSA Journal - Wiley Online Library](#)

EFSA (2011b) 'Scientific Opinion on Tetrabromobisphenol A (TBBPA) and its derivatives in food'. [Scientific Opinion on Tetrabromobisphenol A \(TBBPA\) and its derivatives in food - - 2011 - EFSA Journal - Wiley Online Library](#)

EFSA (2011c) 'Statement on tolerable weekly intake for cadmium', EFSA Journal, 9(2), p. 1975. doi:10.2903/j.efsa.2011.1975. [Statement on tolerable weekly intake for cadmium \(wiley.com\)](#)

EFSA (2012a) 'Cadmium dietary exposure in the European population', EFSA Journal, 10(1), p. 2551. doi:10.2903/j.efsa.2012.2551. [Cadmium dietary exposure in the European population - - 2012 - EFSA Journal - Wiley Online Library](#)

EFSA (2012b) 'Lead dietary exposure in the European population', EFSA Journal, 10(7), p. 2831. doi:10.2903/j.efsa.2012.2831. [Lead dietary exposure in the European population - - 2012 - EFSA Journal - Wiley Online Library](#)

EFSA (2012c) 'Scientific Opinion on the risk for public health related to the presence of mercury and methylmercury in food', EFSA Journal, 10(12), p. 2985. doi:10.2903/j.efsa.2012.2985. [Scientific Opinion on the risk for public health related to the presence of mercury and methylmercury in food - - 2012 - EFSA Journal - Wiley Online Library](#)

EFSA (2014) 'Scientific Opinion on the risks to public health related to the presence of perchlorate in food, in particular fruits and vegetables', EFSA Journal, 12(10), p. 3869. doi:10.2903/j.efsa.2014.3869. [Scientific Opinion on the risks to public health related to the presence of perchlorate in food, in particular fruits and vegetables - - 2014 - EFSA Journal - Wiley Online Library](#)

EFSA (2015a) 'Risks for public health related to the presence of chlorate in food', EFSA Journal, 13(6), p. 4135. doi:10.2903/j.efsa.2015.4135. [Risks for public health related to the presence of chlorate in food - - 2015 - EFSA Journal - Wiley Online Library](#)

EFSA (2015b) 'Scientific Opinion on the risks to public health related to the presence of bisphenol A (BPA) in foodstuffs', EFSA Journal, 13(1), p. 3978. doi:10.2903/j.efsa.2015.3978. [Scientific Opinion on the risks to public health related to the presence of bisphenol A \(BPA\) in foodstuffs - - 2015 - EFSA Journal - Wiley Online Library](#)

EFSA (2017) 'Dietary exposure assessment to perchlorate in the European population', EFSA Journal, 15(10), p. e05043. doi:10.2903/j.efsa.2017.5043. [Dietary exposure assessment to perchlorate in the European population - - 2017 - EFSA Journal - Wiley Online Library](#)

EFSA (2018) 'Risk for animal and human health related to the presence of dioxins and dioxin-like PCBs in feed and food', EFSA Journal, 16(11). doi:10.2903/j.efsa.2018.5333. [Risk for animal and human health related to the presence of dioxins and dioxin-like PCBs in feed and food - - 2018 - EFSA Journal - Wiley Online Library](#)

EFSA (2019) 'Update of the risk assessment of di-butylphthalate (DBP), butyl-benzyl-phthalate (BBP), bis(2-ethylhexyl)phthalate (DEHP), di-isononylphthalate (DINP) and di-isodecylphthalate (DIDP) for use in food contact materials', EFSA Journal, 17(12). doi:10.2903/j.efsa.2019.5838. [Update of the risk assessment of di-butylphthalate \(DBP\), butyl-benzyl-phthalate \(BBP\), bis\(2-ethylhexyl\)phthalate \(DEHP\), di-isononylphthalate \(DINP\) and di-isodecylphthalate \(DIDP\) for use in food contact materials - - 2019 - EFSA Journal - Wiley Online Library](#)

EFSA (2020a) 'Risk assessment of aflatoxins in food', EFSA journal. European Food Safety Authority, 18(3), pp. e06040–e06040. doi:10.2903/j.efsa.2020.6040. [Risk assessment of aflatoxins in food - - 2020 - EFSA Journal - Wiley Online Library](#)

EFSA (2020b) 'Risk to human health related to the presence of perfluoroalkyl substances in food', EFSA Journal, 18(9). doi:10.2903/j.efsa.2020.6223. [Risk to human health related to the presence of perfluoroalkyl substances in food - - 2020 - EFSA Journal - Wiley Online Library](#)

EFSA (2021a) 'Chronic dietary exposure to inorganic arsenic', EFSA Journal, 19(1), p. e06380. doi: /10.2903/j.efsa.2021.6380. [Chronic dietary exposure to inorganic arsenic - - 2021 - EFSA Journal - Wiley Online Library](#)

EFSA (2021b) 'Update of the risk assessment of hexabromocyclododecanes (HBCDDs) in food'. [Update of the risk assessment of hexabromocyclododecanes \(HBCDDs\) in food - - 2021 - EFSA Journal - Wiley Online Library](#)

European Chemicals Agency (ECHA) (2019) 'Annex XV Restriction Report Proposal for a Restriction for intentionally added microplastics'. [Microsoft Word - rest\\_microplastics\\_axvreport\\_en.docx \(europa.eu\)](#)

European Union (EU) (2011) 'Commission Regulation (EU) No 574/2011 of 16 June 2011 amending Annex I to Directive 2002/32/EC of the European Parliament and of the Council as regards maximum levels for nitrite, melamine, Ambrosia spp. and carry-over of certain coccidiostats and histomonostats and consolidating Annexes I and II thereto'. [Commission Regulation \(EU\) No 574/2011 of 16 June 2011 amending Annex I to Directive 2002/32/EC of the European Parliament and of the Council as regards maximum levels for nitrite, melamine, Ambrosia spp. and carry-over of certain coccidiostats and histomonostats and consolidating Annexes I and II theretoText with EEA relevance \(europa.eu\)](#)

EVM (2003) Safe Upper Levels for Vitamins and Minerals. [vitmin2003.pdf \(food.gov.uk\)](#)

FAO/WHO (1989) monograph on iodine. [661. Iodine \(WHO Food Additives Series 24\) \(inchem.org\)](#)

FAO/WHO (2000) Estradiol-17 $\beta$ , progesterone and testosterone in "WHO Food Additives Series 43. Toxicological evaluation of certain veterinary drug residues in food. Prepared by the fifty-second meeting of the Joint FAO/WHO Expert Committee on Food Additives." 43. Geneva. [962. Estradiol-17beta-, progesterone, and testosterone \(WHO Food Additives Series 43\) \(inchem.org\)](#)

FAO/WHO (2016) 'Safety evaluation of certain food additives and contaminants: Supplement 1, Non-dioxin-like polychlorinated biphenyls'. [v71s1je01.pdf \(inchem.org\)](#)

Farebrother, J., Zimmermann, M.B. and Andersson, M. (2019) 'Excess iodine intake: sources, assessment, and effects on thyroid function', *Annals of the New York Academy of Sciences*, 1446(1), pp. 44-65. doi:10.1111/nyas.14041. <https://doi.org/10.1111/nyas.14041>

Fernandes, A. *et al.* (2012) Organic Environmental Contaminants in the 2012 Total Diet Study Samples. [Research report: Total Diet Study: dietary exposure to organic environmental contaminants \(food.gov.uk\)](#)

FSA (2016) Infant Metals Survey. [View PDF \(food.gov.uk\)](#)

Heudorf, U., Mersch-Sundermann, V. and Angerer, J. (2007) 'Phthalates: toxicology and exposure', *International Journal of Hygiene and Environmental Health*, 210(5), pp. 623–634. doi:10.1016/j.ijheh.2007.07.011. <https://doi.org/10.1016/j.ijheh.2007.07.011>

Hill, N.I., Becanova, J. and Lohmann, R. (2021) 'A sensitive method for the detection of legacy and emerging per- and polyfluorinated alkyl substances (PFAS) in dairy milk', *Analytical and Bioanalytical Chemistry* [Preprint]. doi:10.1007/s00216-021-03575-2. <https://doi.org/10.1007/s00216-021-03575-2>

Höjer, A. *et al.* (2012) 'Effects of feeding dairy cows different legume-grass silages on milk phytoestrogen concentration', *Journal of Dairy Science*, 95(8), pp. 4526–4540. doi:10.3168/jds.2011-5226. <https://doi.org/10.3168/jds.2011-5226>

Kowalczyk, J. *et al.* (2013) 'Absorption, Distribution, and Milk Secretion of the Perfluoroalkyl Acids PFBS, PFHxS, PFOS, and PFOA by Dairy Cows Fed Naturally Contaminated Feed', *Journal of Agricultural and Food Chemistry*, 61(12), pp. 2903–2912. doi:10.1021/jf304680j. <https://doi.org/10.1021/jf304680j>

Li, R. *et al.* (2014) 'Postweaning dietary genistein exposure advances puberty without significantly affecting early pregnancy in C57BL/6J female mice', *Reproductive Toxicology (Elmsford, N.Y.)*, 44, pp. 85–92. doi:10.1016/j.reprotox.2013.12.003. <https://doi.org/10.1016/j.reprotox.2013.12.003>

Malekinejad, H., Scherpenisse, P. and Bergwerff, A.A. (2006) 'Naturally Occurring Estrogens in Processed Milk and in Raw Milk (from Gestated Cows)', *Journal of Agricultural and Food Chemistry*, 54(26), pp. 9785–9791. doi:10.1021/jf061972e. <https://doi.org/10.1021/jf061972e>

McCarver, G. *et al.* (2011) 'NTP-CERHR expert panel report on the developmental toxicity of soy infant formula', *Birth Defects Research Part B: Developmental and Reproductive Toxicology*, 92(5), pp. 421–468. doi:10.1002/bdrb.20314. [NTP-CERHR expert panel report on the developmental toxicity of soy infant formula - McCarver - 2011 - Birth Defects Research Part B: Developmental and Reproductive Toxicology - Wiley Online Library](https://doi.org/10.1002/bdrb.20314)

Montgomery, H., Haughey, S.A. and Elliott, C.T. (2020) 'Recent food safety and fraud issues within the dairy supply chain (2015–2019)', *Global Food Security*, 26, p. 100447. doi:10.1016/j.gfs.2020.100447. [Recent food safety and fraud issues within the dairy supply chain \(2015-2019\) - PubMed \(nih.gov\)](https://doi.org/10.1016/j.gfs.2020.100447)

NHS (2018) 'Drinks and cups for babies and young children'. (Accessed: 28 October 2021) Drinks and cups for babies and young children - NHS ([www.nhs.uk](http://www.nhs.uk))

Nordic Council of Ministers (2020) 'Soy intake and possible adverse health effects in Nordic children and pregnant women'. [temanord2020-532.pdf \(norden.org\)](#)

Nørskov, N.P. *et al.* (2019) 'Concentrations of phytoestrogens in conventional, organic and free-range retail milk in England', *Food Chemistry*, 295, pp. 1-9. doi:10.1016/j.foodchem.2019.05.081. [Concentrations of phytoestrogens in conventional, organic and free-range retail milk in England - PubMed \(nih.gov\)](#)

O'Kane, S.M. *et al.* (2018) 'The Effect of Processing and Seasonality on the Iodine and Selenium Concentration of Cow's Milk Produced in Northern Ireland (NI): Implications for Population Dietary Intake', *Nutrients*, 10(3), p. 287. doi:10.3390/nu10030287. [The Effect of Processing and Seasonality on the Iodine and Selenium Concentration of Cow's Milk Produced in Northern Ireland \(NI\): Implications for Population Dietary Intake - PubMed \(nih.gov\)](#)

Payling, L.M. *et al.* (2015) 'Effect of milk type and processing on iodine concentration of organic and conventional winter milk at retail: Implications for nutrition', *Food Chemistry*, 178, pp. 327-330. doi:10.1016/j.foodchem.2015.01.091. [Effect of milk type and processing on iodine concentration of organic and conventional winter milk at retail: implications for nutrition - PubMed \(nih.gov\)](#)

Pesticide Residues in Food (PRiF) (2015) 'The expert committee on Pesticide Residues in Food (PRiF) Annual Report 2015'. [Available at: PDF](#)

Pesticide Residues in Food (PRiF) (2016) 'The expert committee on Pesticide Residues in Food (PRiF) Annual Report 2016'. [Available at: PDF](#)

Pesticide Residues in Food (PRiF) (2017) 'The expert committee on Pesticide Residues in Food (PRiF) Annual Report 2017'. [The Expert Committee on Pesticide Residues in Food - Annual Report 2017 \(publishing.service.gov.uk\)](#)

Pesticide Residues in Food (PRiF) (2018) 'The expert committee on Pesticide Residues in Food (PRiF) Annual Report 2018'. [Available at: PDF](#)

Pesticide Residues in Food (PRiF) (2019) 'The expert committee on Pesticide Residues in Food (PRiF) Annual Report 2019'. [PRiF Annual Report 2019 \(publishing.service.gov.uk\)](#)

Pesticide Residues in Food (PRiF) (2020) 'The expert committee on Pesticide Residues in Food (PRiF) Annual Report 2020'. [Annual Report 2020 \(publishing.service.gov.uk\)](https://publishing.service.gov.uk)

Ráduly, Z. et al. (2020) 'Toxicological and Medical Aspects of Aspergillus-Derived Mycotoxins Entering the Feed and Food Chain', *Frontiers in microbiology*, 10, pp. 2908–2908. doi:10.3389/fmicb.2019.02908. [Toxicological and Medical Aspects of Aspergillus-Derived Mycotoxins Entering the Feed and Food Chain - PubMed \(nih.gov\)](https://pubmed.ncbi.nlm.nih.gov/)

Reijden, O.L. van der, Zimmermann, M.B. and Galetti, V. (2017) 'Iodine in dairy milk: Sources, concentrations and importance to human health', *Best Practice & Research Clinical Endocrinology & Metabolism*, 31(4), pp. 385–395. doi:10.1016/j.beem.2017.10.004. [Iodine in dairy milk: Sources, concentrations and importance to human health - PubMed \(nih.gov\)](https://pubmed.ncbi.nlm.nih.gov/)

Roberts, C. et al. (2018) 'National Diet and Nutrition Survey Results from Years 7 and 8 (combined) of the Rolling Programme (2014/2015 – 2015/2016)'. [National Diet and Nutrition Survey \(publishing.service.gov.uk\)](https://publishing.service.gov.uk)

Scaglioni, P.T. et al. (2014) 'Aflatoxin B<sub>1</sub> and M<sub>1</sub> in milk', *Analytica Chimica Acta*, 829, pp. 68–74. doi:10.1016/j.aca.2014.04.036. [Aflatoxin B<sub>1</sub> and M<sub>1</sub> in milk - PubMed \(nih.gov\)](https://pubmed.ncbi.nlm.nih.gov/)

SCVPH (2002) Opinion of the Scientific Committee on Veterinary Measures relating to Public Health on Review of previous SCVPH opinions of 30 April 1999 and 3 May 2000 on the potential risks to human health from hormone residues in bovine meat and meat products. [Opinion of the SCVPH \(europa.eu\)](https://ec.europa.eu/scvph/)

Socas-Rodríguez, B. et al. (2015) 'Core-shell polydopamine magnetic nanoparticles as sorbent in micro-dispersive solid-phase extraction for the determination of estrogenic compounds in water samples prior to high-performance liquid chromatography-mass spectrometry analysis', *Journal of chromatography A*, 1397, pp. 1–10. doi:10.1016/j.chroma.2015.04.010. [Journal of Chromatography A | ScienceDirect.com by Elsevier](https://www.sciencedirect.com/journal/journal-of-chromatography-a)

The Organisation for Economic Co-operation and Development (OECD) (2021) 'Reconciling Terminology of the Universe of Per- and Polyfluoroalkyl Substances: Recommendations and Practical Guidance'. [Reconciling Terminology of the Universe of Per- and Polyfluoroalkyl Substances: Recommendations and Practical Guidance \(oecd.org\)](https://www.oecd.org/)

Toussaint, B. et al. (2019) 'Review of micro- and nanoplastic contamination in the food chain', Food Additives & Contaminants. Part A, Chemistry, Analysis, Control, Exposure & Risk Assessment, 36(5), pp. 639-673.

doi:10.1080/19440049.2019.1583381. [Full article: Review of micro- and nanoplastic contamination in the food chain \(tandfonline.com\)](#)

Veterinary Medicines Directorate (VMD) (2014) 'Code of Practice on the responsible use of animal medicines on the farm'.

[Responsible use COP April 2015.pdf \(publishing.service.gov.uk\)](#)

Veterinary Medicines Directorate (VMD) (2015) 'National statutory surveillance scheme for veterinary residues in animals and animal products: 2015'.

[Microsoft Word - PCDOCS-#726877-v8-2015\\_Published\\_Results\\_Paper .doc \(publishing.service.gov.uk\)](#)

Veterinary Medicines Directorate (VMD) (2016) 'National statutory surveillance scheme for veterinary residues in animals and animal products: 2016'.

[VETERINARY RESIDUES COMMITTEE \(publishing.service.gov.uk\)](#)

Veterinary Medicines Directorate (VMD) (2017) 'National statutory surveillance scheme for veterinary residues in animals and animal products: 2017'.

Veterinary Medicines Directorate (VMD) (2018) 'National statutory surveillance scheme for veterinary residues in animals and animal products: 2018'.

[National statutory surveillance scheme for veterinary residues in animals and animal products 2018 \(publishing.service.gov.uk\)](#)

Veterinary Medicines Directorate (VMD) (2019) 'National statutory surveillance scheme for veterinary residues in animals and animal products: 2019'. [National statutory surveillance scheme for veterinary residues in animals and animal products 2019 \(publishing.service.gov.uk\)](#)

Veterinary Medicines Directorate (VMD) (2020) 'National statutory surveillance scheme for veterinary residues in animals and animal products: 2019'.

[National statutory surveillance scheme for veterinary residues in animals and animal products: 2020 \(publishing.service.gov.uk\)](#)

Veterinary Products Committee (2006) Risks Associated with the Use of Hormonal Substances in Food-Producing Animals.

[\[ARCHIVED CONTENT\] Other Reports \(nationalarchives.gov.uk\)](#)

Villarrubia-Gómez, P., Cornell, S.E. and Fabres, J. (2018) 'Marine plastic pollution as a planetary boundary threat - The drifting piece in the sustainability puzzle', *Marine Policy*, 96, pp. 213-220. doi:10.1016/j.marpol.2017.11.035.

[Marine plastic pollution as a pla... preview & related info | Mendeley](#)

Wogan, G.N., Paglialunga, S. and Newberne, P.M. (1974) 'Carcinogenic effects of low dietary levels of aflatoxin B1 in rats', *Food and Cosmetics Toxicology*, 12(5), pp. 681-685. doi:10.1016/0015-6264(74)90239-9.

[Carcinogenic effects of low dietary levels of aflatoxin B1 in rats. - Abstract - Europe PMC](#)

Xiao, C.W. (2008) 'Health Effects of Soy Protein and Isoflavones in Humans', *The Journal of Nutrition*, 138(6), pp. 1244S-1249S. doi:10.1093/jn/138.6.1244S. [Health effects of soy protein and isoflavones in humans - PubMed \(nih.gov\)](#)