Sub-statement on the potential risk(s) from exposure to microplastics: Inhalation route

Microplastics - Inhalation route -Research priorities for risk assessment

In this guide

In this guide

- 1. Microplastics Inhalation route Background
- 2. Microplastics Inhalation route Scope and purpose
- 3. Microplastics Inhalation route Analytical detection methodologies
- 4. Microplastics Inhalation route Toxicity
- 5. <u>Microplastics Inhalation route Toxicokinetics</u>
- 6. <u>Microplastics Inhalation route Exposure</u>
- 7. Microplastics Inhalation route Potential new approaches
- 8. Microplastics Inhalation route COT evaluation
- 9. Microplastics Inhalation route Research priorities for risk assessment
- 10. Microplastics Inhalation route -COT conclusions
- 11. <u>Microplastics Inhalation route Abbreviations</u>
- 12. Microplastics Inhalation route References

Data gaps

94. For the inhalation route the significant data gaps include the lack of:

- Harmonised analytical methods for detection of different NMPs during sample collection.
- Methods of detection of NMPs in tissues and their systemic effects.
- Understanding the contribution and effects of different exposure scenarios (*e.g.* indoor and outdoor environments).
- Understanding how different pre-existing lung and other disease states may be involved in any effects from microplastic exposure.

- How available occupational data and on other particle types should be extrapolated to the general population.
- Data on inhalation exposure to NMPs that are resuspended in an indoor environment.

Priorities for risk assessment

95. The COT recommends the following research priorities ranked in order of importance for addressing the data gaps in the potential toxicity of NMPs in humans and suggest a call should be put out to researchers (Table 1). Information in these areas will assist in the future risk assessment of these particles by inhalation and other routes of exposure.

Table 1. Table of future priorities for risk assessment divided into opportunities for improved study design and reporting, as well as research needs.

Opportunities for Ranking Field improved study **Research needs** Order design and reporting Development of reference standards and an appropriate fit for purpose quantification and detection Physicochemical methodologies for 1 properties /Analytical No information. NMPs in different methodologies matrices.

Standardisation of the NMPs used and reported.

		Studies to explore the effect (s) of the same type of NMP on different tissues and of different types of NMP on the same target tissue.	Investigations of the extent to which NMPs with a range of sizes and compositions are assimilated into human tissues and development of techniques capable of identifying the presence of microplastics in the
2	Biological effects	Consistency in the use and reporting of the standards used, the source of the NMPs, clear characterisation of the NMPs used and standardised reporting of dosimetry.	human body (e.g. in biopsies, samples from tissue banks, if possible, histopathology sections; residual controls at point of sampling).

Exploration of steady states.

Standards in			
measuring and			
reporting size of			
particulates.			

Physicochemical properties

Consistency in the use and reporting of the standards used, the source of the NMPs and clear characterisation of the NMPs used.

From studies of particles at the nanoscale, it has been reported that nanoplastics can deposit lower down in the lung and have been shown to translocate across the pulmonary cellular barrier to secondary organs. Further research needed to confirm. More studies needed looking into the potential effects of nanoplastics are needed to understand size related effects.

Studies on the absorption, distribution, digestion and removal (excretion) of different particle types and sizes.

Studies on the Steady states studies. persistence and potential Adsorption, Distribution, accumulation of NMPs Metabolism and in the human body Provide and maintain a **Excretion Toxicokinetics.** and on the extent to data base of which NMPs are information assessing digestible. biodistribution of NMPs.

Evaluation of current dosimetry models for use with NMPs.

Assessment of the degradation of novel/emerging plastic-based materials on the market such as biobased plastics (e.g. bamboo ware, polylactic acid, chitin) and other advanced polymer matrix composite materials during their use and end of life for their contribution to NMPs.

It is unclear whether, and by how much, they already contribute to the burden of NMPs or similar particles.

Research is needed to explore this and apply to future novel materials.

Migration and degradation.

5

Migration/degradationand relevanand analyticalstuff as wemethodology/detection.establish a

Regular assessment of NMPs in water, air and relevant food stuff as well as establish a monitoring programme. Microplastic concentrations in the environment are expected to increase in the future. In addition, increased and widespread use of single-use plastic personal protective equipment due to the COVID-19 pandemic may also be a significant contributing source of plastic pollution.

The quantification methods for microplastic particulate matter are currently limited and can only be estimated, thus improved technology is required. Regular assessment of NMPs in water, air and relevant food stuff as well as establish a monitoring programme. This can then be shared between academia, researchers and government bodies both nationally and internationally.