

**TOX/2022/57**

## **Committee on Toxicity of Chemicals in Food, Consumer Products and the Environment**

### **First draft statement on the potential risk to human health of turmeric and curcumin**

#### **Introduction**

1. The Food Standards Agency (FSA) has been monitoring incidents related to consumption of raw and powdered turmeric and its supplements. In light of these incidents and due to the uncertainties surrounding the composition and possible contamination of these commodities, the Committee on the Toxicity of Chemicals in Food, Consumer Products and the Environment (COT) has been asked to comment on the risk to human health from turmeric and curcumin in their various forms.
2. A discussion paper (TOX/2019/52) was presented to the Committee on 17th September 2019 providing information on the safety of curcumin in supplements and historical turmeric contamination issues, particularly in relation to lead.
3. Two draft statements were presented to the Committee, on the 3<sup>rd</sup> of December 2019 and on the 10<sup>th</sup> of March 2020 which summarised the exposure to raw and powdered turmeric both in the diet and as used in higher quantities for their purported health benefits. The draft statements also covered potential contamination of curcumin and turmeric, which has been associated with adverse health effects in the past.
4. From the COT meeting on the 17<sup>th</sup> of September 2019, it was concluded by members that given past reported contamination issues with turmeric supplements, there would be value in commissioning a chemical analysis of turmeric supplements available on the UK market.

This is a draft statement for discussion.  
It does not reflect the final views of the Committee and should not be cited.

5. A survey of 30 products was undertaken by Fera Science Ltd in Summer 2021. All samples were analysed for the curcuminoids: curcumin, bisdemethoxycurcumin (BDMC) and demethoxycurcumin (DMC) as well as the black pepper derived alkaloid, piperine; and a comprehensive analysis of 69 trace elements which included the heavy metals lead (Pb), mercury (Hg), arsenic (As) and cadmium (Cd). These results were presented in a discussion paper to the COT on the 29<sup>th</sup> of March 2022 (TOX/2022/19).

6. Since the recent 2021 Fera survey, the FSA have commissioned a further 70 turmeric spice powder analyses for Pb. These were included as part of the annual FSA retail chemical safety and imported food surveillance programs.

7. In July 2022 a discussion paper was discussed by COT which covered a review of recent literature on bioavailability impact / pharmacokinetics (PK) with adjuvant compounds delivered in supplements with curcuminoids and an updated review of curcuminoid safety when these supplements are taken long term. The discussion paper in July 2022 also covered a market 'snapshot' of 'novel' supplements, i.e. synthetic curcuminoids or curcuminoids within micro or nanoparticles for marketing claims of 'greater absorption'.

8. From the discussion of July 2022 members highlighted that some of the other 'novel' supplement types such as micellar nano and micro formulations should be looked at in further detail, regarding their pharmacokinetics and therefore their impact on the active chemicals (e.g. curcuminoids) bioavailability. It has been decided that this will form a new discussion paper in the future as its own topic.

9. This statement provided as Annex A summarises the discussions and conclusions of the Committee from 2019 to date regarding turmeric in food and supplement consumption as 'conventional' supplements, i.e. as a curcuminoid / oleoresin extract with or without the adjuvant compound piperine.

This is a draft statement for discussion.  
It does not reflect the final views of the Committee and should not be cited.

10. This statement also includes extra detail on the adulteration of turmeric powders by other Curcuma species, azo dyes and further assessment of trace element data reported after the 30-product survey discussed in the COT meeting of March 2022.

11. This statement references the opinion from the recently published report on 'Adverse effects associated with the consumption of food supplements containing turmeric' by the French Agency for Food, Environmental and Occupational Health & Safety (ANSES). At this moment in time only the headline conclusions can be referenced as the full report is currently only available in French. An update by the Italian health authorities on the topic is also included.

## **Questions for the Committee**

12. The Committee are asked to consider:

- i) Does the Committee have any comments on the structure or content of the draft Statement?
- ii) Given the recent publication of the ANSES opinion, would members like to see a full translation of the ANSES report at a future meeting?
- iii) Is there enough advice in this statement regarding the consumption of supplements for vulnerable groups, e.g. the immunosuppressed, people with existing liver conditions or people taking other medication
- iv) In the medium to long term safety effects of curcumin section, does the committee have an opinion on how the medical trials with ill patients, sometimes involving other treatments, should be

This is a draft statement for discussion.  
It does not reflect the final views of the Committee and should not be cited.

considered? i.e. should they be given equal weighting to other studies in healthy individuals where only curcumin is tested?

## **TOX/2022/57/Annex A**

### **Committee on Toxicity of Chemicals in Food, Consumer Products and the Environment**

#### **First draft statement on the potential risk to human health of turmeric and curcumin**

##### **Introduction**

1. The Food Standards Agency (FSA) has been monitoring incidents related to consumption of raw and powdered turmeric and its supplements. In light of these incidents and due to the uncertainties surrounding the composition and possible contamination of these commodities, the Committee on the Toxicity of Chemicals in Food, Consumer Products and the Environment (COT) has been asked to comment on the risk to human health from turmeric and curcumin in their various forms.

2. Turmeric is the common name for the rhizome (underground stem) of *Curcuma longa* L., a perennial herb cultivated in tropical and subtropical regions of the world. India is the largest producer of turmeric, supplying over 90 % of the world's demand (Olojede et al., 2009). There are approximately 70 varieties of *C. longa* cultivated in India (Sasikumar, 2005). For centuries, turmeric has been widely used for imparting colour and flavour to food, and in Indian and Chinese traditional medicine as a remedy for the treatment of inflammation and other diseases (Ammon and Wahl, 1991).

3. Many of the purported pharmacological properties of turmeric have been attributed to curcumin (chemical name: diferuloylmethane). These properties include antioxidant, analgesic, anti-inflammatory, antiseptic, anticarcinogenic, chemopreventive, chemotherapeutic, antiviral, antibacterial, antifungal and antiplatelet activities (Alok et al., 2015). Curcumin is a polyphenol compound naturally present within turmeric rhizomes. Its derivatives desmethoxycurcumin (DMC) and bisdemethoxycurcumin (BDMC)

This is a draft statement for discussion.  
It does not reflect the final views of the Committee and should not be cited.

are also present within turmeric rhizomes. These compounds are collectively called “curcuminoids”.

4. Due to its purported health benefits, the consumption of curcumin/turmeric supplements is increasingly popular. However, in recent years there have been a number of reports of hepatotoxicity linked to the consumption of curcumin supplements.

5. The FSA’s Novel Foods Team consider turmeric food supplements, comprising of turmeric oleoresin extract or pure curcumin powder, to be novel. These products were not significantly used as a food or food ingredient before 15<sup>th</sup> of May 1997. Therefore, before these products may be placed on the market in the UK or EU as a food or supplement, authorisation, which includes a safety assessment, under the Novel Food Regulation, is required.

6. Curcumin (E 100) is a dicinnamoylmethane dye authorised as a food additive in the EU. It has been evaluated by the Joint FAO/WHO Expert Committee on Food Additives (JECFA), the Scientific Committee on Food (SCF) and the European Food Safety Authority (EFSA). An Acceptable Daily Intake (ADI) of 3 mg/kg bw/d had been established by JECFA in 2004 based on a reproductive toxicity study and this was re-confirmed in the evaluation by EFSA in 2010 (FAO/WHO, 2004a; EFSA Panel on Food Additives and Nutrient Sources added to Food (ANS), 2010).

7. It is the usage that defines if curcuminoids are a food additive or a novel food. If used as a food additive, then this falls under the food additive legislation. When used as a stand-alone ingredient instead of as an additive then it would be novel, and subject to the novel food regulations.

8. It is estimated that supplement intake potentially leads to exposures that are several magnitudes higher than through dietary exposure. Furthermore, synthetic forms of turmeric and curcumin, or conjugation with other chemicals such as piperine, are used to potentially increase absorption,

This is a draft statement for discussion.  
It does not reflect the final views of the Committee and should not be cited.

thus altering the toxicokinetic (TK) profile. The addition of piperine into turmeric supplements is a very common practice. Therefore, the COT in 2020 questioned the relevance of comparing exposures from supplement intake to health-based guidance values for dietary curcumin. It was decided that it would not be appropriate because synthetic forms or adjuvated curcumin, which may be used in supplements, could have altered TK profiles and increased bioavailability. Thus, the levels determined as of low safety concern in food may not be relevant for supplements.

9. A known safety issue with curcumin and/or turmeric is contamination. Contamination with heavy metals for example is a result of either turmeric grown on contaminated soil or intentional adulteration with, for example, lead chromate. Often lead (Pb) chromate, a Pb-based colour, is used to enhance the appearance of turmeric. Other yellow chromate compounds such as zinc, sodium, potassium or strontium chromate could also be used as adulterants. As a result, raw or ground turmeric could potentially contain high levels of Pb or other metals.

10. Turmeric powder can be intentionally or unintentionally adulterated with chemical dyes or powders of other species of *Curcuma* which may be toxic. For example, the powder of *Curcuma zedoaria*, a common adulterant in turmeric powder, is known to be toxic; the high-protein flour of *C. zedoaria* caused 100 % mortality within 6 days when given at 320 g/kg diet to 5 week-old rats (Latif et al., 1979). Furthermore, in supplements, there has been a number of reported cases that involved adulteration with nimesulide, a nonsteroidal anti-inflammatory drug known to cause liver problems.

11. Based on Grand Views Research business intelligence report, the global curcumin market size was approximately \$58 million in 2020 and is forecast towards progression at a global Compound Annual Growth Rate of 16.1% during the forecast period (2020-2028). Europe is the second biggest market and projected to have the fastest estimated compound annual growth rate of 16.7% between 2020-2028. Regarding market shares, pharmaceutical

This is a draft statement for discussion.  
It does not reflect the final views of the Committee and should not be cited.

applications are dominating the curcumin market by revenue and are estimated to have an approximate 50% market share. Food applications were projected to grow by 16% by 2028 (Grand Views Research, 2022).

12. Curcuminoid supplements are generally sold as a capsule containing powder or a turmeric extract, often containing the adjuvant compound piperine. Based on a review of the current curcumin supplement market by FSA risk assessors, counting 'novel' supplement products as an approximate percentage of all products on sale from major high street retailers, supermarkets and one major online retailer the proportion of supplements with 'novel' micro or nano formulations of curcumin came to approximately 10% of the market. This is with a high degree of uncertainty and it is uncertain how popular these products are, i.e. how well they sell compared to the standard supplements. From these novel supplements on the market the largest proportion of products were colloidal suspensions, with the use of micelles (e.g. surfactant phospholipids) to deliver the curcuminoids. Many of these products claim that this increased the bioavailability of the curcuminoids. Supplements containing synthetic curcuminoids were rare, developed potentially for pharmacotherapy (e.g. cancer treatment (He et al., 2018)) rather than general health use.

13. Stohs et al., (2020) review of modified forms of curcumin supplement products states 'micelles, liposomes, phospholipid complexes, microemulsions, nano-emulsions, emulsions, solid lipid nanoparticles, nanostructured lipid carriers, biopolymer nanoparticles and microgels' offer the greatest potential for delivery systems to increase bioavailability. The mechanism being through 'enhancing small intestine permeation, preventing possible degradation in the microenvironment, increasing plasma half-life and enhancing curcumin efficacy. (Stohs et al., 2020).

14. This draft statement summarises the discussions and conclusions of the Committee to date regarding turmeric consumption in food and



This is a draft statement for discussion.  
It does not reflect the final views of the Committee and should not be cited.

supplement consumption as 'conventional' supplements, i.e. as a curcuminoid / oleoresin extract with or without the adjuvant compound piperine.

### Supplements and reported hepatotoxicity

15. Between December 2018 and 20<sup>th</sup> of July 2019, a total of 21 individual cases of acute cholestatic hepatitis "likely to be linked to the consumption of food supplements based on curcumin and piperine" were reported on Italian territory. A total of 18 turmeric supplements have been associated with this hepatitis outbreak, one of which ("Curcuma Liposomal & black pepper" by Nutrimea) was recalled by Belgium's Federal Agency for Food Chain Safety (AFSCA) (Chu, W, 2019).

16. Whilst the AFSCA stated that "the exact source of contamination had not yet been established", an update from Italy's National Institute of Health indicated that "the interdisciplinary group, section dietetics, and the technical committee for animal nutrition and health concluded that, to date, the causes are likely to be related to individual susceptibility, pre-existing alterations, latent hepato-biliary function or even the use of drugs". The institute did not believe the hepatitis was linked to a contamination with heavy metals such as Pb. The Institute adopted a warning for the labelling of the supplements in question (to take effect from 31st December 2019), advising against their use for subjects with altered hepato-biliary function, and recommending medical advice when other medications are being taken. The Institute added that for turmeric powder, which was implicated in one hepatitis case, no particular recommendations were needed especially considering its history of consumption as a food.

17. As reported by Danielles (2022), Italy's Ministry of Health updated its advice regarding labelling of any products derived from *C.longa* due to turmeric's potential hepatotoxicity and persistent recorded adverse effects in the Italian population. All products must now state 'In case of liver, biliary or calculus abnormalities in the biliary tract, the use of this product is not recommended. Do not use during pregnancy and lactation. Do not use for

This is a draft statement for discussion.  
It does not reflect the final views of the Committee and should not be cited.

prolonged periods without consulting your doctor. If you are taking medications, it is advisable to hear the opinion of the doctor.'

18. ANSES (the French Agency for Food, Environmental and Occupational Health & Safety) report that their 'nutrivigilance scheme has received over 100 reports of adverse effects, including 15 reports of hepatitis, potentially related to the consumption of food supplements containing turmeric or curcumin.' This has led the French agency to provide a recent opinion, in June 2022, on turmeric safety (ANSES, 2022).

## **Toxicokinetics**

19. In both humans and animals, curcumin when consumed as a food additive has been shown to have low oral bioavailability.

20. Approximately 75% of the administered dose was excreted in the faeces with negligible amounts appearing in the urine following oral administration of 1 g/kg bw of curcumin in rats (Wahlström and Blennow, 1978). Oral bioavailability is similarly low in humans, due to poor absorption and extensive first-pass metabolism in the intestine and liver (Ireson et al., 2002). Curcuminoids are sensitive to phase II metabolism in the gastrointestinal tract, with glucuronides identified as being the dominant metabolite (Wang et al., 2019).

21. Furthermore, it is also reported that the low bioavailability of curcuminoids is due to low membrane permeability and hence low absorption. With the underlying mechanism relating to an active reflux by P-glycoprotein and a n-octanol/water partition coefficient (log P) of approximately 4 (Jamwal, 2018; Ji et al., 2016). Therefore, the low bioavailability of curcuminoids can be attributed to a fast metabolism, rapid excretion and low membrane permeability.

22. Numerous studies in animals have evaluated the level of curcumin after administration and found that either no curcumin or its metabolites, or

This is a draft statement for discussion.  
It does not reflect the final views of the Committee and should not be cited.

only low levels, were detected in serum or tissue (Ravindranath and Chandrasekhara, 1981; Shen and Ji, 2012).

23. In supplements it is common practice to alter the curcumin product to change its metabolism and enhance its bioavailability, by addressing the metabolism using metabolism inhibitors, membrane permeability or both. This can be achieved with methods such as the use of liposomal curcumin, curcumin nanoparticles (as part of a nano or microemulsion (Liu et al., 2020)), the use of micelles, i.e. a curcumin phospholipid complex and / or the use of synthetic structural analogues of curcumin that are water soluble.

24. The use of adjuvants is currently the most widely adopted modification of turmeric supplements. For example, piperine is the most widely used adjuvant, is the major active ingredient in black pepper, and is a known inhibitor of glucuronidation in the liver and intestine (Di et al., 2015). Hence, piperine may provide a corresponding decrease in the metabolism of curcuminoids. Furthermore, it is reported that piperine may interfere with efflux mechanisms in the epithelial cells that expel compounds back into the intestine for excretion (Chen et al., 2020). This results in a change of permeation properties of the intestine due to the alteration of membrane dynamics (Khajuria et al., 2002) and therefore compounds with low bioavailability due to lower membrane permeability such as curcuminoids can subsequently be better absorbed. Piperine may also directly increase the intestinal absorptive surface due to the induction of the synthesis of proteins associated with cytoskeletal function of the epithelial cells of the intestine (Khajuria et al., 2002). Although the findings from this paper are highly questionable with the use of high doses and small changes reported.

25. In an in vitro study using caco-2 cells Wang et al., (2019) showed that the absorption of curcumin was significantly increased (by approximately 2.5-fold) ( $p < 0.01$ ), when cells were exposed alongside piperine, suggesting piperine could promote the intestinal absorption of curcumin. Although this

This is a draft statement for discussion.  
It does not reflect the final views of the Committee and should not be cited.

study used a co-amorphous formulation that may not be representative of 'real world' use.

26. Piperine has been shown to increase the absorption of curcumin by up to 154% in rats and up to 2,000% in a human study (Shoba et al., 1998). Within this study, piperine was administered at 20 mg/kg concomitantly with curcumin at 2 g/kg to Wistar rats (single dose), and at 20 mg/kg in humans with curcumin at 2 g total (also a single dose). The 'curcumin only' control group returned results < LOD for curcumin in serum for the absorption half-life and the 2000% increase conclusion was derived only from the AUC measurement.

27. There is a lack of any further studies providing direct evidence of piperine enhancing the bioavailability of curcumin, however there are several examples of piperine being used to aid the bioavailability of other compounds. (Bano et al., 1991; Di et al., 2015; Lambert et al., 2004).

28. Based on the in vitro and in vivo studies above, the TK of curcuminoids, taken as a supplement alongside piperine, could be very different to the kinetics when consumed via conventional dietary exposure as a flavouring. However, based on the evidence available it seems unlikely that any difference in consumers would be marked.

29. There is evidence in the recent literature that shows piperine may have less of an effect on bioavailability as an adjuvant compound than previously reported. Fanca-Berthon et al., (2021) undertook a study using a cohort of 30 human volunteers assessing several different curcumin supplement delivery mechanisms. The piperine-curcuminoid dose combination included approximately 15 mg of piperine with 1,500 mg of curcuminoids. The study monitored blood plasma concentrations over a 24-hour period after a single dose and included metabolites of curcuminoids as well as the parent compounds curcumin, BDMC and DMC to provide a 'total curcuminoids' blood plasma concentration. The piperine-curcuminoid

This is a draft statement for discussion.  
It does not reflect the final views of the Committee and should not be cited.

combination did not show any significant differences to the curcuminoid only standard extract, contrary to the Shoba et al., (1998) human study (Fança-Berthon et al., 2021).

30. Fanca-Berthon et al., (2021) gave several potential reasons for the differences in these findings from the Shoba et al., (1998) human study on the effect of piperine bioavailability when provided concomitantly with curcuminoids:

- The Shoba study had only 8 study participants compared to 30 in the Fanca-Berthon et al. study
- Shoba et al. studied a shorter kinetic duration, sampling after 3 hours compared to 24 hours in the Fanca-Berthon study, which found a T<sub>max</sub> of 6 hours for the parent curcuminoid compounds.
- Shoba et al. only measured the single compound curcumin, compared to the metabolites and other related curcuminoids analysed in the Fanca-Berthon study and used for their conclusions, which state: 'Based on the individual quantification of 15 curcuminoid metabolites, this study demonstrated that unconjugated curcumin, DMC, and BDMC represented only 1% of the total plasma curcuminoids following oral administration of a variety of turmeric formulations. Curcumin plasma concentration alone only reached a maximum of 18–21.5 ng/mL in contrast to >400 ng/mL for all metabolites combined.'

31. No other studies could be found in the literature that reported a negative or 'no effect' of piperine when used as an adjuvant compound to aid the bioavailability of curcuminoids. This may be due to the common drawbacks of assessing the peer reviewed literature for negative or 'no effect' conclusions, i.e. these studies are often unreported (Joobar et al., 2012).

This is a draft statement for discussion.  
It does not reflect the final views of the Committee and should not be cited.

## Toxicity

### Derivation of a Health Based Guidance Value (further details provided in Annex B)

32. In 1975, the Scientific Committee for Food (SCF) evaluated curcumin. No ADI was set by the SCF as they considered that curcumin (from natural foods) could be classified as a colour for which an ADI could not be established, because this material is excluded from the provisions of the Community Directive on colours, but which is nevertheless acceptable for use in food (SCF, 1975).

33. In 1995, JECFA on the basis of the NOAEL of 220 mg/kg bw/day in the carcinogenicity study of mice and a safety factor of 200, issued a temporary ADI of 0 - 1 mg/kg bw for curcumin pending the submission of the results of a reproductive toxicity study (FAO/WHO, 1995).

34. In 2004, JECFA withdrew the temporary ADI and established an ADI for curcumin of 0 - 3 mg/kg bw/day based on significant decreases in the average bodyweights of Wistar rat F2 generation pups in a reproductive toxicity study (FAO/WHO, 2004a).

35. In 2010, based on the study used by JECFA, the EFSA ANS panel concluded that the present database supported an ADI of 3 mg/kg bw/day, also based on significant decreases in the average bodyweights of Wistar rat F2 generation pups (EFSA Panel on Food Additives and Nutrient Sources added to Food (ANS), 2010). EFSA also reported human studies where volunteers were exposed to relatively high doses of curcumin either via single dose or for a few months. Based on the results, for dose levels up to 12,000 mg/day, only short-term and semi-chronic adverse effects, such as gastrointestinal effects, headache and rash were observed, but without a clear dose-relationship.

This is a draft statement for discussion.  
It does not reflect the final views of the Committee and should not be cited.

36. In the meeting of the COT on the 10<sup>th</sup> March 2020 Members questioned the relevance of comparing exposures from supplement intake to the ADI for dietary curcumin. It was decided that it would not be appropriate because synthetic forms or adjuvated curcumin, which may be used in supplements, could have altered TK profiles and increased bioavailability. Thus, the levels determined as of low safety concern in food may not be relevant for supplements.

#### Hepatitis and idiosyncratic drug hepatotoxicity (further details provided in Annex B)

37. Hepatitis is the general term for inflammation of the liver. This has a range of clinical presentations varying in duration, severity and eventual outcome. The initial symptoms of hepatitis are often non-specific but in the later stages of the disease the symptoms reflect impairment of various liver functions. Laboratory evidence of liver cell damage can often be detected in asymptomatic patients but significant impact on the synthetic, metabolic and excretory functions of the liver eventually leads to symptoms such as bruising secondary to lack of clotting factors, encephalopathy caused by failure to convert ammonia to urea, and itching when bile salts are deposited in the skin instead of being eliminated in the bile.

38. In the UK, the most common causes of liver injury are fatty infiltration of the liver or viral infection, but toxicants (including alcohol), genetic storage disease and autoimmune processes can also lead to liver damage. In a proportion of patients, no ready explanation can be found for liver damage however severe. Toxicant-induced hepatitis, usually caused by drugs, is common and often resolves when the relevant chemical exposure ceases. In some cases, however, cellular damage is severe and the outcome can be fatal.

39. Identifying a cause for an episode of hepatitis depends upon a knowledge of the history of exposure to chemicals, drugs or contact with sources of hepatitis infection, together with laboratory investigations. Infection

This is a draft statement for discussion.  
It does not reflect the final views of the Committee and should not be cited.

with many of the hepatitic viruses can be identified either by demonstrating an antigenic part of the virus or a specific antibody response to the virus in the blood. Autoimmune disease can be diagnosed from the pattern of antibodies to specific cellular components such as mitochondria and from the clinical picture of other organ involvement. Damaged liver cells tend to leak enzymes into the blood and some clue as to the site of greatest damage within the liver can be gleaned from the pattern of enzymes in the blood, with transaminases, particularly alanine aminotransferase (ALT), being released from damaged parenchymal cells and alkaline phosphatase being released from cells lining the bile ducts.

40. Idiosyncratic drug hepatotoxicity (IDH) occurs in 1/500 to 1/50,000 individuals exposed to a particular drug (the prevalence of idiopathic hepatitis in the community is estimated to be 1/100,000) (Kaplowitz, 2005). IDH has been associated with a variety of pharmaceutical drugs as well as food supplements, notably kava kava. IDH is variable, person specific and occurs for many drugs, but also does not occur for many others. Idiosyncratic events are not caused by the drug itself but by reactions unique to the individual who is exposed to them (Apica and Lee, 2014). However, without the drug there would be no effect. IDH is generally too rare to be detected in clinical trials, though elevated ALT levels may be an indicator. As a general rule, an ALT level greater than three times the upper level of normal is considered to be a sensitive indicator of liver toxicity (the marker is not completely specific since muscle injury may elevate ALT levels). While this is nearly universally described for idiosyncratic liver toxicants, it is not always predictive of overt idiosyncratic toxicity.

41. Two types of IDH occur. Allergic IDH occurs with a short latency period and involves the adaptive immune system. Symptoms may include fever, rash or eosinophilia. It is not well understood why some individuals have or subsequently develop allergic IDH. However, it is likely due to an individual's genetics, body chemistry at a particular time of life, frequent or multiple drug exposures and/or the presence of an underlying disease (American Academy of Allergy Asthma & Immunology, 2020). Also, having an



This is a draft statement for discussion.  
It does not reflect the final views of the Committee and should not be cited.

allergy to one drug predisposes an individual to have an allergy to another unrelated drug. Non-allergic IDH has none of the above features. There can be a long latency period, where there may have been months of normal liver function test results prior to the occurrence of IDH.

### Curcuminoids and hepatotoxicity

42. Full details of the relevant animal and human studies were provided in the COT discussion paper TOX/2019/52 and summarised in Annex B.

43. In short there are a number of animal and human studies covering acute, sub-chronic and chronic toxicity of curcuminoids. There is no key study but there is a weight of evidence for hepatotoxicity being the critical toxicological endpoint, which the COT previously considered.

44. There is evidence that the hepatotoxicity of curcuminoids has a reversible nature, both in rats (Chavalittumrong et al., 2002) and in human cases where liver function tests normalised after ceasing consumption of the supplement (Luber et al., 2019; Lukefahr et al., 2018; Suhail et al., 2020).

45. Between December 2018 and 20th July 2019, a total of 21 individual cases of acute cholestatic hepatitis “likely to be linked to the consumption of food supplements based on curcumin and piperine” were reported on Italian territory. A total of 18 turmeric supplements have been associated with this hepatitis outbreak, one of which (“Curcuma Liposomal & black pepper” by Nutrimea) was recalled by Belgium’s Federal Agency for Food Chain Safety (AFSCA) (Chu, W, 2019).

46. A long-term study undertaken by the US Drug-Induced Liver Injury Network (DILIN, 2022 (currently a poster presentation due for journal publication in February 2023)) between 2004 and 2017 examined the clinical, histologic and human leukocyte antigen (HLA) associations of turmeric related hepatotoxicity in patients. They concluded of the 1,697 recorded drug induced liver injuries, 9 (0.5%) could be attributed to turmeric consumption. Of these 9

This is a draft statement for discussion.  
It does not reflect the final views of the Committee and should not be cited.

cases, turmeric was taken on average 102 days before onset of injury with a range of 30 – 425 days. Eight of the 9 cases caused hepatocellular injury, 5 of the patients required hospitalisation and 1 died of their injuries.

47. ANSES (the French Agency for Food, Environmental and Occupational Health & Safety) report that their 'nutrivigilance scheme has received over 100 reports of adverse effects, including 15 reports of hepatitis, potentially related to the consumption of food supplements containing turmeric or curcumin'. This has led the French agency to provide a recent opinion, in June 2022, on turmeric safety (ANSES, 2022). In their opinion, ANSES advises against 'the consumption of food supplements containing turmeric by people with bile duct disease' and that 'there is a risk of curcumin interacting with certain medications such as anticoagulants, cancer drugs and immunosuppressants.'

48. Pancholi et al., (2021) dispute the potential hepatotoxic effects reported in human studies. They state that only a handful of human cases reporting toxic symptoms, out of millions of humans exposed to curcuminoid supplements daily, is not a significant finding, with those that have symptoms having other personal reasons for these. Conversely, they report curcuminoids have potential hepatoprotective properties. For example, they quote Rahmani et al., (2016) who show a reduced liver fat content and liver transaminases of humans with non-alcoholic fatty liver disease taking 70 mg of curcumin daily for 8 weeks. However, it should be noted that 3 of the authors of the Pancholi et al., (2021) publication are employed by a supplements company.

#### Medium to long term safety of curcuminoids

49. Pancholi et al., (2021) present a 90-day safety study in humans taking supplements containing curcuminoids. Twenty healthy human volunteers were given 380 mg of curcuminoids daily (adjuvanted with fenugreek derived galactomannans as a curcumin-galactomannoside complex) for 90 days. Aspartate aminotransferase (AST), ALT, alkaline phosphatase (ALP), gamma-

This is a draft statement for discussion.  
It does not reflect the final views of the Committee and should not be cited.

glutamyl transferase (GGT) and bilirubin were all within a normal range after the 90-day exposure, indicating no liver effects after this time period.

50. Gupta, Patchva and Aggarwal, (2012) report that 'safety, tolerability, and nontoxicity of curcumin at high doses are well established by human clinical trials'. They reference a study providing 8 g of curcuminoids daily to 21 pancreatic cancer patients for between 3 and 14 months alongside their regular chemotherapy treatment. Adverse effects reported, including, for example, neutropenia and fatigue were attributed to the chemotherapy treatment and/or disease progression rather than the curcuminoid supplement (Kanai et al., 2011).

51. In a double-blind, placebo-controlled study by Petracca et al., (2021) 80 multiple sclerosis patients were enrolled onto a trial, 40 of which received 500 mg of curcumin twice daily for 24 months. Only 53 patients completed the full time period of the trial. No differences in the occurrence of adverse effects were reported in the patient group taking curcumin supplements compared to the controls.

52. Amalraj et al., (2021) describe a double-blind, placebo-controlled study in 30 healthy volunteers (15 volunteers per two groups). The test group were given 500 mg of curcumin daily for 8 weeks. The supplement was provided as an asafoetida (an oleo gum resin) - curcumin complex. No adverse effects were reported in any of the volunteers during the study.

53. There are numerous other similar clinical trials in the literature showing the potential long-term safety of curcuminoid consumption at similar or higher concentrations provided in supplements. For example trials reported by (Appelboom et al., 2014; Haroyan et al., 2018; Nakagawa et al., 2022; Sterzi et al., 2016).

[Contamination of raw, ground turmeric and curcumin supplements with lead \(Pb\)](#)

This is a draft statement for discussion.

It does not reflect the final views of the Committee and should not be cited.

54. Raw turmeric can be contaminated with Pb as a result of either turmeric grown on Pb rich soil or intentional adulteration with lead chromate (Cowell et al., 2017). It has been reported that lead chromate, a Pb-based colour, may be used to enhance the appearance of turmeric (Forsyth et al., 2019). As a result, raw or ground turmeric could potentially contain high levels of Pb. Forsyth et al., (2019) found Pb as high as 1,150 mg/kg in an extensive turmeric survey in Bangladesh.

55. Pb in the body is distributed to the brain, liver, kidney and bones. It is stored in the teeth and bones, where it accumulates over time. Human exposure is usually assessed through the measurement of Pb in blood. Pb in bone is released into blood during pregnancy and becomes a source of exposure to the developing fetus (WHO, 2022).

56. The EFSA Panel on Contaminants in the Food Chain (CONTAM Panel) identified developmental neurotoxicity in young children and cardiovascular effects and nephrotoxicity in adults as the critical effects for the risk assessment of Pb exposure. The respective BMDLs derived from blood Pb levels in µg/L (corresponding dietary intake values in µg/kg bw/d) were: developmental neurotoxicity BMDL<sub>01</sub>, 12 (0.50); effects on systolic blood pressure (SBP) BMDL<sub>01</sub>, 36 (1.50); effects on prevalence of chronic kidney disease BMDL<sub>10</sub>, 15 (0.63). The Panel highlighted that by protecting children, who are far more sensitive, from the developmental effects of Pb, the general population would also be protected from any adverse effects (EFSA Panel on Contaminants in the Food Chain (CONTAM), 2010). In young children, EFSA concluded that a MOE of 10 or greater (of the corresponding dietary intake values) should be sufficient to ensure that there was no appreciable risk of a clinically significant effect on IQ. At lower MOEs, but greater than 1.0, the risk was likely to be low, but not such that it could be dismissed as of no potential concern.

57. After review of the EFSA and JECFA 2011 (FAO/WHO, 2011) evaluations, COT (COT, 2013) agreed that neurodevelopmental effects of Pb represent the most sensitive endpoint for effects in the developing foetus,

This is a draft statement for discussion.  
It does not reflect the final views of the Committee and should not be cited.

whilst also being protective of the other toxicological end points.

### Contamination with other potential adulterants

#### C.zedoaria

58. Turmeric may be adulterated with other curcuma species such as *C. zedoaria*., also known as zedoary or white turmeric (Dhakal et al., 2019). This is a plant in the same genus that has lower curcumin content and cost but produces a higher yield.

59. The literature on the toxicity of *C. zedoaria* is limited. In a subacute toxicity study *C. zedoaria* caused elevation of liver enzymes at levels of 62.5 mg/kg bw and above in mice (Lakshmi et al., 2011). The essential oil of *C. zedoaria* was also found to significantly increase the levels of ALT and ALP at 100 and 200 mg/kg bw/d in pregnant rats treated with the essential oil from Gestation Day (GD) 7 to GD17. The levels of AST were significantly increased at 200 mg/kg bw/d. In the same paper, rat embryos treated ex vivo at doses of 10, 20 and 40 µg/ml with the essential oil exhibited developmental toxicity as observed from changes in yolk sac diameter, crown-rump length, head length, number of somite and score of flexion, heart, fore and hind brain (Zhou et al., 2013). The high-protein flour of *C. zedoaria* caused 100 % mortality within 6 days when given at 320 g/kg diet to 5 week-old rats (Latif et al., 1979). It has been reported that despite causing central nervous system depression and affecting liver enzymes at high doses, at the levels present in turmeric mixes as well as from traditional use in food *C. zedoaria* would not be of concern (Bejar Ezra, 2018). However, quantification of the levels present in adulterated turmeric would be needed in order to establish the risk.

#### Azo dyes

60. The literature reports adulteration of turmeric powders with azo dyes which are prohibited in foods, such as Sudan dye and Metanil yellow (Sasikumar, 2019). These are aromatic compounds containing an azo group (R-N=N-R) and are not permitted as food additives due to their potential

This is a draft statement for discussion.

It does not reflect the final views of the Committee and should not be cited.

carcinogenicity. The International Agency for Research on Cancer (IARC) has classified the azo dyes Sudan I–IV as Class 3 carcinogens (Pan et al., 2012).

61. A number of animal studies have shown potential tumour formation given the exposure of different Sudan dyes in food products (Nisa et al., 2016). Tsuda et al., (2000), from an in vivo comet assay using ddY mice, showed that 17 azo compounds were positive for genotoxicity.

62. The German Federal Institute for Risk Assessment (BfR, 2003) concluded that for the Sudan dyes, given their mechanism of action, a concentration cannot be determined for these chemical compounds at which a carcinogenic action occurs (the 'threshold value'). Therefore, the BfR could not propose a tolerable daily intake (TDI) for the Sudan dyes.

63. EFSA (2005) concluded that there are insufficient toxicology data on any of the illegal azo dyes to undertake a full risk assessment. However, EFSA did state that there is enough experimental evidence to conclude that Sudan I is both genotoxic and carcinogenic.

64. Regarding the yellow azo dyes, there is evidence to suggest they may be neurotoxic and hepatotoxic. Nagaraja and Desiraju, (1993) report that in Wistar rats the administration of Metanil yellow affected the developing and adult brain. 'In the treated rats the amine levels in the hypothalamus, striatum and brain stem were significantly affected, and the changes were not generally reversible even after withdrawal of the dye.'

65. Saxena and Sharma, (2015) describe hepatotoxic effects in Swiss albino rats when administering a mixture of the three yellow azo dyes Metanil yellow, Sunset yellow and Tartrazine. 'Significantly increased concentrations of serum total protein, serum albumin, serum ALP and hepatic malondialdehyde and significantly lowered levels of superoxide dismutase, reduced glutathione and catalase in the liver tissue of treated animals were observed when compared with control animals.' Furthermore, in the treated

This is a draft statement for discussion.  
It does not reflect the final views of the Committee and should not be cited.

groups infiltration of hepatocytes were observed along with necroscopy and vacuolation.

66. In EFSA's 2005 evaluation on consideration of Metanil yellow the committee stated that there was indication of genotoxic activity however concluded there was 'some indication of tumour promoting activity but probably only through enzyme induction.'

67. In 2017 JECFA evaluated the yellow azo dyes Quinoline Yellow and D&C Yellow No. 10. They concluded that due to their similar chemical structures and manufacturing process that it would be acceptable to take the toxicology data for D&C Yellow No. 10 to support conclusions for Quinoline Yellow. Considering the two similar long-term studies on D&C Yellow No. 10 in albino Sprague Dawley rats (Hogan, G.K. and Knezevich, A.L., 1982a, 1982b) a dietary NOAEL of 0.5% (equivalent to 250 mg/kg bw per day), based on effects on body and organ weights was concluded. Using this NOAEL and an uncertainty factor of 100, JECFA established an ADI of 0–3 mg/kg bw for Quinoline Yellow (FAO/WHO, 2017).

## **Exposure assessment**

### **Exposure from food**

68. Curcumin powder is authorised for use as a colouring agent in food (E 100), where its purity is specified as "not less than 90 % total colouring matters" (i.e. curcumin, DMC, and BDMC) (EC, 2008). Directive 94/36/EC states the maximum permitted levels (MPLs) for E 100 in foodstuffs, range from 20 to 500 mg/kg depending on the food item and beverages (which range from 100 to 200 mg/L) (EC, 1994).

69. A previous discussion paper (TOX/2019/52) discussed dietary exposure using turmeric (curcumin) consumption data from the EFSA ANS 2010 evaluation which used data from the UK National Diet and Nutrition Survey (NDNS) 2000 - 2001 and the European (EXPOCHI) project (EFSA ANS, 2010). Further details can be found in Annex B.

This is a draft statement for discussion.  
It does not reflect the final views of the Committee and should not be cited.

70. Dietary curcumin exposure was below the JECFA 2004 ADI of 3 mg/kg bw/day (FAO/WHO, 2004a) for both adults and children when using UK NDNS data and taking the mean exposure using both maximum reported occurrence concentrations and maximum permitted levels.

#### Exposure through turmeric supplements

71. In addition to exposure to curcumin through a normal diet, turmeric supplements can also be taken. These can be bought as 'over the counter' supplements or by 'self-dosing', through consumption of spices in large quantities. The exposures for curcumin and heavy metals from supplements have been calculated using data from a recent survey by Fera (Fera, 2022) and are described in paragraphs 75 – 81.

72. Curcuminoids can be extracted from ground turmeric powder using organic solvents to create a turmeric oleoresin extract. JECFA lists several solvents permitted for extraction: acetone, methanol, ethanol, and isopropanol (FAO/WHO, 2004b). The European Commission, however, has a different list of permitted solvents: acetone, carbon dioxide, ethyl acetate, dichloromethane, n-butanol, methanol, ethanol, and hexane (EC, 2008). According to JECFA specifications, residual solvent concentrations in turmeric oleoresin intended for use in food are limited to 25 mg/kg for hexane, 30 mg/kg for acetone and 50 mg/kg for ethanol, methanol, and isopropanol (FAO/WHO, 2004b). The extraction methodology used affects the curcuminoid content (37-55 %) (Li, et al., 2011), and the essential oil content (< 25%) (Braga et al., 2003) of the turmeric oleoresin.

73. Turmeric oil extract can be prepared in various ways, for example through the treatment of turmeric powder with steam distillation, supercritical CO<sub>2</sub> extraction (Li et al., 2011), or by evaporating the organic solvent of a crude turmeric oleoresin extract (Funk et al., 2010).



This is a draft statement for discussion.  
It does not reflect the final views of the Committee and should not be cited.

74. Curcumin powder can be obtained through the purification of turmeric oleoresin by crystallisation (Li et al., 2011). However, there can be limited commercial availability of authentic samples of pure curcumin, since its separation from DMC and BDMC can be difficult and time consuming. Thus, commercial “pure” curcumin is, in many cases, a mixture of at least these three curcuminoids (Li et al., 2011). For example, a sample of commercial “pure” curcumin (labelled as 94 % purity) was, after HPLC analysis, found to have approximately 70 % purity (Li et al., 2011). In addition, the composition of a sample of commercial “curcumin” was found to be approximately 71.5 % curcumin, 19.4 % DMC, and 9.1 % BDMC (Pfeiffer et al., 2003).

75. As mentioned earlier in this statement and described in paragraphs 22 - 30, in supplements it is common practice to alter the curcumin product to change its metabolism and enhance its bioavailability, by addressing the metabolism, membrane permeability or both. This can be achieved with methods such as the use of liposomal curcumin, curcumin nanoparticles, the use of curcumin phospholipid complex and the use of structural analogues of curcumin that are water soluble.

#### [Assessment after 2021 sample survey - Curcuminoids](#)

76. Previously, in discussion paper TOX/2019/52, a range of supplement information was taken to estimate exposure to curcuminoids. Since these papers were written a sample survey had been commissioned by the FSA and undertaken by Fera Science Ltd in Summer 2021. The final report for this survey can be found on the FSA website (Fera Science Ltd, 2022).

77. Thirty samples were purchased from a variety of sources (online suppliers, large supermarkets and small retailers) and analysed by Fera Science Ltd using mass spectrometry. These consisted of supplements (n=15), ground/powdered turmeric (n=10) and fresh turmeric root (n=5). One of the fresh samples arrived dried.

This is a draft statement for discussion.  
It does not reflect the final views of the Committee and should not be cited.

78. All samples were analysed in duplicate for the curcuminoids: curcumin, BDMC and DMC as well as the black pepper derived alkaloid piperine.

79. Of the supplements sampled, 5 had total curcuminoids over 10%, with one at almost 30%, and one at almost 50% absolute concentration. Of the 5 supplements providing concentrations of total curcuminoids on the label all results were within  $\pm 20\%$  of the stated concentration.

80. From the survey, taking the recommended doses daily according to the supplement's label, exposure concentrations for a 70 kg adult would range from 0.1 to 7 mg/kg bw/day (mean of 1.7 mg/kg bw/day). Taking the supplement that provides 7 mg/kg bw/day for a 70 kg adult, would contribute a further approximate 3-fold increase in exposure to curcuminoids than would be expected from a high dietary exposure (2.6 mg/kg bw/day), highlighted and discussed in Annex B to this statement (Annex B, Table 2.)

81. Ten of the 15 supplements contained detectable concentrations of piperine with 6 of those  $> 1\%$ , which could potentially alter the TK of the curcumin compounds consumed within the same supplement. One of the samples containing piperine did not state this on the label. Three of the supplements contained piperine at approximately 10% or higher.

82. From the powder samples analysed, i.e. where turmeric is sold as a spice ingredient, if these samples were to be taken as a supplement rather than a food ingredient, e.g. at a teaspoon (5 g) a day exposures were all below the ADI of 3 mg/kg bw/day.

#### [Assessment after 2021 sample survey - Heavy Metals and other trace elements](#)

83. From the recent survey undertaken by Fera Science Ltd (Fera Science Ltd, 2022) described above, all samples were analysed for 69 trace elements which included the heavy metals Pb, Hg, As, and Cd.

This is a draft statement for discussion.  
It does not reflect the final views of the Committee and should not be cited.

84. Twenty-nine of the 30 samples tested had heavy metal concentrations of low concern, i.e. below the maximum level (ML) set for supplements by retained EU legislation EC 1881/2006 or below the EU ML set for root spices by the amendment EU 2021/1317 (due to the recent date, this amendment to the EU regulation is not in UK legislation). For supplements, the MLs are 3 mg/kg for Pb, 1 mg/kg for Cd and 0.1 mg/kg for Hg. For root spice powders the EU ML is 1.5 mg/kg for Pb. Arsenic does not have a ML set for supplements, but all concentrations, bar two samples, were below the 0.2 mg/kg ML set for white rice by EU 2015/1006.

85. One sample, a turmeric spice powder, contained a Pb concentration approximately 10 times higher than the majority of other samples analysed, at 2.25 mg/kg. This would be over the amended recent EU ML of 1.5 mg/kg for root spice powders. This sample also had the second highest concentration of Chromium (Cr) at 2.11 mg/kg which may indicate potential adulteration with lead chromate. If this sample was taken as a supplement at, for example, a high estimate of 2.5 teaspoons a day (approximately 7g) the total exposure of lead from this consumption alone would be 0.23 µg/kg bw/day for a 70 kg adult. This is approximately 2 fold lower than the estimated dietary lead exposure of 0.5 µg/kg bw day for the effects of Pb on developmental neurotoxicity, derived from the BMDL<sub>01</sub> of 12 µg/L blood Pb concentration (EFSA Panel on Contaminants in the Food Chain (CONTAM), 2010).

86. When comparing heavy metal results for supplement samples against spice powders and fresh turmeric there were no clear trends or significant differences between the groups when evaluating the results using a students t-test (supplements against powder & fresh samples).

87. On evaluation of the other trace element results from the 30 samples, results that differed by greater than the mean plus 2 x the standard deviation and 5 x the mean of that sample type (i.e. fresh, powder or supplement) are summarised in Annex B. If taking the supplements as described on their

This is a draft statement for discussion.  
It does not reflect the final views of the Committee and should not be cited.

labels, no calculated exposures for any trace elements would exceed any health-based guidance values (HBGVs), where HBGVs exist.

88. Overall, the trace element profile of each sample was variable, this is explained by

a) the different geographical sources of the products and therefore the differing background trace element concentrations from the environment the products were derived

b) The varying chemical nature of the different supplement formulations

89. Since the recent 2021 Fera survey, the FSA in 2022 have commissioned a further 70 turmeric spice powder sample analyses for Pb. All results were < 0.5 mg/kg and of no health concern.

#### Exposure of other potential adulterants

##### C.zedoaria

90. The occurrence data within the literature of adulteration of *C. longa* with *C. zedoaria* is extremely limited. Sasikumar et al., (2004) describe 3 samples of turmeric powder bought from an Indian market that were found to contain more of the contaminant *C. zedoaria* than the expected *C. longa*. Due to the much lower price and easier availability of the potential adulterant *C. zedoaria* (Dhanya et al., 2011), adulteration of products appears to be assumed rather than much direct evidence in the available literature from chemical analysis. Sasikumar (2019) describes wild curcuma species as common adulterants of turmeric products.

##### Azo dyes

91. The occurrence data of Sudan red and Metanil yellow dyes in turmeric powders is primarily from surveys undertaken from Indian and Pakistan spice

This is a draft statement for discussion.  
It does not reflect the final views of the Committee and should not be cited.

markets. Dixit, Khanna and Das, (2008) with a sample size of 15 turmeric powders from an Indian market, report concentrations of adulteration with Sudan dyes between 0.09% and 1.2% and Metanil yellow between 0.15% and 0.46%.

92. Dixit et al., (2009) report results for a much larger survey across Indian markets, analysing for Metanil yellow only. This included 100 branded turmeric powders and 612 non branded products. None of the branded products contained the dye, 17% (105 samples) of the non-branded powders contained Metanil yellow ranging in concentration between 0.1% to 0.85%.

93. In other surveys from the literature , from a survey of 9 non-branded products Rao et al., (2021) found two products contained Metanil yellow at approximately 0.1%. Ullah et al., (2022) from a survey of branded and non-branded turmeric powders from Pakistan markets (total sample numbers unknown) found Sudan dyes between 0.00015% and 0.84% in the non-branded samples only.

94. From a recent FSA survey of 30 turmeric spice products (this included 6 non-branded products) in 2022, no illegal dyes were detected. This included the Sudan red dyes.

## **Risk Characterisation**

### **Curcuminoids**

95. As detailed in Annex B, consumption as part of the normal diet (from its use as an additive and spice) would lead to curcumin exposures that are generally within the ADI of 3 mg/kg bw/day.

96. There is high uncertainty regarding the risk from the intake of raw and powdered turmeric in high quantities for their purported health benefits. The literature review of human studies within the 2019 COT discussion paper (TOX/2019/52) suggests oral intake of curcuminoids in humans is well

This is a draft statement for discussion.  
It does not reflect the final views of the Committee and should not be cited.

tolerated up to doses of 114 mg/kg bw/day, though minor symptoms of nausea or diarrhoea may occur. Longer term clinical studies summarised in this statement suggest that daily consumption of curcuminoids at concentrations at or above those found in supplements result in no adverse effects.

97. With regard to dietary turmeric supplements, a recent 2021 survey shows that a small proportion (2 from 15 samples) of the tested products would lead to high exposure of curcuminoids i.e. exposures at or above the ADI of 3 mg/kg bw/day. COT concluded from previous discussions that this ADI may not be relevant for supplements. This is because as also shown in the 2021 survey, 10 of the 15 supplements analysed contained piperine which may alter the TK of the curcuminoids, potentially increasing the bioavailability and hepatotoxicity. After a review of the current evidence, it is concluded that the claim with many supplements that piperine improves the bioavailability of curcuminoids is sceptical with high uncertainty. The addition of piperine at these relatively low concentrations are a potential promotional or marketing tool from the producers with limited scientific evidence supporting these claims.

98. Regarding the curcuminoid concentrations in supplements and the safety implications of long-term intake of these based on the studies presented, it is concluded that if consumption was based on the label guidance then minor exceedances of the dietary ADI should not pose a significant risk to the population.

99. However, substantial exceedances of the ADI represent a potential health risk to humans. Lower consumption rates also pose the risk of adverse effects if other medicines are being taken concomitantly and particularly in people with underlying conditions such as latent impairment of biliary function.

100. Regarding the recent reports of hepatotoxicity, the Committee has reviewed all available data and have concluded that there is a link to turmeric

This is a draft statement for discussion.  
It does not reflect the final views of the Committee and should not be cited.

because the effects occurred upon challenge and were reversed after withdrawal. The symptoms are consistent with an idiosyncratic drug reaction.

101. Furthermore, some of the other 'novel' supplement types such as micellar nano and micro formulations should be looked at in further detail, in regards their pharmacokinetics and therefore their impact on curcuminoid bioavailability. This further detail is requested regardless that they only potentially make up a likely small percentage of the supplement market, as they may become more popular in the future.

#### Contamination of raw, ground turmeric and curcumin supplements

102. Contamination of raw turmeric with Pb is a result of either turmeric grown on lead rich soil or intentional adulteration with lead chromate. It has been reported that lead chromate, a Pb-based colour, is sometimes used to enhance the appearance of turmeric. As a result, raw or ground turmeric could potentially contain high levels of Pb.

103. After the Italian incident in 2018 and 2019 (see paragraphs 15 to 17), Italy's National Institute of Health did not believe a contaminant was the underlying cause. Furthermore, the recent 2021 survey of 30 turmeric products and the 2022 survey of 70 turmeric spice powder products did not show any Pb concentrations above the ML of 3 mg/kg for supplements, and only one spice powder sample with Pb concentrations at 2.25 mg/kg was above a recent ML set by the EU of 1.5 mg/kg. These data add to the evidence base that the hepatotoxic effects noted with taking turmeric is more likely due to the curcuminoids rather than heavy metal contamination. Therefore, it is concluded that after the results of the recent product surveys, lead (Pb) contamination of turmeric products is likely not the reason for incidents such as the hepatotoxicity reported in the Italian incident. However, it is still uncertain why this incident occurred as a cluster of adverse effects.

104. On evaluation of the other trace element concentrations from the recent study by Fera Science Ltd (Fera Science Ltd, 2022), the metal

This is a draft statement for discussion.  
It does not reflect the final views of the Committee and should not be cited.

concentrations across all of the products tested was extremely variable. This can be attributed to the geographical variation of the source of the products, and / or the other varying additives within supplements. Concentrations of trace elements (metals) that are relatively higher than the average across that product type have been described and evaluated in Annex B to this statement. No calculated exposures for any trace elements if taking the supplements as described on their labels would exceed any HBGVs, where HBGVs exist.

## **Summary and conclusions**

105. Turmeric is the common name for the rhizome (underground stem) of *Curcuma longa* L., a perennial herb cultivated in tropical and subtropical regions of the world.

106. Curcumin (E 100) is a dicinnamoylmethane dye authorised as a food additive in the EU. It has been evaluated by JECFA, the SCF and (EFSA). An ADI of 3 mg/kg bw/d had been established based on a reproductive toxicity study by JECFA in 2004 (FAO/WHO, 2004a) and was re-confirmed in the evaluation by EFSA in 2010.

107. The consumption of turmeric and/or curcumin either raw, powdered or in supplements has become increasingly popular due to the purported health benefits.

108. Curcumin has low bioavailability, however, in supplements, synthetic forms of curcumin or chemical alterations are used to increase its bioavailability, thus potentially altering its toxicity profile. The claim of many supplements that piperine improves the bioavailability of curcuminoids is sceptical with high uncertainty. Ten of the 15 supplements recently surveyed in 2021 contained piperine, 6 of which at > 1% concentration.

109. Consumption of turmeric/curcumin as part of the diet from its use as a food additive or as a spice generally leads to exposures that are below the



This is a draft statement for discussion.  
It does not reflect the final views of the Committee and should not be cited.

dietary ADI. However, when consumed in high quantities for its purported health benefits, or via the intake of supplements, occasional exceedances of the ADI can occur. From a recent curcuminoid survey of 15 supplements and when following the dosage advice on the label, 2 of these would lead to concentrations above the ADI set for dietary exposure. It is concluded that if consumption was based on the label guidance then minor exceedances of the dietary ADI should not pose a significant risk to the population.

110. Regarding the recent reports of hepatotoxicity, the Committee have reviewed all available data and have concluded that, despite the limited data available, there is a link to turmeric consumption because the effects occurred upon challenge were reversed after withdrawal. The symptoms are consistent with an idiosyncratic drug reaction. The occurrence of a possible contaminant (e.g. heavy metals) being the reason for the recent incidents of hepatotoxicity is unlikely.

111. The Committee agree that substantial exceedances of the ADI represent a potential health risk to humans, especially if other medicines are being taken concomitantly and for individuals with altered hepato-biliary function.

112. Other 'novel' supplement types such as micellar nano and micro formulations should be looked at in further detail, in regards their pharmacokinetics and therefore their impact on curcuminoid bioavailability. This is the area of largest concern. This further detail is requested regardless that they only potentially make up a likely small percentage of the supplement market, as they may become more popular in the future.

This is a draft statement for discussion.  
It does not reflect the final views of the Committee and should not be cited.

## List of Abbreviations and Technical terms

ADI	Acceptable Daily Intake
AFSCA	Belgium's Federal Agency for Food Chain Safety
ALP	Alkaline Phosphatase
ALT	Alanine Aminotransferase
AST	Aspartate Aminotransferase
ANS Panel	Scientific Panel on Food Additives and Nutrient Sources Added to Food
ANSES	French Agency for Food, Environmental and Occupational Health & Safety
AUC	Area Under the Curve
As	Arsenic
BDMC	Bisdemethoxycurcumin
BfR	German Federal Institute for Risk Assessment
BMDL	Benchmark Dose Level
Cd	Cadmium
CO <sub>2</sub>	Carbon Dioxide
CONTAM Panel	Panel on Contaminants in the Food Chain
COT	Committee on Toxicity
Cr	Chromium
DILIN	United States Drug-Induced Liver Injury Network
DMC	Desmethoxycurcumin
EC	European Commission
EFSA	European Food Safety Authority
EU	European Union
EXPOCHI	Exposure Assessments For Children In Europe
FAO	Food and Agricultural Organisations of the United Nations
GD	Gestation Day
GGT	Gamma-Glutamyl Transferase
HBGVs	Health-Based Guidance Values
Hg	Mercury
HPLC	High Performance Liquid Chromatography
IARC	International Agency for Research on Cancer

This is a draft statement for discussion.  
 It does not reflect the final views of the Committee and should not be cited.

IDH	Idiosyncratic Drug Hepatotoxicity
JECFA	Joint FAO-WHO Expert Committee Report on Food Additives
mg	Milligrams
mg/kg bw/day	Milligrams per Kilogram Bodyweight per Day
MPL	Maximum Permitted Levels
ML	Maximum Level
NDA Panel	Panel on Nutrition, Novel Foods and Food Allergens
NDS	National Diet and Nutrition Survey
NOAEL	No-Observed-Adverse-Effect Level
Pb	Lead
PK	Pharmacokinetic
SBP	Systolic Blood Pressure
SCF	Scientific Committee for Food
SD	Standard Deviation
TDI	Tolerable Daily Intake
TK	Toxicokinetic
UK	United Kingdom
WHO	World Health Organisation

This is a draft statement for discussion.  
It does not reflect the final views of the Committee and should not be cited.

## References

Alok, A., Singh, I.D., Singh, S., Kishore, M., Jha, P.C., 2015. Curcumin – Pharmacological Actions And its Role in Oral Submucous Fibrosis: A Review. J Clin Diagn Res 9, ZE01–ZE03. [JCDR - Antioxidant, Fibrinolytic property, Herbal medicine, Nanoparticles, Turmeric](#)

Amalraj, A., Varma, K., Jacob, J., Kuttappan, S., 2021. Efficacy and safety of a gut health product (Actbiome) prepared by incorporation of asafoetida-curcumin complex onto the turmeric dietary fiber in the management of gut health and intestinal microflora in healthy subjects: A randomized, double-blind, placebo controlled study. Bioactive Carbohydrates and Dietary Fibre 26, 100280. [Efficacy and safety of a gut health product \(Actbiome\) prepared by incorporation of asafoetida-curcumin complex onto the turmeric dietary fiber in the management of gut health and intestinal microflora in healthy subjects: A randomized, double-blind, placebo controlled study - ScienceDirect](#)

American Academy of Allergy Asthma & Immunology, 2020. Medications and Drug Allergic Reactions. [Medications and Drug Allergic Reactions | AAAAI](#)

Ammon, H.P., Wahl, M.A., 1991. Pharmacology of Curcuma longa. Planta Med 57, 1–7. [Thieme E-Journals - Planta Medica / Abstract \(thieme-connect.de\)](#)

ANSES (AGENCE NATIONALE DE SÉCURITÉ SANITAIRE de l'alimentation, de l'environnement et du travail), 2022. L'évaluation des risques liés à la consommation de compléments alimentaires contenant du curcuma. [AVIS révisé de l'Anses relatif à l'évaluation des risques relatifs à la consommation de compléments alimentaires contenant du curcuma](#)

Apica, B.S., Lee, W.M., 2014. Drug-Induced Liver Injury, in: McManus, L.M., Mitchell, R.N. (Eds.), Pathobiology of Human Disease. Academic Press, San Diego, pp. 1825–1837. [Drug-Induced Liver Injury - ScienceDirect](#)

This is a draft statement for discussion.  
It does not reflect the final views of the Committee and should not be cited.

Appelboom, T., Maes, N., Albert, A., 2014. A New Curcuma Extract (Flexofytol®) in Osteoarthritis: Results from a Belgian Real-Life Experience. *Open Rheumatol J* 8, 77–81. [A New Curcuma Extract \(Flexofytol®\) in Osteoarthritis: Results from a Belgian Real-Life Experience \(openrheumatologyjournal.com\)](#)

Bano, G., Raina, R.K., Zutshi, U., Bedi, K.L., Johri, R.K., Sharma, S.C., 1991. Effect of piperine on bioavailability and pharmacokinetics of propranolol and theophylline in healthy volunteers. *European Journal of Clinical Pharmacology* 41, 615–617. [Effect of piperine on bioavailability and pharmacokinetics of propranolol and theophylline in healthy volunteers | SpringerLink](#)

Bejar Ezra, 2018. Adulteration of turmeric (*Curcuma longa*) root and rhizome, and root and rhizome extracts. *Botanical Adulterants Bulletin*, 2018 (abstract only)

Braga, M.E.M., Leal, P.F., Carvalho, J.E., Meireles, M.A.A., 2003. Comparison of Yield, Composition, and Antioxidant Activity of Turmeric (*Curcuma longa* L.) Extracts Obtained Using Various Techniques. *J. Agric. Food Chem.* 51, 6604–6611. [Comparison of Yield, Composition, and Antioxidant Activity of Turmeric \(Curcuma longa L.\) Extracts Obtained Using Various Techniques | Journal of Agricultural and Food Chemistry \(acs.org\)](#)

Chavalittumrong, P., Chivapat, S., Rattanajarasroj, S., Punyamong, S., Chuthaputti, A., Phisalaphong, C., 2002. Chronic toxicity study of curcuminoids in rats. *The Songklanakarin Journal of Science and Technology* 24, 16.

Chen, S., Li, Q., McClements, D.J., Han, Y., Dai, L., Mao, L., Gao, Y., 2020. Co-delivery of curcumin and piperine in zein-carrageenan core-shell nanoparticles: Formation, structure, stability and in vitro gastrointestinal digestion. *Food Hydrocolloids* 99, 105334. [Co-delivery of curcumin and](#)

This is a draft statement for discussion.  
It does not reflect the final views of the Committee and should not be cited.

[piperine in zein-carrageenan core-shell nanoparticles: Formation, structure, stability and in vitro gastrointestinal digestion - ScienceDirect](#)

Chu, W, 2019. Belgium recall same curcumin-based supplement linked to Italian hepatitis cases. [Belgium recall same curcumin-based supplement linked to Italian hepatitis cases \(nutraingredients.com\)](#)

COT, 2013. Statement on the potential risks from lead in the infant diet. [\[ARCHIVED CONTENT\] COT statement on lead | Food Standards Agency \(nationalarchives.gov.uk\)](#)

Cowell, W., Ireland, T., Vorhees, D., Heiger-Bernays, W., 2017. Ground Turmeric as a Source of Lead Exposure in the United States. Public Health Rep 132, 289–293. [Ground Turmeric as a Source of Lead Exposure in the United States - Whitney Cowell, Thomas Ireland, Donna Vorhees, Wendy Heiger-Bernays, 2017 \(sagepub.com\)](#)

Danielles, S, 2022. Italy prohibits all health claims linked to turmeric, issues warning for labels. URL [Italy prohibits all health claims linked to turmeric, issues warning for labels \(nutraingredients.com\)](#)

Dhakar, S., Schmidt, W.F., Kim, M., Tang, X., Peng, Y., Chao, K., 2019. Detection of Additives and Chemical Contaminants in Turmeric Powder Using FT-IR Spectroscopy. Foods 8, E143. [Foods | Free Full-Text | Detection of Additives and Chemical Contaminants in Turmeric Powder Using FT-IR Spectroscopy \(mdpi.com\)](#)

Dhanya, K., Syamkumar, S., Siju, S., Sasikumar, B., 2011. Sequence characterized amplified region markers: A reliable tool for adulterant detection in turmeric powder. Food Research International 44, 2889–2895. [Sequence characterized amplified region markers: A reliable tool for adulterant detection in turmeric powder - ScienceDirect](#)

This is a draft statement for discussion.  
It does not reflect the final views of the Committee and should not be cited.

Di, X., Wang, X., Di, X., Liu, Y., 2015. Effect of piperine on the bioavailability and pharmacokinetics of emodin in rats. Journal of Pharmaceutical and Biomedical Analysis 115, 144–149. [Effect of piperine on the bioavailability and pharmacokinetics of emodin in rats - ScienceDirect](#)

DILIN, 2022. Liver Injury Associated with Turmeric-a Growing Problem: Nine Cases from the Drug-Induced Liver Injury Network [DILIN] (poster presentation). PowerPoint Presentation (dilin.org) [PowerPoint Presentation \(dilin.org\)](#)

Dixit, S., Khanna, S.K., Das, M., 2008. A Simple 2-Directional High-Performance Thin-Layer Chromatographic Method for the Simultaneous Determination of Curcumin, Metanil Yellow, and Sudan Dyes in Turmeric, Chili, and Curry Powders. Journal of AOAC INTERNATIONAL 91, 1387–1396. [A Simple 2-Directional High-Performance Thin-Layer Chromatographic Method for the Simultaneous Determination of Curcumin, Metanil Yellow, and Sudan Dyes in Turmeric, Chili, and Curry Powders | Journal of AOAC INTERNATIONAL | Oxford Academic \(oup.com\)](#)

Dixit, S., Purshottam, S.K., Khanna, S.K., Das, M., 2009. Surveillance of the quality of turmeric powders from city markets of India on the basis of curcumin content and the presence of extraneous colours. Food Additives & Contaminants: Part A 26, 1227–1231. [Surveillance of the quality of turmeric powders from city markets of India on the basis of curcumin content and the presence of extraneous colours: Food Additives & Contaminants: Part A: Vol 26, No 9 \(tandfonline.com\)](#)

EC, 2008. Commission Directive 2008/128/EC of 22 December 2008 laying down specific purity criteria concerning colours for use in foodstuffs (Codified version) (Text with EEA relevance), OJ L. [EUR-Lex - 32008L0128 - EN - EUR-Lex \(europa.eu\)](#)

This is a draft statement for discussion.  
It does not reflect the final views of the Committee and should not be cited.

EC, 1994. EUROPEAN PARLIAMENT AND COUNCIL DIRECTIVE 94/36/EC of 30 June 1994 on colours for use in foodstuffs. [EUROPEAN PARLIAMENT AND COUNCIL DIRECTIVE 94/36/EC](#)

EFSA, 2005. Opinion of the Scientific Panel on food additives, flavourings, processing aids and materials in contact with food (AFC) to review the toxicology of a number of dyes illegally present in food in the EU. EFSA Journal. [Opinion of the Scientific Panel on food additives, flavourings, processing aids and materials in contact with food \(AFC\) to review the toxicology of a number of dyes illegally present in food in the EU | EFSA \(europa.eu\)](#)

EFSA Panel on Contaminants in the Food Chain (CONTAM), 2010. Scientific Opinion on Lead in Food. EFS2 8. [Scientific Opinion on Lead in Food - - 2010 - EFSA Journal - Wiley Online Library](#)

EFSA Panel on Food Additives and Nutrient Sources added to Food (ANS), 2010. Scientific Opinion on the re-evaluation of curcumin (E 100) as a food additive. EFS2 8. [Scientific Opinion on the re-evaluation of curcumin \(E 100\) as a food additive - - 2010 - EFSA Journal - Wiley Online Library](#)

Fança-Berthon, P., Tenon, M., Bouter-Banon, S.L., Manfré, A., Maudet, C., Dion, A., Chevallier, H., Laval, J., van Breemen, R.B., 2021. Pharmacokinetics of a Single Dose of Turmeric Curcuminoids Depends on Formulation: Results of a Human Crossover Study. The Journal of Nutrition 151, 1802–1816. [Pharmacokinetics of a Single Dose of Turmeric Curcuminoids Depends on Formulation: Results of a Human Crossover Study | The Journal of Nutrition | Oxford Academic \(oup.com\)](#)

FAO/WHO, 2017. Safety evaluation of certain food additives: prepared by the eighty-second meeting of the Joint FAO/WHO Expert Committee on Food Additives (JECFA), World Health Organization, Geneva. [Safety evaluation of](#)



This is a draft statement for discussion.  
It does not reflect the final views of the Committee and should not be cited.

[certain food additives: prepared by the eighty-second meeting of the Joint FAO/WHO Expert Committee on Food Additives \(JECFA\)](#)

FAO/WHO, 2011. Safety evaluation of certain food additives and contaminants, The 74<sup>th</sup> Meeting of the Joint FAO WHO Expert Committee on Food Additives. WHO, Geneva. [Compendium of Food Additive Specifications - Joint FAO/WHO Expert Committee on Food Additives \(JECFA\) 74th Meeting 2011. FAO JECFA Monographs 11](#)

FAO/WHO, 2004a. Evaluation of certain food additives and contaminants. Sixty-first report of the Joint FAO/WHO Expert Committee on Food Additives. WHO Technical Report Series 922. [WHO TRS 922.pdf](#)

FAO/WHO, 2004b. Curcumin Chemical and Technical Assessment. [Microsoft Word - 2004-02-24 CTA 61 Curcumin.doc \(fao.org\)](#)

FAO/WHO, 1995. Evaluation of certain food additives and naturally occurring toxicants (Fourty-fourth report of the Joint FAO/WHO Expert Committee on Food Additives).WHO Technical Report Series, No. 859.  
Federal Institute for Risk Assessment (BfR), 2003. Dyes Sudan I to IV in food. [WHO TRS 859.pdf](#)

Fera Science Ltd, 2022. Turmeric survey for the FSA (FS430403). [Background to the study | Food Standards Agency](#)

Forsyth, J.E., Nurunnahar, S., Islam, S.S., Baker, M., Yeasmin, D., Islam, M.S., Rahman, M., Fendorf, S., Ardoin, N.M., Winch, P.J., Luby, S.P., 2019. Turmeric means “yellow” in Bengali: Lead chromate pigments added to turmeric threaten public health across Bangladesh. Environmental Research 179, 108722. [Turmeric means “yellow” in Bengali: Lead chromate pigments added to turmeric threaten public health across Bangladesh - ScienceDirect](#)

This is a draft statement for discussion.  
It does not reflect the final views of the Committee and should not be cited.

Funk, J.L., Frye, J.B., Oyarzo, J.N., Zhang, H., Timmermann, B.N., 2010. Anti-Arthritic Effects and Toxicity of the Essential Oils of Turmeric (*Curcuma longa* L.). *J Agric Food Chem* 58, 842–849. [Anti-Arthritic Effects and Toxicity of the Essential Oils of Turmeric \(\*Curcuma longa\* L.\) | Journal of Agricultural and Food Chemistry \(acs.org\)](#)

Grand View Research (2022). GVR Report cover Curcumin Market Size, Share & Trends Report Curcumin Market Size, Share & Trends Analysis Report By Application (Pharmaceutical, Food, Cosmetics), By Region (North America, Europe, Asia Pacific, CSA, MEA), And Segment Forecasts, 2020 - 2028. [Curcumin Market Size & Share Analysis Report, 2028 \(grandviewresearch.com\)](#)

Gupta, S.C., Patchva, S., Aggarwal, B.B., 2012. Therapeutic Roles of Curcumin: Lessons Learned from Clinical Trials. *AAPS J* 15, 195–218. [Therapeutic Roles of Curcumin: Lessons Learned from Clinical Trials | SpringerLink](#)

Haroyan, A., Mukuchyan, V., Mkrtychyan, N., Minasyan, N., Gasparyan, S., Sargsyan, A., Narimanyan, M., Hovhannisyanyan, A., 2018. Efficacy and safety of curcumin and its combination with boswellic acid in osteoarthritis: a comparative, randomized, double-blind, placebo-controlled study. *BMC Complementary and Alternative Medicine* 18, 7. [Efficacy and safety of curcumin and its combination with boswellic acid in osteoarthritis: a comparative, randomized, double-blind, placebo-controlled study | BMC Complementary Medicine and Therapies | Full Text \(biomedcentral.com\)](#)

He, Y., Li, W., Hu, G., Sun, H., Kong, Q., 2018. Bioactivities of EF24, a Novel Curcumin Analog: A Review. *Frontiers in Oncology* 8. <https://www.frontiersin.org/article/10.3389/fonc.2018.00614> (Accessed: 7 June 2022). [Frontiers | Bioactivities of EF24, a Novel Curcumin Analog: A Review \(frontiersin.org\)](#)

This is a draft statement for discussion.  
It does not reflect the final views of the Committee and should not be cited.

Hogan, G.K. and Knezevich, A.L., 1982a. Long-term dietary toxicity/carcinogenicity study of D&C Yellow #10 in rats. Bio/dynamics Inc. Project no. 77-1884. Submitted by International Association of Color Manufacturers (IACM) to WHO, January 2016

Hogan, G.K. and Knezevich, A.L., 1982b. Long-term dietary toxicity/carcinogenicity study of D&C Yellow #10 in rats, Bio/dynamics Inc. Project No. 78-2163. Submitted by International Association of Color Manufacturers (IACM) to WHO, January 2016

Ireson, C.R., Jones, D.J.L., Orr, S., Coughtrie, M.W.H., Boocock, D.J., Williams, M.L., Farmer, P.B., Steward, W.P., Gescher, A.J., 2002. Metabolism of the Cancer Chemopreventive Agent Curcumin in Human and Rat Intestine. *Cancer Epidemiol Biomarkers Prev* 11, 105–111.

Jamwal, R., 2018. Bioavailable curcumin formulations: A review of pharmacokinetic studies in healthy volunteers. *Journal of Integrative Medicine* 16, 367–374. [Bioavailable curcumin formulations: A review of pharmacokinetic studies in healthy volunteers - ScienceDirect](#)

Ji, H., Tang, J., Li, M., Ren, J., Zheng, N., Wu, L., 2016. Curcumin-loaded solid lipid nanoparticles with Brij78 and TPGS improved in vivo oral bioavailability and in situ intestinal absorption of curcumin. *Drug Delivery* 23, 459–470. [Full article: Curcumin-loaded solid lipid nanoparticles with Brij78 and TPGS improved in vivo oral bioavailability and in situ intestinal absorption of curcumin \(tandfonline.com\)](#)

Joober, R., Schmitz, N., Annable, L., Boksa, P., 2012. Publication bias: What are the challenges and can they be overcome? *J Psychiatry Neurosci* 37, 149–152. [Publication bias: What are the challenges and can they be overcome? | JPN](#)

Kanai, M., Yoshimura, K., Asada, M., Imaizumi, A., Suzuki, C., Matsumoto, S., Nishimura, T., Mori, Y., Masui, T., Kawaguchi, Y., Yanagihara, K., Yazumi, S.,

This is a draft statement for discussion.  
It does not reflect the final views of the Committee and should not be cited.

Chiba, T., Guha, S., Aggarwal, B.B., 2011. A phase I/II study of gemcitabine-based chemotherapy plus curcumin for patients with gemcitabine-resistant pancreatic cancer. *Cancer Chemother Pharmacol* 68, 157–164. [A phase I/II study of gemcitabine-based chemotherapy plus curcumin for patients with gemcitabine-resistant pancreatic cancer | SpringerLink](#)

Kaplowitz, N., 2005. Idiosyncratic drug hepatotoxicity. *Nat Rev Drug Discov* 4, 489–499. [Idiosyncratic drug hepatotoxicity | Nature Reviews Drug Discovery](#)

Khajuria, A., Thusu, N., Zutshi, U., 2002. Piperine modulates permeability characteristics of intestine by inducing alterations in membrane dynamics: Influence on brush border membrane fluidity, ultrastructure and enzyme kinetics. *Phytomedicine* 9, 224–231. [Piperine modulates permeability characteristics of intestine by inducing alterations in membrane dynamics: Influence on brush border membrane fluidity, ultrastructure and enzyme kinetics - ScienceDirect](#)

Lakshmi, S., Padmaja, G., Remani, P., 2011. Antitumour Effects of Isocurcumenol Isolated from *Curcuma zedoaria* Rhizomes on Human and Murine Cancer Cells. *International Journal of Medicinal Chemistry* 2011, e253962. [Antitumour Effects of Isocurcumenol Isolated from \*Curcuma zedoaria\* Rhizomes on Human and Murine Cancer Cells \(hindawi.com\)](#)

Lambert, J.D., Hong, J., Kim, D.H., Mishin, V.M., Yang, C.S., 2004. Piperine enhances the bioavailability of the tea polyphenol (-)-epigallocatechin-3-gallate in mice. *J Nutr* 134, 1948–1952. [Piperine Enhances the Bioavailability of the Tea Polyphenol \(-\)-Epigallocatechin-3-gallate in Mice | The Journal of Nutrition | Oxford Academic \(oup.com\)](#)

Latif, M.A., Morris, T.R., Miah, A.H., Hewitt, D., Ford, J.E., 1979. Toxicity of shoti (Indian arrowroot: *Curcuma zedoaria*) for rats and chicks. *Br J Nutr* 41, 57–63. [Toxicity of shoti \(Indian arrowroot: \*Curcuma zedoaria\*\) for rats and chicks | British Journal of Nutrition | Cambridge Core](#)

This is a draft statement for discussion.  
It does not reflect the final views of the Committee and should not be cited.

Li Shiyou, Yuan Wei, Deng Guangrui, Wang Ping, Yang Peiyong,, Aggarwal Bharat, 2011. Chemical Composition and Product Quality Control of Turmeric (Curcuma longa L.). TOPHARMCJ 5, 28–54. [Chemical Composition and Product Quality Control of Turmeric \(Curcuma longa L.\) \(benthamopen.com\)](#)

Liu, Z., Smart, J.D., Pannala, A.S., 2020. Recent developments in formulation design for improving oral bioavailability of curcumin: A review. Journal of Drug Delivery Science and Technology 60, 102082. [Recent developments in formulation design for improving oral bioavailability of curcumin: A review - ScienceDirect](#)

Luber, R.P., Rentsch, C., Lontos, S., Pope, J.D., Aung, A.K., Schneider, H.G., Kemp, W., Roberts, S.K., Majeed, A., 2019. Turmeric Induced Liver Injury: A Report of Two Cases. Case Reports Hepatol 2019, 6741213. [Turmeric Induced Liver Injury: A Report of Two Cases \(hindawi.com\)](#)

Lukefahr, A.L., McEvoy, S., Alfafara, C., Funk, J.L., 2018. Drug-induced autoimmune hepatitis associated with turmeric dietary supplement use. BMJ Case Reports bcr-2018-224611. [Drug-induced autoimmune hepatitis associated with turmeric dietary supplement use | BMJ Case Reports](#)

Nagaraja, T.N., Desiraju, T., 1993. Effects of chronic consumption of metanil yellow by developing and adult rats on brain regional levels of noradrenaline, dopamine and serotonin, on acetylcholine esterase activity and on operant conditioning. Food and Chemical Toxicology 31, 41–44. [Effects of chronic consumption of metanil yellow by developing and adult rats on brain regional levels of noradrenaline, dopamine and serotonin, on acetylcholine esterase activity and on operant conditioning - ScienceDirect](#)

Nakagawa, Y., Mori, K., Yamada, S., Mukai, S., Hirose, A., Nakamura, R., 2022. The Oral Administration of Highly-Bioavailable Curcumin for One Year Has Clinical and Chondro-Protective Effects: A Randomized, Double-Blinded,

This is a draft statement for discussion.  
It does not reflect the final views of the Committee and should not be cited.

Placebo-Controlled Prospective Study. *Arthroscopy, Sports Medicine, and Rehabilitation* 4, e393–e402. [The Oral Administration of Highly-Bioavailable Curcumin for One Year Has Clinical and Chondro-Protective Effects: A Randomized, Double-Blinded, Placebo-Controlled Prospective Study - Arthroscopy, Sports Medicine, and Rehabilitation \(arthroscopysportsmedicineandrehabilitation.org\)](#)

Nisa, A., Zahra, N., B., Y., 2016. Sudan dyes and their potential health effects. *Pakistan Journal of biochemistry and molecular biology* 49, 29–35.

Olojede, A.O., Nwokocha, C.C., Akinpelu, A.O., Dalyop, T., 2009. Effect of Variety, Rhizome and Seed Bed Types on Yield of Turmeric (*Curcuma longa* L) under a Humid Tropical Agro-Ecology. *Advances in Biological Research* 3 (1-2): 40-42. [8.xps \(idosi.org\)](#)

Pan, H., Feng, J., He, G.-X., Cerniglia, C.E., Chen, H., 2012. Evaluation of impact of exposure of Sudan azo dyes and their metabolites on human intestinal bacteria. *Anaerobe* 18, 445–453. [Evaluation of impact of exposure of Sudan azo dyes and their metabolites on human intestinal bacteria - ScienceDirect](#)

Pancholi, V., Smina, T.P., Kunnumakkara, A.B., Maliakel, B., Krishnakumar, I.M., 2021. Safety assessment of a highly bioavailable curcumin-galactomannoside complex (CurQfen) in healthy volunteers, with a special reference to the recent hepatotoxic reports of curcumin supplements: A 90-days prospective study. *Toxicology Reports* 8, 1255–1264. [Safety assessment of a highly bioavailable curcumin-galactomannoside complex \(CurQfen\) in healthy volunteers, with a special reference to the recent hepatotoxic reports of curcumin supplements: A 90-days prospective study - ScienceDirect](#)

Petracca, M., Quarantelli, M., Moccia, M., Vacca, G., Satelliti, B., D'Ambrosio, G., Carotenuto, A., Ragucci, M., Assogna, F., Capacchione, A., Lanzillo, R.,

This is a draft statement for discussion.  
It does not reflect the final views of the Committee and should not be cited.

Morra, V.B., 2021. Prospective study to evaluate efficacy, safety and tolerability of dietary supplement of Curcumin (BCM95) in subjects with Active relapsing Multiple Sclerosis treated with subcutaneous Interferon beta 1a 44 mcg TIW (CONTAIN): A randomized, controlled trial. Multiple Sclerosis and Related Disorders 56, 103274. [Prospective study to evaluate efficacy, safety and tolerability of dietary supplement of Curcumin \(BCM95\) in subjects with Active relapsing Multiple Sclerosis treated with subcutaneous Interferon beta 1a 44 mcg TIW \(CONTAIN\): A randomized, controlled trial - Multiple Sclerosis and Related Disorders \(msard-journal.com\)](#)

Pfeiffer, E., Höhle, S., Solyom, A.M., Metzler, M., 2003. Studies on the stability of turmeric constituents. Journal of Food Engineering 56, 257–259.

Rahmani, S., Asgary, S., Askari, G., Keshvari, M., Hatamipour, M., Feizi, A., Sahebkar, A., 2016. Treatment of Non-alcoholic Fatty Liver Disease with Curcumin: A Randomized Placebo-controlled Trial. Phytotherapy Research 30, 1540–1548. [Treatment of Non-alcoholic Fatty Liver Disease with Curcumin: A Randomized Placebo-controlled Trial - Rahmani - 2016 - Phytotherapy Research - Wiley Online Library](#)

Rao, S.N., Vennapusa, C.S.R., Patel, S., Meti, S., Huggar, B., 2021. Determination of banned adulterants in turmeric and chilli powders using ultra-high-performance liquid chromatography. Journal of Liquid Chromatography & Related Technologies 44, 235–243. [Determination of banned adulterants in turmeric and chilli powders using ultra-high-performance liquid chromatography: Journal of Liquid Chromatography & Related Technologies: Vol 44, No 3-4 \(tandfonline.com\)](#)

Ravindranath, V., Chandrasekhara, N., 1981. Metabolism of curcumin--studies with [3H]curcumin. Toxicology 22, 337–344. [Metabolism of curcumin--studies with \[3H\]curcumin - ScienceDirect](#)

This is a draft statement for discussion.  
It does not reflect the final views of the Committee and should not be cited.

Sasikumar, B., 2019. Advances in adulteration and authenticity testing of turmeric (*Curcuma longa* L.). *J Spices Arom Crops* 96–105. [Advances in adulteration and authenticity testing of turmeric \(Curcuma longa L.\) | Journal of Spices and Aromatic Crops \(updatepublishing.com\)](#)

Sasikumar, B., 2005. Genetic resources of *Curcuma*: diversity, characterization and utilization. *Plant Genetic Resources* 3, 230–251. [Genetic resources of Curcuma: diversity, characterization and utilization | Plant Genetic Resources | Cambridge Core](#)

Sasikumar, B., Syamkumar, S., Remya, R., John Zachariah, T., 2004. PCR Based Detection of Adulteration in the Market Samples of Turmeric Powder. *Food Biotechnology* 18, 299–306. [PCR Based Detection of Adulteration in the Market Samples of Turmeric Powder: Food Biotechnology: Vol 18, No 3 \(tandfonline.com\)](#)

Saxena, B., Sharma, S., 2015. Food Color Induced Hepatotoxicity in Swiss Albino Rats, *Rattus norvegicus*. *Toxicol Int* 22, 152–157. [Food Color Induced Hepatotoxicity in Swiss Albino Rats, Rattus norvegicus - PubMed \(nih.gov\)](#)

SCF (Scientific Committee for Food), 1975. Reports from the Scientific Committee for Food (1st series), opinion expressed 27 June 1975. [REPORTS OF THE SCIENTIFIC COMMITTEE FOR FOOD : First series \(pitt.edu\)](#)

Shen, L., Ji, H.-F., 2012. The pharmacology of curcumin: is it the degradation products? *Trends Mol Med* 18, 138–144. [The pharmacology of curcumin: is it the degradation products?: Trends in Molecular Medicine \(cell.com\)](#)

Shoba, G., Joy, D., Joseph, T., Majeed, M., Rajendran, R., Srinivas, P.S., 1998. Influence of piperine on the pharmacokinetics of curcumin in animals and human volunteers. *Planta Med* 64, 353–356. [Thieme E-Journals - Planta Medica / Abstract \(thieme-connect.de\)](#)



This is a draft statement for discussion.  
It does not reflect the final views of the Committee and should not be cited.

Sterzi, S., Giordani, L., Morrone, M., Lena, E., Magrone, G., Scarpini, C., Milighetti, S., Pellicciari, L., Bravi, M., Panni, I., Ljoka, C., Bressi, F., Foti, C., 2016. The efficacy and safety of a combination of glucosamine hydrochloride, chondroitin sulfate and bio-curcumin with exercise in the treatment of knee osteoarthritis: a randomized, double-blind, placebo-controlled study. *Eur J Phys Rehabil Med* 52, 321–330.

Stohs, S.J., Chen, O., Ray, S.D., Ji, J., Bucci, L.R., Preuss, H.G., 2020. Highly Bioavailable Forms of Curcumin and Promising Avenues for Curcumin-Based Research and Application: A Review. *Molecules* 25, 1397. [Molecules | Free Full-Text | Highly Bioavailable Forms of Curcumin and Promising Avenues for Curcumin-Based Research and Application: A Review \(mdpi.com\)](#)

Suhail, F.K., Masood, U., Sharma, A., John, S., Dhamoon, A., 2020. Turmeric supplement induced hepatotoxicity: a rare complication of a poorly regulated substance. *Clin Toxicol (Phila)* 58, 216–217. [Turmeric supplement induced hepatotoxicity: a rare complication of a poorly regulated substance: Clinical Toxicology: Vol 58, No 3 \(tandfonline.com\)](#)

Tsuda, S., Matsusaka, N., Madarame, H., Ueno, S., Susa, N., Ishida, K., Kawamura, N., Sekihashi, K., Sasaki, Y.F., 2000. The comet assay in eight mouse organs: results with 24 azo compounds. *Mutation Research/Genetic Toxicology and Environmental Mutagenesis* 465, 11–26. [The comet assay in eight mouse organs: results with 24 azo compounds - ScienceDirect](#)

Ullah, A., Chan, M.W.H., Aslam, S., Khan, A., Abbas, Q., Ali, Shamsheer, Ali, M., Hussain, A., Mirani, Z.A., Sibte-e-Hassan, S., Kazmi, M.R., Ali, Shaukat, Hussain, S., Khan, A.M., 2022. Banned Sudan dyes in spices available at markets in Karachi, Pakistan. null 1–9. [Banned Sudan dyes in spices available at markets in Karachi, Pakistan: Food Additives & Contaminants: Part B: Vol 0, No 0 \(tandfonline.com\)](#)

This is a draft statement for discussion.  
It does not reflect the final views of the Committee and should not be cited.

Wahlström, B., Blennow, G., 1978. A study on the fate of curcumin in the rat. Acta Pharmacol Toxicol (Copenh) 43, 86–92. [A Study on the Fate of Curcumin in the Rat - Wahlström - 1978 - Acta Pharmacologica et Toxicologica - Wiley Online Library](#)

Wang, R., Han, J., Jiang, A., Huang, R., Fu, T., Wang, L., Zheng, Q., Li, W., Li, J., 2019. Involvement of metabolism-permeability in enhancing the oral bioavailability of curcumin in excipient-free solid dispersions co-formed with piperine. International Journal of Pharmaceutics 561, 9–18. [Involvement of metabolism-permeability in enhancing the oral bioavailability of curcumin in excipient-free solid dispersions co-formed with piperine - ScienceDirect](#)

WHO, 2017. Lead poisoning. [Lead poisoning \(who.int\)](#)

Zhou, L., Zhang, K., Li, J., Cui, X., Wang, A., Huang, S., Zheng, S., Lu, Y., Chen, W., 2013. Inhibition of vascular endothelial growth factor-mediated angiogenesis involved in reproductive toxicity induced by sesquiterpenoids of Curcuma zedoaria in rats. Reprod Toxicol 37, 62–69. [Inhibition of vascular endothelial growth factor-mediated angiogenesis involved in reproductive toxicity induced by sesquiterpenoids of Curcuma zedoaria in rats - ScienceDirect](#)

This is a draft statement for discussion.  
It does not reflect the final views of the Committee and should not be cited.

## **TOX/2022/57 Annex B**

### **Committee on Toxicity of Chemicals in Food, Consumer Products and The Environment**

#### **First draft statement on the potential risk to human health of turmeric and curcumin**

##### **Introduction**

1. The information presented in this annex should be read in conjunction with the main draft statement on the potential risk to human health of turmeric and curcumin supplements. It contains further detailed information relating to past COT discussions on the safety of curcuminoids and turmeric. This includes discussion on the derivation of a Health Based Guidance Value for curcumin, hepatitis, idiosyncratic drug hepatotoxicity and curcuminoids in relation to toxicity covering both animal and human studies. Detailed exposure data for turmeric in food and evaluation of trace element contamination from a recent survey is also included.

##### **Toxicity**

###### [Derivation of a Health Based Guidance Value](#)

2. In 1975, the SCF evaluated curcumin (chemical name: diferuloylmethane). No ADI was set by SCF as they considered that curcumin (from natural foods) could be classified as colour for which an ADI could not be established but which is nevertheless acceptable for use in food (SCF, 1975).

3. In 1995, JECFA evaluated the results of toxicology and carcinogenicity studies in rats and mice administered turmeric oleoresin containing 79-85 % curcuminoids (National Toxicology Program (NTP), 1993). After 15 months of treatment, absolute and relative liver weights were increased in both male and female mice in the mid- and highest-dose groups relative to controls. The No Observed Adverse Effect Level (NOAEL) for liver enlargement was 2000

This is a draft statement for discussion.  
It does not reflect the final views of the Committee and should not be cited.

mg/kg in the diet, equal to 220 mg/kg bw/ day. On the basis of the NOAEL of 220 mg/kg bw/day in the carcinogenicity study of mice and a safety factor of 200, JECFA established a temporary ADI of 0-1 mg/ kg bw and extended it, pending the submission of the results of a reproductive toxicity study with curcumin (FAO/WHO, 1995).

4. In 2004, JECFA noted that the turmeric oleoresin used in the NTP (1993) study did not comply with the current specification for curcumin. JECFA withdrew the temporary ADI and established an ADI for curcumin of 0-3 mg/kg bw based on significant decreases in the average bodyweights of Wistar rat F2 generation pups in a reproductive toxicity study (FAO/WHO, 2004).

5. In 2010, based on the study used by JECFA, the EFSA ANS panel concluded that the present database supported an ADI of 3 mg/kg bw/day, also based on significant decreases in the average bodyweights of Wistar rat F2 generation pups (EFSA Panel on Food Additives and Nutrient Sources added to Food (ANS), 2010). In this study, Wistar rats were fed diets containing curcumin (> 90 % purity) at doses equal to 0, 130-140, 250-290 and 850-960 mg/kg bw/day in males, and 0, 160, 310-320 and 1000-1100 mg/kg bw/day in females. The total period of treatment was 21 weeks for the parental generation and 24 weeks for the F1 generation. No treatment-related clinical signs of toxicity during the study were reported.

6. Significant decreases in the average weights of the F2 generation pups were observed at days 1 and 7 at the intermediate dose, and on days 7, 14 and 21 at the high dose. These decreases represented < 10 % of the average weight of the concurrent controls and were reported to be within the range of the historical control data. There were no other effects on general health, body weight, pup survival or fertility indices in either generation. JECFA considered the effect on pup weight seen at the intermediate dose (equal to 250-320 mg/kg bw/day) to be incidental and therefore a NOEL. JECFA allocated an ADI for curcumin of 0-3 mg/kg bw/day based on the application of an uncertainty factor of 100 to this NOEL (FAO/WHO, 2004).

This is a draft statement for discussion.  
It does not reflect the final views of the Committee and should not be cited.

7. EFSA ANS panel (2010) also reported human studies where volunteers were exposed to relatively high doses of curcumin either via single dose or for a few months. Based on the results, for dose levels up to 12,000 mg/day, only short-term and semi-chronic adverse effects, such as gastrointestinal effects, headache and rash were observed, but without clear dose-relationship.

### Hepatitis

8. Hepatitis is the general term for inflammation of the liver. This has a range of clinical presentations varying in duration, severity and eventual outcome. The initial symptoms of hepatitis are often non-specific but in the later stages of the disease the symptoms reflect impairment of various liver functions. Laboratory evidence of liver cell damage can often be detected in asymptomatic patients but significant impact on the synthetic, metabolic and excretory functions of the liver eventually leads to symptoms such as bruising secondary to lack of clotting factors, encephalopathy caused by failure to convert ammonia to urea, and itching when bile salts are deposited in the skin instead of being eliminated in the bile. The liver has a remarkable ability to regenerate after damage but often fails to replicate the original complex cellular architecture necessary for normal function and instead produces cirrhosis, a combination of fibrous tissue and regenerative nodules.

9. In the UK, the most common causes of liver injury are fatty infiltration of the liver or viral infection, but toxicants (including alcohol), genetic storage disease and autoimmune processes can also lead to liver damage. In a proportion of patients, no ready explanation can be found for liver damage however severe. Toxicant-induced hepatitis, usually caused by drugs, is common and often resolves when the relevant chemical exposure ceases. In some cases, however, cellular damage is severe and the outcome can be fatal.

This is a draft statement for discussion.  
It does not reflect the final views of the Committee and should not be cited.

10. Identifying a cause for an episode of hepatitis depends upon a history of exposure to chemicals, drugs or contact with sources of hepatitis infection, together with laboratory investigations. Infection with many of the hepatitic viruses can be identified either by demonstrating an antigenic part of the virus or a specific antibody response to the virus in the blood. Autoimmune disease can be diagnosed from the pattern of antibodies to specific cellular components such as mitochondria and from the clinical picture of other organ involvement. Damaged liver cells tend to leak enzymes into the blood and some clue as to the site of greatest damage within the liver can be gleaned from the pattern of enzymes in the blood, with transaminases, particularly alanine aminotransferase (ALT), being released from damaged parenchymal cells and alkaline phosphatase being released from cells lining the bile ducts.

#### Morphology of hepatitis

11. The morphological appearances of different types of hepatitis are often similar (Ferrell, 2000). Pathological features of acute hepatitis include swelling and ballooning of hepatocytes and cell death affecting single cells, groups of cells adjacent to portal tracts, or extensive confluent areas. Kupffer cells are actively phagocytic and within the portal tracts there are increased numbers of chronic inflammatory cells. There may also be increased numbers of inflammatory cells in the hepatic parenchyma.

12. The defining feature of active chronic hepatitis is infiltration of lymphocytes from portal tracts with associated death of liver cells, so called interface hepatitis. This in time is associated with fibrosis. Sometimes the amount of inflammation is less, and a biopsy fails to show interface hepatitis. The presence of plasma cells or discrete lymphoid aggregates may suggest the possibility of a viral cause. Some storage disorders, for example Wilson's disease and copper accumulation, and alpha 1 antitrypsin deficiency, show morphological evidence of a chronic active hepatitis. (Ferrell, 2000)

#### Idiosyncratic drug hepatotoxicity

This is a draft statement for discussion.  
It does not reflect the final views of the Committee and should not be cited.

13. Idiosyncratic drug hepatotoxicity (IDH) occurs in 1/500 to 1/50,000 individuals exposed to a particular drug (the prevalence of idiopathic hepatitis in the community is estimated to be 1/100,000) (Kaplowitz, 2005). IDH has been associated with a variety of pharmaceutical drugs as well as food supplements, notably kava kava. IDH is variable, person specific and occurs for many drugs, but also does not occur for many others. Idiosyncratic events are not caused by the drug itself but by reactions unique to the individual who is exposed to them (Apica and Lee, 2014). However, without the drug there would be no effect. IDH is generally too rare to be detected in clinical trials, though elevated ALT levels may be an indicator. As a general rule, an ALT level greater than three times the upper level of normal is considered to be a sensitive indicator of liver toxicity (the marker is not completely specific since muscle injury may elevate ALT levels). While this is nearly universally described for idiosyncratic liver toxicants, it is not always predictive of overt idiosyncratic toxicity.

14. Two types of IDH occur. Allergic IDH occurs with a short latency period and involves the adaptive immune system. Symptoms may include fever, rash or eosinophilia. It is not well understood why some individuals have or subsequently develop allergic IDH. However, it is likely due to an individual's genetics, body chemistry at a particular time of life, frequent or multiple drug exposures and/or the presence of an underlying disease (American Academy of Allergy Asthma & Immunology, 2020). Also, having an allergy to one drug predisposes an individual to have an allergy to another unrelated drug. Non-allergic IDH has none of the above features. There can be a long latency period, where there may have been months of normal liver function test results prior to the occurrence of IDH.

## **Curcuminoids and hepatotoxicity: Animal studies**

### [Subacute toxicity](#)

#### Rats

This is a draft statement for discussion.

It does not reflect the final views of the Committee and should not be cited.

15. Liju et al. (2013) investigated the acute and sub-chronic toxicity of turmeric essential oil (TEO) from *Curcuma longa* L. Acute administration of TEO was done as a single oral dose of up to 5 g/kg bw, and a 13-week sub-chronic toxicity study was done at doses of 0.1, 0.25 and 0.5 g/kg bw/day in Wistar rats. There were no mortalities, adverse clinical signs or changes in body weight, water and food consumption during the acute and sub-chronic studies. Indicators of hepatic function such as aspartate aminotransferase (AST), ALT and alkaline phosphatase (ALP) were unchanged in treated animals compared to untreated animals. Oral administration of TEO for 13 weeks did not alter total cholesterol, triglycerides, markers of renal function, serum electrolyte parameters and histopathology of tissues.

#### Mice

16. Dietary turmeric powder (0.2 %, 1.0 %, 5.0 %, equivalent to 400, 2000, 10000 mg/kg bw/day) or ethanolic turmeric extract (ETE, 0.05 %, 0.25 %, equivalent to 100, 500 mg/kg bw/day) for 14 days, at doses reported to be cancer preventive in model systems, were found to be hepatotoxic in mice. The diferuloylmethane contents of the turmeric powder and ETE were not reported. Exposure of mice to dietary turmeric or ETE did not have any significant effect on body weight/ liver weight or liver to body weight ratios. Animals exposed to 0.2 % turmeric showed coagulative necrosis in liver. Livers from mice receiving 1.0 % and 5.0 % turmeric showed extreme degenerative changes with necrosis. Coagulative necrotic foci surrounded by a zone of regeneration were also evident. Similar changes were also seen in animals treated with 0.05 % and 0.25 % ETE. Necrotic changes in liver, the principal effect, was seen in 6/6 animals from 5.0 % turmeric, 3/6 from 1.0 % turmeric and 3/6 animals from 0.2 % turmeric. Similarly, 3/6 animals from 0.05 % as well as 0.25 % ETE also showed these alterations. Liver from mice receiving control diet showed normal ultrastructure. In the liver of the animals receiving 5.0 % or 1.0 % turmeric diet or 0.25 % or 0.05 % ETE, some of the parenchymal cells had round nuclei consisting of clumped or densely margined chromatin. The cytoplasm consisted of numerous pleomorphic vacuolated mitochondria filled with dense bodies, surrounded by rough or



This is a draft statement for discussion.  
It does not reflect the final views of the Committee and should not be cited.

smooth endoplasmic reticulum and free ribosomes, and an increased number of glycogen particles and Golgi complexes with vesicles (Kandarkar et al., 1998).

### Sub-chronic toxicity

#### Rats

17. In a 28 day study, curcumin (purity > 98 %) was administered to Sprague Dawley rats through the diet at dose levels of 26.1, 84.8, 224.8, 459.7 and 1117.8 mg/kg bw/day. Clinical chemistry did not reveal major signs of liver damage associated with administration of diferuloylmethane. cDNA microarray experiments were performed on hepatic RNA. Diferuloylmethane altered the expression of 12 genes. Three of these were related to peroxisomes (phytanoyl-CoA dioxygenase, enoyl-CoA hydratase; CYP4A3). Increased cyanide insensitive palmitoyl-CoA oxidation was observed. The authors concluded that these data suggest that diferuloylmethane is a weak peroxisome proliferator in rats (Stierum et al., 2008).

18. In a 13 week study, groups of 10 male and 10 female F344/N rats were fed diets containing 0, 1000, 5000, 10000, 25000, or 50000 ppm turmeric oleoresin (79 to 85 % diferuloylmethane), estimated to deliver average daily doses of 50, 250, 480, 1300, or 2600 mg/kg bw to males and 60, 300, 550, 1450, or 2800 mg/kg bw to females. All rats survived until the end of the study. The final mean body weight of males receiving 50,000 ppm was 5 % lower than that of the controls. Feed consumption by exposed male and female rats was similar to that of controls. The absolute and relative liver weights of female rats and the relative liver weights of male rats receiving 5000, 10,000, 25,000, and 50,000 ppm were significantly greater than those of the controls. According to the NTP, these increases may have been due to mild hepatocellular swelling or hypertrophy. In the clinical chemistry, urinalysis, and other hematologic parameters, no differences were observed that were considered by the NTP to be biologically significant. Clinical findings included stained fur, and discoloured faeces and urine of exposed animals. Mild to moderate hyperplasia of the mucosal epithelium was observed in the

This is a draft statement for discussion.

It does not reflect the final views of the Committee and should not be cited.

cecum and colon of male and female rats that received 50000 ppm (NTP 1993).

19. A six-month toxicity study of curcuminoids extracted from the powdered dried rhizome of *Curcuma longa* was performed in six groups of 15 Wistar rats of each sex. The extract was reported to contain 58-67 % curcuminoids. The water control group received 5 ml of water/kg bw/day, while the tragacanth (plant-derived gum) control group received 5 ml of 0.5 % tragacanth suspension/ kg bw/day orally. Three treatment groups were given the suspension of curcuminoids powder at doses of 10, 50 and 250 mg/kg bw/day. The fourth treatment group, or the recovery group, also received 250 mg/kg bw/day of curcuminoids for six months, but two weeks of no curcuminoids treatment elapsed before the time of sacrifice. It was found that the growth rate of male rats receiving curcuminoids at 50 mg/kg bw/day was significantly higher than that of the tragacanth control group. Curcuminoids did not produce any significant dose-related changes of haematological parameters. In the group of male animals receiving 250 mg/kg bw/day of curcuminoids, actual and relative liver weights and the level of alkaline phosphatase (ALP) were significantly higher than those of the two controls, but the ALP level was still within a normal range. There appeared to be a higher incidence of mild degree of liver fatty degeneration and adrenocortical fatty degeneration in this group of animals, however the incidence was not significantly different from that of the two controls. The authors concluded that at higher doses, curcumin may affect the function and morphology of the liver in a reversible manner. The EFSA Panel concluded that due to the lack of knowledge on the diferuloylmethane content and nature of the extract tested, the study could not be used to assess the safety of diferuloylmethane (Chavalittumrong et al., 2002).

Mice

20. Sub-chronic oral toxicity of turmeric and ETE was studied in female Swiss mice and Wistar rats fed turmeric powder (0, 1 and 5 %) and ethanolic turmeric extract (0, 0.05 and 0.25 %) through the diet for 14 and/or 90 days.

This is a draft statement for discussion.  
It does not reflect the final views of the Committee and should not be cited.

The curcuminoid content of the ETE was approximately 98%. The administration of a high dose of turmeric (5 %) for longer duration (90 days) showed a significant reduction in body weight gain, alterations in absolute and/ or relative liver weights, and hepatotoxicity i.e. focal necrosis or focal necrosis with regeneration both in mice and rats. In mice, lower doses of turmeric (i.e. 0.2 or 1 % for 14 days) also showed hepatotoxicity and they were found to be more vulnerable to turmeric-induced hepatotoxicity than rats (Deshpande et al., 1998).

21. In a 13 week study, groups of 10 male and 10 female B6C3F1 mice were fed diets containing 0, 1000, 5000, 10000, 25000, or 50000 ppm turmeric oleoresin, which were estimated to deliver average daily doses of 150, 750, 1700, 3850, or 7700 mg/kg bw to males and 200, 1000, 1800, 4700 or 9300 mg/kg bw to females (NTP 1993). The major component of the oleoresin was identified as curcumin (79 to 85 %). The percent composition was monitored periodically at the study laboratory with free-acid titration and high-performance liquid chromatography methods, and no change in composition was observed. There were no deaths attributed to turmeric oleoresin and the final mean body weight gains and final mean body weights of all exposed groups of male and female mice were similar to those of the controls. Feed consumption by exposed male and female mice was similar to that by the controls. Absolute and relative liver weights of male mice that received 5000 ppm and male and female mice that received 10000, 25000 and 50000 ppm were significantly greater than those of the controls. Clinical findings in mice included stained fur, and discoloured faeces and urine. According to NTP, there were no biologically significant differences in hematologic, clinical chemistry, or urinalysis parameters, and there were no chemical related histopathologic lesions. EFSA ANS panel (2010) concluded that the no-effect level with respect to gross and microscopic pathological changes was 7700 and 9280 mg/kg bw/day in males and females, respectively, which were the highest doses tested.

#### Chronic toxicity

This is a draft statement for discussion.  
It does not reflect the final views of the Committee and should not be cited.

#### Two years study in rats

22. Groups of 60 male and 60 female F344/N rats were fed diets containing 2000, 10000, or 50000 ppm turmeric oleoresin (79 to 85 % diferuloylmethane) for 104 (males) or 103 (females) weeks, which were estimated to deliver average daily doses of 80, 460, or 2000 mg/kg bw to males and 90, 440, or 2400 mg/kg bw to females. Nine or 10 rats from each exposure group were evaluated after 15 months. Survival of exposed male and female rats was similar to that of the controls. The final mean body weights of all exposed male rats and female rats receiving 2000 and 10000 ppm were similar to those of the controls. The final mean body weights of male and female rats that received 50000 ppm were slightly lower (up to 10 %) than those of the controls throughout much of the study. Feed consumption by exposed male and female rats was similar to that of controls throughout the study. The absolute and relative liver weights of female rats receiving 10000 or 50000 ppm were significantly greater than those of controls at the 15-month interim evaluation. There were no clinical findings related to toxicity. In male and female rats receiving 50000 ppm the haematocrit values, haemoglobin concentrations and erythrocyte counts at the 15-month interim evaluation were significantly lower than those in the controls. In addition, platelet counts in male and female rats that received 50000 ppm and reticulocyte counts in male rats that received 50000 ppm were significantly higher than those in the controls. No biologically significant differences were observed in clinical chemistry parameters. (NTP 1993).

#### Two years study in mice

23. Groups of 60 male and 60 female B6C3F1 mice were fed diets containing 2000, 10000, or 50000 ppm turmeric oleoresin (79 to 85 % diferuloylmethane) for 103 weeks, which were estimated to deliver average daily doses of 220, 520, or 6000 mg/kg bw to males and 320, 1620, or 8400 mg/kg bw to females (NTP 1993). Nine or ten mice from each exposure group were evaluated after 15 months. Survival of exposed male and female mice was similar to that of the controls (male: 0 ppm, 43/50; 2000 ppm, 43/50;

This is a draft statement for discussion.

It does not reflect the final views of the Committee and should not be cited.

10000 ppm, 37/50; 50000 ppm 42/50; female: 39/50, 41/50, 34/50, 42/50). The mean body weight of female mice receiving 50000 ppm was slightly lower (up to 12 %) than that of the controls from about week 25. The final mean body weights of males that received 50000 ppm and females that received 10000 and 50000 ppm were significantly lower than those of controls. The final mean body weights of other exposed groups of male and female mice were similar to those of the controls. Feed consumption by exposed male and female mice was similar to that of the controls throughout the study. The absolute and relative liver weights of male and female mice receiving 10000 and 50000 ppm were significantly greater than those of the controls at the 15-month interim evaluation. There were no clinical findings related to toxicity. No biologically significant differences were observed in hematologic parameters. The alkaline phosphatase values of male and female mice that received 10000 and 50000 ppm were significantly higher than those of controls at the 15-month interim evaluation. The incidences of hepatocellular adenoma in male and female mice receiving 10000 ppm (the mid dose group), but not those in mice receiving 2000 or 50000 ppm, were significantly increased (male: 25/50, 28/50, 35/50, 30/50; female: 7/50, 8/50, 19/51, 14/50). The number of male and female mice in the 10000 and 50000 ppm groups with multiple hepatocellular neoplasms compared with controls was of statistical significance. However, JECFA concluded that this effect was not dose-related and that curcumin is not a carcinogen (FAO/WHO 1995). In addition, the EFSA Panel noted that all statistically significant effects noted by the NTP refer to benign neoplastic lesions (adenomas) (EFSA 2010). The EFSA Panel noted that the effects observed were not dose-dependent, were in line with historical control values and were not consistent across sexes and/or species. The Panel noted moreover that hepatocellular tumours occurring in untreated and treated B6C3F1 mice are not relevant for humans. Therefore, the Panel agreed with JECFA that curcumin was not carcinogenic.

### **Curcuminoids and hepatotoxicity: Case studies in humans**

This is a draft statement for discussion.  
It does not reflect the final views of the Committee and should not be cited.

24. In addition to the reported cases in Italy, several cases of liver toxicity associated with curcumin supplements intake have been reported in the literature.

25. In Australia, from Luber et al., (2019) a 52 year old woman presented to her general practitioner with a one-week history of nausea, pruritus, and painless jaundice with associated pale stools and dark urine. This occurred approximately one month following commencement of one tablet per day of Ancient Wisdom Modern Medicine High Potency Turmeric (375 mg curcuminoids and 4 mg black pepper per tablet), along with a flaxseed oil supplement and occasional diclofenac use for arthritic pain. (N.B. this supplement leads to an intake of 204.6 % of the ADI of 3 mg/kg bw/day). She had no prior history of liver disease and had normal liver function tests (LFTs) three months before. Her medical history was notable only for oligoarticular osteoarthritis.

26. Upon admission, all oral medications and supplements were ceased. She was found to have a bilirubin of 162  $\mu\text{mol/L}$  with a hepatocellular profile on liver function tests (ALT 2591 U/L, AST 1770 U/L, ALP 263 U/L, and GGT 370 U/L). With progressive jaundice over the subsequent days she was referred to the emergency department, at which point her bilirubin peaked at 536  $\mu\text{mol/L}$ .

27. Due to lack of significant improvement by day four of admission, a liver biopsy was performed. Histology showed nonspecific inflammatory changes with generally preserved hepatic architecture and no fibrosis. She was discharged day 12 of admission (bilirubin 260  $\mu\text{mol/L}$ , ALT 1232 U/L) with the presumptive diagnosis of diclofenac induced liver injury. By two months after admission her liver function tests had normalised (bilirubin 21  $\mu\text{mol/L}$ , ALT 33 U/L) and she was discharged from the clinic. At this point she recommenced the turmeric supplement as sole therapy for her arthritis. Three weeks later her nausea recurred and repeat liver function tests showed an acute hepatitis (ALT 2093 U/L, AST 1030 U/L, and bilirubin 60  $\mu\text{mol/L}$ ). She was advised to

This is a draft statement for discussion.  
It does not reflect the final views of the Committee and should not be cited.

cease the turmeric supplement, and two months later her liver function tests had again normalised.

28. The turmeric supplements were sent for analysis by a validated liquid chromatography mass spectrometry method. Results were compared to a toxicology library containing approximately 1400 compounds, including medications, illicit drugs, and over-the-counter medicines. A further sample was analysed by inductively coupled plasma mass spectrometry for the presence of trace elements. The turmeric supplement tested negative for drugs, adulterants and toxic heavy metals. (Luber et al. 2019).

29. A 55 year old man was found to have an asymptomatic transaminitis at a routine check-up. His background history included idiopathic thrombocytopenic purpura, hypertension, gout, and osteoarthritis, with regular medications including long-term telmisartan, atenolol, and lercanidipine. He had no known liver history with normal liver function tests one year prior. His only new medication was commencement of a turmeric supplement five months prior to testing. He was referred to a hepatologist and underwent a screen for causes of acute hepatitis. Abdominal ultrasonography showed diffuse steatosis. A drug reaction was suspected, and the turmeric supplement was ceased. Follow-ups occurred over the subsequent four months. Near normalisation of liver function tests occurred by one month (ALT 96 U/L, bilirubin 10 µmol/L) with further improvement by four months after cessation (ALT 46 U/L, bilirubin 11 µmol/L). The turmeric supplement was the presumed cause of the hepatitis. The turmeric supplement was not identified, thus further analysis on it could not be performed and thus dose information is unknown. (Luber et al. 2019).

30. In the US, reported by Suhail et al., (2020) a 61 year old female with polycystic liver disease presented with fatigue, dark urine and polyarthralgias for one week. She denied alcohol use. Physical examination demonstrated right upper quadrant abdominal tenderness. Laboratory findings were notable for aspartate aminotransferase 1553 mg/dL, alanine aminotransferase 2607

This is a draft statement for discussion.

It does not reflect the final views of the Committee and should not be cited.

mg/dL, alkaline phosphatase 246 mg/dL and total bilirubin 1.6 mg/dL with a direct component of 1 mg/dL. Hepatic synthetic function was intact.

Medications included naproxen and ergocalciferol (vitamin D) with no changes four years prior to the onset of transaminitis. Viral infections were ruled out.

Autoimmune work-up yielded positive antinuclear antibody (1:250) with normal anti-smooth muscle antibody and serum IgG levels. Abdominal Doppler was negative for portal or hepatic vein thrombosis. Magnetic resonance cholangiopancreatography did not show any biliary ductal pathology. It was discovered that the patient was taking turmeric supplements for 6 months. A liver biopsy demonstrated pan lobular hepatitis with early parenchymal collapse suggestive of a morphologic counterpart of acute hepatitis and hepatocellular pattern of injury. The patient was thought to have drug-induced liver injury from turmeric pills that were discontinued and she was discharged with prednisone. Her LFTs normalized after 3 weeks, after which, her prednisone was tapered off.

31. Suhail et al. (2019) noted that the temporal association of liver injury in the patient, normalisation of LFTs upon withdrawal and improvement with steroids implicate the turmeric supplement as the likely causative agent of liver injury. The Roussel-Uclaf Causality Assessment Method (RUCAM) scale, which attempts to codify causality of drug toxicity into objective criteria, was eight in the patient indicating a probable adverse drug reaction. A negative serological workup and normalisation of LFTs following the discontinuation of steroids further solidify this conclusion.

32. In another case in the US reported by Lukefahr et al., (2018) a 71 year old woman was found to have an asymptomatic transaminitis at routine check-up. Her medical history included hypothyroidism, Raynaud's syndrome, osteoarthritis, hypertension, dyslipidaemia, irritable bowel syndrome, and diverticulosis, with regular medications including amlodipine, metoprolol, and atenolol. A 'low cost' turmeric supplement (of unknown product identification) was taken according to label recommendations for a period of 8 months prior to transaminitis testing. The patient was referred to a gastroenterologist.



This is a draft statement for discussion.  
It does not reflect the final views of the Committee and should not be cited.

Laboratory and biopsy findings led to a diagnosis of autoimmune hepatitis. Treatment was limited to withdrawing use of turmeric supplements. AST and ALT decreased significantly within 30 days of discontinuation and normalised by 13 months. In this case, the patient (not the clinicians) hypothesised that the turmeric may have been the cause of the elevated liver transaminases and elected to cease its use.

33. Subsequent reassessment of the liver biopsy by Lukefahr et al., (2018) revealed auto-fluorescent inclusions in the pigment-laden histiocytes, with an excitation/emission spectrum consistent with curcumin, or possibly lipofuscin. Histiocyte fluorescence, which was not noted in liver biopsy specimens from patients with unrelated disorders, was quenchable by treatment with Sudan Black B (SBB), as has been reported for lipofuscin. However, because the authors also documented complete SBB quenching of curcumin autofluorescence in fixed cultured cells specifically loaded with curcuminoids, the authors could not ascertain with certainty whether the histiocyte inclusions were composed of lipofuscin, a lysosomal degradation product and/or curcuminoids derived from the turmeric supplement that the patient was still consuming at the time of the biopsy.

34. In September 2018, the Committee on Herbal Medicinal Products (HMPC) of the European Medicines Agency (EMA) considered several clinical studies with turmeric preparations involving patients with digestive disorders. Although a possible effect in improving symptoms was seen, firm conclusions could not be drawn due to limitations in the study design, for example only one study compared turmeric with a control group. However, the HMPC considered the effectiveness of these preparations for relief of mild digestive problems to be plausible and there is evidence that they have been used safely in this way for at least 30 years (including at least 15 years within the European Union). Thus, the HMPC concluded that turmeric preparations (which cover turmeric ground powders and extracts) can be used (only by adults) for relief of mild problems with digestion, such as feelings of fullness, slow digestion and flatulence. The EMA considered that side effects of dry

This is a draft statement for discussion.  
It does not reflect the final views of the Committee and should not be cited.

mouth, flatulence and stomach irritation may occur. If symptoms continue for longer than two weeks or worsen while taking the medicine, a qualified healthcare practitioner should be consulted (EMA (European Medicines Agency), 2018).

## Exposure from curcuminoids in food

35. The relative proportions and total concentration of curcuminoids within turmeric rhizomes vary depending on the variety grown and the conditions of cultivation (Li et al., 2011) (Table 1).

Table 1. Percentage composition of curcuminoids in turmeric powders and oleoresin extracts (adapted from Li et al., 2011).

Curcuminoid	Composition in turmeric powders (mean) (% dry weight)	Composition in turmeric oleoresin extracts (mean $\pm$ s.d.) (% dry weight)
Diferuloylmethane	2.86	19.51 $\pm$ 2.07
Demethoxycurcumin	1.47	8.31 $\pm$ 1.13
Bisdemethoxycurcumin	1.36	6.22 $\pm$ 0.88
Total	5.69	34.04 $\pm$ 4.08

36. EFSA (EFSA Panel on Food Additives and Nutrient Sources added to Food (ANS), 2010) estimated dietary exposure to diferuloylmethane in children and adults using national consumption data with maximum permitted levels (MPLs) specified in Directive 94/36/EC (EC 1994) (tier 2 approach), and maximum reported use levels (tier 3 approach). Estimates of dietary exposure to diferuloylmethane obtained from these approaches are presented in Table 2.

37. For adults (> 18 years old), the EFSA ANS Panel (2010) estimated the exposure based on the UK consumption survey as the UK population is

This is a draft statement for discussion.  
It does not reflect the final views of the Committee and should not be cited.

considered to be one of the highest consumers of soft drinks in Europe and individual food consumption data (UK National Diet and Nutrition Survey (NDNS), 2000-2001) are available (Tennant D., 2007, 2006). For children (1-10 years old), the Panel estimated exposure based on dietary exposure assessments for children in Europe (EXPOCHI) project. The EXPOCHI project details individual food consumption data from eleven European countries (Belgium, France, the Netherlands, Spain, Italy, Finland, Sweden, Czech Republic, Cyprus, Greece and Germany). As the UK is not included in the EXPOCHI consortium, estimates for UK children (1.5 - 4.5 years old) were made by the Panel with the use of individual food consumption data (UK NDNS, 1992-1993) (Tennant D., 2007, 2006).

Table 2: Estimates of dietary exposure to curcumin in the UK adult population and in children from the EXPOCHI study and UK NDNS data.

<b>Maximum permitted level (tier 2):</b>	<b>UK adult exposure (&gt; 18 years old) to curcumin (mg/kg bw/day)</b>	<b>Children (UK &amp; EXPOCHI , 1-10 years old) exposure to curcumin (mg/kg bw/day)</b>
Mean exposure	0.9	0.5-3.8
Exposure 97.5 <sup>th</sup> %ile	3.3	1.2-7.2
<b>Maximum reported use levels (tier 3):</b>	<b>Children (UK &amp; EXPOCHI , 1-10 years old) exposure to curcumin (mg/kg bw/day)</b>	<b>Children (UK &amp; EXPOCHI , 1-10 years old) exposure to curcumin (mg/kg bw/day)</b>
Mean exposure	0.8 (1.0 *)	0.5-3.4 (0.7-3.6 *)
Exposure 97.5 <sup>th</sup> %ile	2.0 (2.6 *)	1.1-7.1 (1.6-7.6 *)

\* Includes dietary exposure to diferuloylmethane from turmeric powder added to food as a spice and curry powder (see Table 3).

38. In tier 2, the main contributor to curcumin exposure from the UK adult diet was non-alcoholic flavoured drinks (46 %). The main contributors to the

This is a draft statement for discussion.  
It does not reflect the final views of the Committee and should not be cited.

estimates of mean curcumin exposure for UK children (and children considered by the EXPOCHI consortium) were non-alcoholic beverages (13-55 %), fine bakery wares (e.g. biscuits, cakes, wafer) (12-43 %), desserts, including flavoured milk products (12-45 %), and sauces, seasonings, pickles, relishes, chutney and piccalilli (11-42 %).

39. In tier 3, the main contributor to curcumin exposure from the UK adult diet was non-alcoholic flavoured drinks (50 %). The main contributors to the estimates of mean curcumin exposure for UK children (and children considered by the EXPOCHI consortium) were fine bakery wares (e.g. biscuits, cakes, wafer) (13-47 %), desserts (including flavoured milk products) (13-52 %), non-alcoholic beverages (15-57 %) and sauces and seasonings (11-45 %).

40. The exposure assessment in tier 3 does not take into account the use of turmeric as a spice in cooking. The addition of turmeric spice in cooking was estimated to contribute to the dietary exposure of diferuloylmethane by the EFSA ANS panel (2010) (Table 3).

Table 3. Estimates of dietary exposure to diferuloylmethane from ingestion of spice added to food and curry powder in adults in children.

<b>Exposure from spice added to food *</b>	<b>Adult (18-64 years old) curcumin exposure (mg/kg bw/day)</b>	<b>Children (5-12 years old) curcumin exposure (mg/kg bw/day)</b>
Mean	0.1 (n = 66)	0.1 (n = 7)
97.5 <sup>th</sup> %ile	0.3 (n = 66)	0.2 (n = 7)
<b>Exposure from curry powder added to food *</b>	<b>Adult (18-64 years old) curcumin exposure (mg/kg bw/day)</b>	<b>Children (5-12 years old) curcumin exposure (mg/kg bw/day)</b>
Mean	0.1 (n = 91)	0.1 (n = 21)
97.5 <sup>th</sup> %ile	0.3 (n = 91)	0.3 (n = 21)

This is a draft statement for discussion.  
It does not reflect the final views of the Committee and should not be cited.

\* The use of turmeric as a spice added to foods and used in home-made recipes was assessed using data from Irish adults (1379 adults, aged 18-64 years) and children (594 children, aged 5-12 years) (Harrington et al., 2001; IUNA (Irish Universities Nutrition Alliance), 2005). The dietary intake of curry powder was also considered, as turmeric powder is a widespread ingredient of it (approximately 30 % depending on the blend).

41. The above information is indicative of the current exposures to curcumin and/or turmeric as part of the diet (food colour and as a spice). However, consumption of raw and powdered turmeric in large quantities to promote wellbeing is becoming increasingly popular. Based on information readily available on the internet it is proposed that to benefit from turmeric's antioxidant effects, one should consume between 500 to 1000 mg of curcuminoids per day. It is suggested that one teaspoon of fresh or powdered turmeric contains 200 mg of curcumin. Consumptions at the proposed levels, could lead to exposures to curcumin above the recommended ADI.

## **Contamination of raw, ground turmeric and curcumin supplements**

### **Heavy metals**

42. From the recent survey undertaken by Fera Science Ltd (Fera Science Ltd, 2022) all samples were analysed for 69 trace elements which included the heavy metals Pb, Hg, As, and Cd.

43. Twenty-nine of the 30 samples tested had heavy metal concentrations of low concern, i.e. below the maximum level (ML) set for supplements by EC 1881/2006 or below the EU ML set for root spices by the amendment EU 2021/1317 (due to the recent date, this amendment to the EU regulation is not in UK legislation). For supplements, the MLs are 3 mg/kg for Pb, 1 mg/kg for Cd and 0.1 mg/kg for Hg. For root spice powders the EU ML is 1.5 mg/kg for Pb. Arsenic does not have a ML set for supplements, but all concentrations, bar two samples) were below the 0.2 mg/kg ML set for white rice by EU 2015/1006.

This is a draft statement for discussion.  
It does not reflect the final views of the Committee and should not be cited.

44. One sample, a turmeric spice powder, contained a Pb concentration approximately 10 times higher than the majority of other samples analysed, at 2.25 mg/kg. This would be over the amended recent EU ML of 1.5 mg/kg for root spice powders. This sample also had the second highest concentration of Chromium (Cr) at 2.11 mg/kg which may indicate potential adulteration with lead chromate. If this sample was taken as a supplement at, for example, a worst case 2.5 teaspoons a day (approximately 7g) the total exposure of lead from this consumption alone would be 0.23 µg/kg bw/day for a 70 kg adult. This is approximately 2 fold lower than the estimated dietary lead exposure of 0.5 µg/kg bw day for the effects of Pb on developmental neurotoxicity, derived from the BMDL<sub>01</sub> of 12 µg/L blood Pb concentration (EFSA Panel on Contaminants in the Food Chain (CONTAM), 2010)

45. When comparing heavy metal results for supplement samples against spice powders and fresh turmeric there were no clear trends or significant differences between the groups when evaluating the results using a students t-test (supplements against powder & fresh samples).

46. On evaluation of the other trace element results from the 30 samples, results that differed by greater than the mean plus 2 x the standard deviation and 5 x the mean of that sample type (i.e. fresh, powder or supplement) are summarised in Table 4. Table 5 summarises the exposure of each of the trace elements if taking the supplement as advised on the label, alongside comparison of a Health Based Guidance Value (HBGV) if applicable. No calculated exposures for any trace elements if taking the supplements as described on their labels would exceed any HBGVs where HBGVs exist.

Table 4. List of other trace elements (i.e. not including Pb, Hg, As or Cd) from turmeric product samples, where the concentration is greater than 5 x the mean concentration and the mean concentrations plus 2 x standard deviations for that product type.

This is a draft statement for discussion.  
It does not reflect the final views of the Committee and should not be cited.

Sample code	Sample type	Element	Concentration (mg/kg)	Mean concentration of product type (mg/kg)	Increased fold change from mean
TU03	Supplement	Titanium	281	29	9.7
TU03	Supplement	Niobium	0.16	0.023	7.0
TU06	Supplement	Molybdenum	1.9	0.21	9.0
TU06	Supplement	Uranium	0.57	0.079	6.6
TU06	Supplement	Tin	0.16	0.025	6.4
TU07	Supplement	Thallium	0.10	0.012	8.3
TU07	Supplement	Caesium	0.09	0.014	6.4
TU07	Supplement	Zinc	38	7.3	5.2
TU07	Supplement	Barium	50	9.8	5.1
TU10	Powder	Tungsten	0.039	0.0072	5.4
TU12	Supplement	Copper	114	8.6	13.3
TU15	Supplement	Yttrium	4.1	0.56	7.3
TU15	Supplement	Antimony	0.24	0.033	7.3
TU15	Supplement	Palladium	0.11	0.019	5.8
TU15	Supplement	Lanthanum	1.3	0.23	5.7
TU15	Supplement	Calcium	323,000	63,913	5.1
TU17	Supplement	Beryllium	0.124	0.016	7.8
TU17	Supplement	Antimony	0.19	0.033	5.8

This is a draft statement for discussion.  
It does not reflect the final views of the Committee and should not be cited.

Table 5. Summary of exposure of each of the trace elements from Table 4 for an adult taking the supplement as advised on the label, alongside comparison of a Health Based Guidance Value (HBGV) if applicable

Element	Concentration (mg/kg)	Advised consumption of supplement (g/day)	Exposure from product* (µg/kg bw/day)	HBGV or other reference value (µg/kg bw day)	HBGV reference
Titanium	281	0.42	1.69	n/a	n/a
Niobium	0.16	0.42	0.0010	n/a	n/a
Molybdenum	1.9	0.35	0.010	26 (TUI)	(Institute of Medicine (US) Panel on Micronutrients, 2001)
Uranium	0.57	0.35	0.0029	0.6 (TDI)	(WHO, 2012)
Tin	0.16	0.35	0.00080	2 (p-TDI)	(FAO/WHO, 2004)
Thallium	0.10	0.92	0.0013	0.01 (p-RFD)	(EPA, 2009)
Caesium	0.09	0.92	0.0012	n/a	n/a
Zinc	38	0.92	0.50	1000 (TDI)	(FAO/WHO, 1982)
Barium	50	0.92	0.66	200 (TDI)	(SCHER, 2012)
Tungsten #	0.039	7	0.0028	n/a	n/a
Copper	114	1	1.6	70 (ADI)	(EFSA, 2022)
Yttrium	4.1	4.7	0.28	145.5 (TDI)	(Kowalczyk et al., 2022)
Antimony	0.24	4.7	0.016	6	(WHO, 2003)
Palladium	0.11	4.7	0.0074	n/a	n/a



This is a draft statement for discussion.  
 It does not reflect the final views of the Committee and should not be cited.

Lanthanum	1.3	4.7	0.087	51.3 (TDI)	(Kowalczyk et al., 2022)
Calcium	323,000	4.7	21,733	35,714 (TUI)**	(SCF, 2003)
Beryllium	0.124	0.3	0.00053	2	(WHO, 2009)
Antimony	0.19	0.3	0.00081	6	(WHO, 2003)

\* Assumes a 70Kg adult consumes the supplement as stated on the label

\*\* Assumes a 70Kg adult to derive the TUI

# Tungsten concentration from a spice powder not a supplement. A conservative estimate of 5g per day if consumed as a supplement has been used for the exposure calculations

TUI = Tolerable Upper Intake

TDI = Tolerable Daily Intake

p-RFD = Provisional Reference Dose

ADI = Acceptable Daily Intake

This is a draft statement for discussion.  
It does not reflect the final views of the Committee and should not be cited.

## List of Abbreviations and Technical terms

ADI	Acceptable Daily Intake
AFSCA	Belgium's Federal Agency for Food Chain Safety
ALP	Alkaline Phosphatase
ALT	Alanine Aminotransferase
AST	Aspartate Aminotransferase
ANS Panel	Scientific Panel on Food Additives and Nutrient Sources Added to Food
As	Arsenic
BDMC	Bisdemethoxycurcumin
BMDL	Benchmark Dose Level
Cd	Cadmium
CONTAM Panel	Panel on Contaminants in the Food Chain
COT	Committee on Toxicity
Cr	Chromium
DMC	Desmethoxycurcumin
EFSA	European Food Safety Authority
ETE	Ethanollic Turmeric Extract
EU	European Union
EXPOCHI	Exposure Assessments For Children In Europe
FAO	Food and Agricultural Organisations of the United Nations
GGT	Gamma-Glutamyl Transferase
HBGVs	Health-Based Guidance Values
Hg	Mercury
HMPC	Committee on Herbal Medicinal Products
IDH	Idiosyncratic Drug Hepatotoxicity
JECFA	Joint FAO-WHO Expert Committee Report on Food Additives
LFTs	Liver Function Tests
mg	Milligrams
mg/kg bw/day	Milligrams per Kilogram Bodyweight per Day
MPL	Maximum Permitted Levels
ML	Maximum Level

This is a draft statement for discussion.  
It does not reflect the final views of the Committee and should not be cited.

NDA Panel	Panel on Nutrition, Novel Foods and Food Allergens
NDNS	National Diet and Nutrition Survey
NOAEL	No-Observed-Adverse-Effect Level
NTP	United States National Toxicology Program
p-RFD	Provisional Reference Dose
Pb	Lead
ppm	Parts Per Million
SCF	Scientific Committee for Food
SD	Standard Deviation
SSB	Sudan Black B
TEO	Turmeric Essential Oil
TDI	Tolerable Daily Intake
TUI	Tolerable Upper Intake
UK	United Kingdom
U/L	Units per Litre
WHO	World Health Organisation

This is a draft statement for discussion.  
It does not reflect the final views of the Committee and should not be cited.

## References

American Academy of Allergy Asthma & Immunology, 2020. Medications and Drug Allergic Reactions. [Medications and Drug Allergic Reactions | AAAAI](#)

Apica, B.S., Lee, W.M., 2014. Drug-Induced Liver Injury, in: McManus, L.M., Mitchell, R.N. (Eds.), Pathobiology of Human Disease. Academic Press, San Diego, pp. 1825–1837. [Drug-Induced Liver Injury - ScienceDirect](#)

Chavalittumrong, P., Chivapat, S., Rattanajarasroj, S., Punyamong, S., Chuthaputti, A., Phisalaphong, C., 2002. Chronic toxicity study of curcuminoids in rats. Songklanakarin J. Sci. Technol. 24, 16.

Deshpande, S.S., Lalitha, V.S., Ingle, A.D., Raste, A.S., Gadre, S.G., Maru, G.B., 1998. Subchronic oral toxicity of turmeric and ethanolic turmeric extract in female mice and rats. Toxicol. Lett. 95, 183–193. [Subchronic oral toxicity of turmeric and ethanolic turmeric extract in female mice and rats - ScienceDirect](#)

EFSA, 2022. Re-evaluation of the existing health-based guidance 44 values for copper and exposure assessment from all sources. [PubCon EFSA-Q-2020-00399 Draft Opinion Copper 23052022 vb \(1\).pdf](#)

EFSA Panel on Contaminants in the Food Chain (CONTAM), 2010. Scientific Opinion on Lead in Food. EFSA J. 8. [Scientific Opinion on Lead in Food - - 2010 - EFSA Journal - Wiley Online Library](#)

EFSA Panel on Food Additives and Nutrient Sources added to Food (ANS), 2010. Scientific Opinion on the re-evaluation of curcumin (E 100) as a food additive. EFSA J. 8. [Scientific Opinion on the re-evaluation of curcumin \(E 100\) as a food additive - - 2010 - EFSA Journal - Wiley Online Library](#)

This is a draft statement for discussion.  
It does not reflect the final views of the Committee and should not be cited.

EMA (European Medicines Agency), 2018. Committee on Herbal Medicinal Products (HMPC) Assessment report on Curcuma longa L., rhizome. [Assessment report on Curcuma longa L., rhizoma \(europa.eu\)](#)

EPA, 2009. Toxicological review of thallium and compounds. [Toxicological Review of Thallium and compounds \(CAS No. 7440-28-0\) \(PDF\) \(epa.gov\)](#)

FAO/WHO, 2004. Evaluation of certain food additives and contaminants. Sixty-first report of the Joint FAO/WHO Expert Committee on Food Additives. WHO Technical Report Series 922. [WHO TRS 922.pdf](#)

FAO/WHO, 1995. Evaluation of certain food additives and naturally occurring toxicants (Forty-fourth report of the Joint FAO/WHO Expert Committee on Food Additives). WHO Technical Report Series, No. 859. [WHO TRS 859.pdf](#)

FAO/WHO, 1982. Evaluation of certain food additives and contaminants, Report of the Joint FAO WHO Expert Committee on Food Additives. World Health Organization, Geneva.

Fera Science Ltd, 2022. Turmeric survey for the FSA (FS430403). [Background to the study | Food Standards Agency](#)

Ferrell, L., 2000. Liver Pathology: Cirrhosis, Hepatitis, and Primary Liver Tumors. Update and Diagnostic Problems. Mod. Pathol. 13, 679–704. [Liver Pathology: Cirrhosis, Hepatitis, and Primary Liver Tumors. Update and Diagnostic Problems | Modern Pathology \(nature.com\)](#)

Harrington, K.E., Robson, P.J., Kiely, M., Livingstone, M.B., Lambe, J., Gibney, M.J., 2001. The North/South Ireland Food Consumption Survey: survey design and methodology. Public Health Nutr. 4, 1037–1042. [The North/South Ireland Food Consumption Survey: survey design and methodology | Public Health Nutrition | Cambridge Core](#)

This is a draft statement for discussion.  
It does not reflect the final views of the Committee and should not be cited.

Institute of Medicine (US) Panel on Micronutrients, 2001. Molybdenum, in: Dietary Reference Intakes for Vitamin A, Vitamin K, Arsenic, Boron, Chromium, Copper, Iodine, Iron, Manganese, Molybdenum, Nickel, Silicon, Vanadium, and Zinc. National Academies Press (US); [Molybdenum - Dietary Reference Intakes for Vitamin A, Vitamin K, Arsenic, Boron, Chromium, Copper, Iodine, Iron, Manganese, Molybdenum, Nickel, Silicon, Vanadium, and Zinc - NCBI Bookshelf \(nih.gov\)](#)

IUNA (Irish Universities Nutrition Alliance), 2005. National Children's Food Survey. Main report. [IUNA | Survey Reports | Dietary Research | Nutritional Research](#)

Kandarkar, S.V., Sawant, S.S., Ingle, A.D., Deshpande, S.S., Maru, G.B., 1998. Subchronic oral hepatotoxicity of turmeric in mice--histopathological and ultrastructural studies. *Indian J. Exp. Biol.* 36, 675–679.

Kaplowitz, N., 2005. Idiosyncratic drug hepatotoxicity. *Nat. Rev. Drug Discov.* 4, 489–499. [Idiosyncratic drug hepatotoxicity | Nature Reviews Drug Discovery](#)

Kowalczyk, E., Givélet, L., Amlund, H., Sloth, J.J., Hansen, M., 2022. Risk assessment of rare earth elements, antimony, barium, boron, lithium, tellurium, thallium and vanadium in teas. *EFSA J.* 20. [Risk assessment of rare earth elements, antimony, barium, boron, lithium, tellurium, thallium and vanadium in teas - Kowalczyk - 2022 - EFSA Journal - Wiley Online Library](#)

Li Shiyou, Yuan Wei, Deng Guangrui, Wang Ping, Yang Peiying,, Aggarwal Bharat, 2011. Chemical Composition and Product Quality Control of Turmeric (*Curcuma longa* L.). *Pharm. Crops* 5, 28–54. [Chemical Composition and Product Quality Control of Turmeric \(Curcuma longa L.\) \(benthamopen.com\)](#)

Liju, V.B., Jeena, K., Kuttan, R., 2013. Acute and subchronic toxicity as well as mutagenic evaluation of essential oil from turmeric (*Curcuma longa* L.).

This is a draft statement for discussion.  
It does not reflect the final views of the Committee and should not be cited.

Food Chem. Toxicol. Int. J. Publ. Br. Ind. Biol. Res. Assoc. 53, 52–61. [Acute and subchronic toxicity as well as mutagenic evaluation of essential oil from turmeric \(Curcuma longa L\) - ScienceDirect](#)

Luber, R.P., Rentsch, C., Lontos, S., Pope, J.D., Aung, A.K., Schneider, H.G., Kemp, W., Roberts, S.K., Majeed, A., 2019. Turmeric Induced Liver Injury: A Report of Two Cases. Case Rep. Hepatol. 2019, 6741213. [Turmeric Induced Liver Injury: A Report of Two Cases \(hindawi.com\)](#)

Lukefahr, A.L., McEvoy, S., Alfafara, C., Funk, J.L., 2018. Drug-induced autoimmune hepatitis associated with turmeric dietary supplement use. BMJ Case Rep. bcr-2018-224611. [Drug-induced autoimmune hepatitis associated with turmeric dietary supplement use | BMJ Case Reports](#)

National Toxicology Program (NTP), 1993. Toxicology and Carcinogenesis Studies of Turmeric Oleoresin (Cas No. 8024-37-1) (Major Component 79%-85% Curcumin, Cas No. 458-37-7) in F344/N Rats and B6C3F1 Mice (Feed Studies). Technical Report Series No. 427, NIH Publication No. 93-3158. U.S. [TR-427: Turmeric Oleoresin \(CASRN 8024-37-1\) \(Major Component 79% Curcumin, CASRN 458-37-7\) in F344/N Rats and B6C3F1Mice \(Feed Studies\) \(nih.gov\)](#)

SCF (Scientific Committee for Food), 2003. Opinion on the Tolerable Upper Intake Level of Calcium. SCF/CS/NUT/UPPLEV/64 Final. [Scientific Opinion on the Tolerable Upper Intake Level of calcium - - 2012 - EFSA Journal - Wiley Online Library](#)

SCF (Scientific Committee for Food), 1975. Reports from the Scientific Committee for Food (1st series), opinion expressed 27 June 1975. [REPORTS OF THE SCIENTIFIC COMMITTEE FOR FOOD : First series \(pitt.edu\)](#)

This is a draft statement for discussion.  
It does not reflect the final views of the Committee and should not be cited.

SCHER (Scientific Committee on Health and Environmental Risks), 2012.,  
2012. Assessment of the tolerable daily intake of barium. [TABLE OF CONTENTS \(europa.eu\)](#)

Stierum, R., Conesa, A., Heijne, W., Ommen, B. van, Junker, K., Scott, M.P., Price, R.J., Meredith, C., Lake, B.G., Groten, J., 2008. Transcriptome analysis provides new insights into liver changes induced in the rat upon dietary administration of the food additives butylated hydroxytoluene, curcumin, propyl gallate and thiabendazole. Food Chem. Toxicol. 46, 2616–2628.  
[Transcriptome analysis provides new insights into liver changes induced in the rat upon dietary administration of the food additives butylated hydroxytoluene, curcumin, propyl gallate and thiabendazole - ScienceDirect](#)

Suhail, F.K., Masood, U., Sharma, A., John, S., Dhamoon, A., 2020. Turmeric supplement induced hepatotoxicity: a rare complication of a poorly regulated substance. Clin. Toxicol. Phila. Pa 58, 216–217.  
<https://doi.org/10.1080/15563650.2019.1632882>

Tennant D., 2007. Screening potential intakes of natural food colours. Report provided for the Natural Food Colours Association (NATCOL).

Tennant D., 2006. Screening of Colour Intakes from Non-Alcoholic Beverages. Report prepared for the Union of European Beverages Associations (UNESDA), December 2006.

WHO, 2003. Antimony in drinking water. Background document for development of WHO Guidelines for Drinking-water Quality.  
[Microsoft Word - Third Edition Antimony.doc \(who.int\)](#)

WHO, 2009. Beryllium in drinking-water. Background document for development of WHO Guidelines for Drinking-water Quality. [Microsoft Word - Fourth Edition Beryllium Final January 2010.doc \(who.int\)](#)



This is a draft statement for discussion.  
It does not reflect the final views of the Committee and should not be cited.

WHO, 2012. Guidelines for drinking water quality, Third Edition. Geneva, Switzerland: WHO. [Guidelines for drinking-water quality, 3rd edition: Volume 1 - Recommendations incorporating the first and second addenda \(who.int\)](#)